

Averages and moving averages

Averages and moving averages

We saw last time that the average value of a function $f(x)$ between times a and b is $\frac{1}{b-a} \int_a^b f(x) dx$.

Averages and moving averages

We saw last time that the average value of a function $f(x)$ between times a and b is $\frac{1}{b-a} \int_a^b f(x) dx$.

Example 1. *Find the average value of the function $f(x) = 2 + x^2$ from $x = 3$ to $x = 4$.*

Averages and moving averages

We saw last time that the average value of a function $f(x)$ between times a and b is $\frac{1}{b-a} \int_a^b f(x) dx$.

Example 1. *Find the average value of the function $f(x) = 2 + x^2$ from $x = 3$ to $x = 4$.*

One of the main uses of averages is as part of a summary of a set of data.

Averages and moving averages

We saw last time that the average value of a function $f(x)$ between times a and b is $\frac{1}{b-a} \int_a^b f(x) dx$.

Example 1. *Find the average value of the function $f(x) = 2 + x^2$ from $x = 3$ to $x = 4$.*

One of the main uses of averages is as part of a summary of a set of data. Moving averages summarize data over periods of time.

Definition 2. *The moving average of a function $f(t)$ with lag-time n -units is the function $\bar{f}(x)$ whose value at time x is the average value of $f(t)$ between times $x - n$ and x .*

Definition 2. *The moving average of a function $f(t)$ with lag-time n -units is the function $\bar{f}(x)$ whose value at time x is the average value of $f(t)$ between times $x - n$ and x .*

Example 3. *Find the four-quarter moving average of the following quarterly sales data.*

1	3	2	4	1.5	3.2	2.8	4.1
1.4	3.6	2.5	5	2	4	3	4.8

Why would you want to do this?

Moving averages are most often used for quickly changing or seasonal data.

Moving averages are most often used for quickly changing or seasonal data. For example, a stock price can change a lot from day-to-day or even hour-to-hour.

Moving averages are most often used for quickly changing or seasonal data. For example, a stock price can change a lot from day-to-day or even hour-to-hour. Most investors hold onto stocks for longer periods, so summaries of changes over weeks or months could be more helpful.

Moving averages are most often used for quickly changing or seasonal data. For example, a stock price can change a lot from day-to-day or even hour-to-hour. Most investors hold onto stocks for longer periods, so summaries of changes over weeks or months could be more helpful. In another example some businesses, like retail, are seasonal. So to get a good comparison of how business is changing, you don't necessarily want to compare from month to month ("gee, another big spike in December followed by a big letdown in January").

Moving averages are most often used for quickly changing or seasonal data. For example, a stock price can change a lot from day-to-day or even hour-to-hour. Most investors hold onto stocks for longer periods, so summaries of changes over weeks or months could be more helpful. In another example some businesses, like retail, are seasonal. So to get a good comparison of how business is changing, you don't necessarily want to compare from month to month ("gee, another big spike in December followed by a big letdown in January"). You want to either compare this December with last

December or - and this is part of what moving averages are about - compare the last twelve months of sales *on average* to the previous twelve months of sales.

December or - and this is part of what moving averages are about - compare the last twelve months of sales *on average* to the previous twelve months of sales. Moving averages allow us to have the best of both worlds - they can be used for month-to-month comparisons, but by taking an average over a whole period, they aren't sensitive to seasonality or large variability.

December or - and this is part of what moving averages are about - compare the last twelve months of sales *on average* to the previous twelve months of sales. Moving averages allow us to have the best of both worlds - they can be used for month-to-month comparisons, but by taking an average over a whole period, they aren't sensitive to seasonality or large variability. You can see these advantages clearly in previous example. Moving averages are used *everywhere* in financial and insurance analysis, in particular so one can understand underlying trends.

Getting back to math, we can find an integral whose value is the average value of $f(t)$ between times $x - n$ and x .

Getting back to math, we can find an integral whose value is the average value of $f(t)$ between times $x - n$ and x . Namely, we plug into the formula for average value and get $\frac{1}{x - (x - n)} \int_{x-n}^x f(t) dt$, which is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.

Getting back to math, we can find an integral whose value is the average value of $f(t)$ between times $x - n$ and x . Namely, we plug into the formula for average value and get $\frac{1}{x - (x - n)} \int_{x-n}^x f(t) dt$, which is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.

Theorem 4. *The n -unit moving average of a function $f(t)$, namely $\bar{f}(x)$, is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.*

Getting back to math, we can find an integral whose value is the average value of $f(t)$ between times $x - n$ and x . Namely, we plug into the formula for average value and get $\frac{1}{x - (x - n)} \int_{x-n}^x f(t) dt$, which is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.

Theorem 4. *The n -unit moving average of a function $f(t)$, namely $\bar{f}(x)$, is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.*

Example 5. *Find the one- and four-unit moving averages of the function $f(t) = t^2 - 1$.*

Getting back to math, we can find an integral whose value is the average value of $f(t)$ between times $x - n$ and x . Namely, we plug into the formula for average value and get $\frac{1}{x - (x - n)} \int_{x-n}^x f(t) dt$, which is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.

Theorem 4. *The n -unit moving average of a function $f(t)$, namely $\bar{f}(x)$, is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.*

Example 5. *Find the one- and four-unit moving averages of the function $f(t) = t^2 - 1$. Graph all three functions and explain the behavior we see.*

Getting back to math, we can find an integral whose value is the average value of $f(t)$ between times $x - n$ and x . Namely, we plug into the formula for average value and get $\frac{1}{x - (x - n)} \int_{x-n}^x f(t) dt$, which is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.

Theorem 4. *The n -unit moving average of a function $f(t)$, namely $\bar{f}(x)$, is equal to $\frac{1}{n} \int_{x-n}^x f(t) dt$.*

Example 5. *Find the one- and four-unit moving averages of the function $f(t) = t^2 - 1$. Graph all three functions and explain the behavior we see.*

Present and future values of a revenue stream

Present and future values of a revenue stream a first example

Present and future values of a revenue stream a first example

This application of the integral is important in finance,

Present and future values of a revenue stream a first example

This application of the integral is important in finance, and according to professors of finance, it is more important to understand that a (Riemann) sum leads to an integral than to memorize the relevant integrals.

Present and future values of a revenue stream a first example

This application of the integral is important in finance, and according to professors of finance, it is more important to understand that a (Riemann) sum leads to an integral than to memorize the relevant integrals. Understanding the technique for getting to the right answer always trumps memorizing a formula,

Present and future values of a revenue stream a first example

This application of the integral is important in finance, and according to professors of finance, it is more important to understand that a (Riemann) sum leads to an integral than to memorize the relevant integrals. Understanding the technique for getting to the right answer always trumps memorizing a formula, since technique can be applied to new problems more often.

Present and future values of a revenue stream a first example

This application of the integral is important in finance, and according to professors of finance, it is more important to understand that a (Riemann) sum leads to an integral than to memorize the relevant integrals. Understanding the technique for getting to the right answer always trumps memorizing a formula, since technique can be applied to new problems more often.

As we saw last term, the value of investments over time

is governed by exponential functions;

is governed by exponential functions; P dollars today becomes Pe^{rt} dollars in t years if the rate of return is (constant at) r .

is governed by exponential functions; P dollars today becomes Pe^{rt} dollars in t years if the rate of return is (constant at) r . We are also familiar with the need to adjust economic statistics for “today’s dollars”.

is governed by exponential functions; P dollars today becomes Pe^{rt} dollars in t years if the rate of return is (constant at) r . We are also familiar with the need to adjust economic statistics for “today’s dollars”.

Definition 6. *The future value of P dollars today after t years, assuming an investment and/or inflation rate of r over that time is Pe^{rt} .*

is governed by exponential functions; P dollars today becomes Pe^{rt} dollars in t years if the rate of return is (constant at) r . We are also familiar with the need to adjust economic statistics for “today’s dollars”.

Definition 6. *The future value of P dollars today after t years, assuming an investment and/or inflation rate of r over that time is Pe^{rt} . The present value of P dollars t years from now is Pe^{-rt} .*

is governed by exponential functions; P dollars today becomes Pe^{rt} dollars in t years if the rate of return is (constant at) r . We are also familiar with the need to adjust economic statistics for “today’s dollars”.

Definition 6. *The future value of P dollars today after t years, assuming an investment and/or inflation rate of r over that time is Pe^{rt} . The present value of P dollars t years from now is Pe^{-rt} .*

Current value is used to determine the worth of a promise of money in the future.

Example 7. *Compute the current value of \$10000, paid five years from now, assuming an inflation rate of 4%.*

Example 7. *Compute the current value of \$10000, paid five years from now, assuming an inflation rate of 4%.*

Often in business and economics, money is paid (back) in installments,

Example 7. *Compute the current value of \$10000, paid five years from now, assuming an inflation rate of 4%.*

Often in business and economics, money is paid (back) in installments, as with student loans, mortgages and retirement accounts.

Example 7. *Compute the current value of \$10000, paid five years from now, assuming an inflation rate of 4%.*

Often in business and economics, money is paid (back) in installments, as with student loans, mortgages and retirement accounts. How can we determine the current or future value of installment payments?

Example 7. *Compute the current value of \$10000, paid five years from now, assuming an inflation rate of 4%.*

Often in business and economics, money is paid (back) in installments, as with student loans, mortgages and retirement accounts. How can we determine the current or future value of installment payments?

Example 8. *\$100 deposited per month in an account earning 5% interest has a present value of $100 + 100e^{-0.05\frac{1}{12}} + \dots + 100e^{-0.05\frac{11}{12}}$ dollars*

Example 7. *Compute the current value of \$10000, paid five years from now, assuming an inflation rate of 4%.*

Often in business and economics, money is paid (back) in installments, as with student loans, mortgages and retirement accounts. How can we determine the current or future value of installment payments?

Example 8. *\$100 deposited per month in an account earning 5% interest has a present value of $100 + 100e^{-0.05\frac{1}{12}} + \dots + 100e^{-0.05\frac{11}{12}}$ dollars or in summation notation $\sum_{i=0}^{11} 100e^{-0.05\frac{i}{12}}$ dollars.*

Digression:

Digression: summation notation is a convenient shorthand for writing sums.

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$,

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month,

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month, \$1200 was deposited as a continuous income stream, evenly over the year?

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month, \$1200 was deposited as a continuous income stream, evenly over the year? Then, over a short fraction of the year

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month, \$1200 was deposited as a continuous income stream, evenly over the year? Then, over a short fraction of the year $\Delta t = \frac{1}{N}$ th of a year,

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month, \$1200 was deposited as a continuous income stream, evenly over the year? Then, over a short fraction of the year $\Delta t = \frac{1}{N}$ th of a year, we would deposit $1200\Delta t$ dollars.

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month, \$1200 was deposited as a continuous income stream, evenly over the year? Then, over a short fraction of the year $\Delta t = \frac{1}{N}$ th of a year, we would deposit $1200\Delta t$ dollars. Its present value would be $(1200\Delta t)e^{-0.5t}$, where t is a time at which this money is deposited.

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month, \$1200 was deposited as a continuous income stream, evenly over the year? Then, over a short fraction of the year $\Delta t = \frac{1}{N}$ th of a year, we would deposit $1200\Delta t$ dollars. Its present value would be $(1200\Delta t)e^{-0.5t}$, where t is a time at which this money is deposited. The sum of all of these pieces is $\sum_{t_i=\frac{1}{N}}^1 (1200\Delta t)e^{-0.5t_i}$.

Digression: summation notation is a convenient shorthand for writing sums. For example, instead of $(3x_1 + 5) + (3x_2 + 5) + \cdots + (3x_{23} + 5)$, we write $\sum_{i=1}^{23} (3x_i + 5)$.

What if, instead of deposited as \$100 per month, \$1200 was deposited as a continuous income stream, evenly over the year? Then, over a short fraction of the year $\Delta t = \frac{1}{N}$ th of a year, we would deposit $1200\Delta t$ dollars. Its present value would be $(1200\Delta t)e^{-0.5t}$, where t is a time at which this money is deposited. The sum of all of these pieces is $\sum_{t_i=\frac{1}{N}}^1 (1200\Delta t)e^{-0.5t_i}$.

We have gone through the steps of setting up a Riemann sum, leading to an integral;

We have gone through the steps of setting up a Riemann sum, leading to an integral; by passing to a limit and using the definite integral, we see that the present value will be $\int_0^1 1200e^{0.5t}$.

We have gone through the steps of setting up a Riemann sum, leading to an integral; by passing to a limit and using the definite integral, we see that the present value will be $\int_0^1 1200e^{0.5t}$. (Note how much this income stream is worth

We have gone through the steps of setting up a Riemann sum, leading to an integral; by passing to a limit and using the definite integral, we see that the present value will be $\int_0^1 1200e^{0.5t}$. (Note how much this income stream is worth - making payments more frequently can make a big difference!)

We have gone through the steps of setting up a Riemann sum, leading to an integral; by passing to a limit and using the definite integral, we see that the present value will be $\int_0^1 1200e^{0.5t}$. (Note how much this income stream is worth - making payments more frequently can make a big difference!)