Fourth Generation

Shufang Su • U. of Arizona

In collaboration with J. Alwall, J.L. Feng, J. Kumar
arXiv: 1002.3366; 1103.xxxx.

Not the usual 4th generation ...





S. Su

Direct searches





$m_{b'}$ > 372 GeV

$m_{t'}$ > 335 GeV

Daniel Whiteson, LHC West Coast Meeting @ Irvine

LHC

<u>3/pb (ICHEP2010)</u>

Just saw top. ~No sensitivity to t′,b′

<u>40/pb (Moriond2011)</u>

t′→ Wq → l+jets ~350 GeV

t′→ Wq → dileptons ~300 GeV

b′→ Wt → ss dileptons ~350 GeV

<u>500/pb+ (LP2011?)</u>

Up to 500 GeV+ ?

b′ and t′





Flacco, DW, Bar-Shalom & Tait
arvix: 1005.1077, PRL 2010

Exotic 4th Generation Mirror Quarks

S. Su

Exotic 4th Generation Mirror Quarks

chiral under SM gauge group

S. Su

Exotic 4th Generation Mirror Quarks

chiral under SM gauge group

same charge
opposite chirality

S. Su

Exotic 4th Generation Mirror Quarks

chiral under SM gauge group

charge under
hidden symmetry

same charge
opposite chirality

S. Su

New after LHC West Coast @ Irvine

- B'B' study @ Tevatron and LHC
- T'T' results from CDF

S. Su

Outline

- Motivation: general / specific
- WIMPless model
- Constraints
- Simulations and cut analysis
- Exclusion and discovery reach
- Conclusion

Dark matter motivation: general

Dark Matter: long-lived on cosmological time scale

Charge under a new unbroken symmetry ⇒ absolutely stable

- have only gravitational interaction with the SM

  can not be discovered at colliders

- couple to SM through connector Y

YY production with $y \to f X$



Dark matter motivation: general

Dark Matter: long-lived on cosmological time scale

Charge under a new unbroken symmetry ⇒ absolutely stable

- have only gravitational interaction with the SM

  can not be discovered at colliders

- couple to SM through connector Y

YY production with $y \to f X$



Dark matter motivation: general

Dark Matter: long-lived on cosmological time scale

Charge under a new unbroken symmetry ⇒ absolutely stable

- have only gravitational interaction with the SM

  can not be discovered at colliders

- couple to SM through connector Y

YY production with $y \to f X$



SM charge & dark charge

SUSY	neutralino	squark	quark	R-parity
ExD	KK gauge boson	KK quark	quark	KK-parity
Our study	DM(no SM charge)	exotic quark	quark	dark charge

WIMP miracle

WIMP

- $m_{WIMP} \sim m_{weak}$
- $g \sim g_{weak}$

$$\left.\begin{aligned} \Omega h^2 &\sim \frac{2.6 \times 10^{-10}\mathrm{GeV}^{-2}}{\langle \sigma_A v \rangle} \\ \langle \sigma_A v \rangle &\sim \frac{\alpha^2}{m_{weak}^2} 0.1 \sim 10^{-9}\mathrm{GeV}^{-2} \end{aligned}\right\} \Rightarrow \Omega h^2 \sim 0.3$$

naturally around the observed value

WIMP miracle

WIMP

- $m_{WIMP} \sim m_{weak}$
- $g \sim g_{weak}$

$$\left.\begin{aligned} \Omega h^2 &\sim \frac{2.6 \times 10^{-10}\mathrm{GeV}^{-2}}{\langle \sigma_A v \rangle} \\ \langle \sigma_A v \rangle &\sim \frac{\alpha^2}{m_{weak}^2} 0.1 \sim 10^{-9}\mathrm{GeV}^{-2} \end{aligned}\right\} \Rightarrow \Omega h^2 \sim 0.3$$

naturally around the observed value

WIMPless miracle

$$\Omega_X \propto \frac{1}{\langle \sigma v \rangle} \sim \frac{m_X^2}{g_X^4}$$

- only fixes one combination of dark matter mass and coupling
- $m_X/g_X^2 \sim m_{weak}/g_{weak}^2$, $\Omega h^2 \sim 0.3$

could have $m_X \neq m_{weak}$ as long as the relation holds

WIMPless DM

J.L. Feng and J. Kumar, PRL 101, 231301 (2008)
J.L. Feng, H. Tu and H. Yu, JCAP 0810, 043 (2008)
J.L. Feng, M. Kaplinghat, H. Tu and H. Yu, JCAP 0907, 004 (2009)

- dark matter: no SM gauge interactions, not WIMP
- naturally obtain right relic density: similar to WIMP

WIMPless model

J.L. Feng and J. Kumar, PRL 101, 231301 (2008)

- Dark matter is hidden no SM interactions
- DM sector has its own particle content, mass m_X, coupling g_X
- Connected to SUSY breaking sector



$$\rightarrow \frac{m_X}{g_X^2} \sim \frac{m}{g^2} \sim \frac{F}{16\pi^2 M} \rightarrow \Omega_X \propto \frac{1}{\langle \sigma v \rangle} \sim \frac{m_X^2}{g_X^4}$$

right relic density !
(irrespective of its mass)

If no direct coupling to SM:

- interact only through gravity
- impact on structure formation
- no direct/indirect/collider signals

WIMPless: not hidden



WIMPless: not hidden



$m_Y \sim \max(m_{weak}, m_X)$ interaction λXYf

WIMPless: not hidden



$m_Y \sim \max(m_{weak}, m_X)$ interaction λXYf

- indirect detection
  $XX \to ff, YY$
- direct detection
  $Xf \to Xf$
- collider: 4th generation fermions

WIMPless: not hidden



$m_Y \sim \max(m_{weak}, m_X)$ interaction λXYf

- indirect detection
  $XX \to ff, YY$
- direct detection
  $Xf \to Xf$
- collider: 4th generation fermions

open new possibility for
- DM model parameters
- new experimental search windows

WIMPless: not hidden



$m_Y \sim \max(m_{weak}, m_X)$ interaction λXYf

- indirect detection
  $XX \to ff, YY$
- direct detection
  $Xf \to Xf$
- collider: 4th generation fermions

light dark matter

open new possibility for
- DM model parameters
- new experimental search windows

XENON100

XENON100 (11.17 live days)

XENON 100: 1005.2615

Reply to the Comments on the XENON100 First Dark Matter Results.



Heated debate goes on and on and on ...

Collar and Mckinsey: 1005.3723 *Response to arXiv:1005.2615*

Dark matter motivation: specific

light DM with large σ_{SI}

- **not generic in typical WIMP**

  σ_{SI} : **chirality flip, proportional to Yukawa coupling**

  A. Bottino, F. Donato, N. Fornengo and S. Scopel, PRD 68, 043506 (2003); PRD 77, 015002 (2008); PRD 78, 083520 (2008).

- **can be easily accommodated in WIMPless model with connector Y**

  J.L. Feng, J. Kumar and L.E. Strigari, PLB 670, 37 (2008)

Explaining DAMA: WIMPless

WIMPless model (discrete symmetry)

$$V = \lambda \left[X \bar{Q}'_L q_L + X \bar{B}'_R b_R + X \bar{T}'_R t_R \right]$$

$$\begin{array}{lcl} Q'_L & : & \left(3, 2, \frac{1}{6}\right) \\ T'_R & : & \left(3, 1, \frac{2}{3}\right) \\ B'_R & : & \left(3, 1, -\frac{1}{3}\right) \ . \end{array}$$

scattering: Xq → Q' → Xq, q=b,t, induce coupling to gluon at 1-loop

M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Phys. Lett. B 78, 443 (1978).



Explaining DAMA: WIMPless

WIMPless model (discrete symmetry)

$$V = \lambda \left[X \bar{Q}'_L q_L + X \bar{B}'_R b_R + X \bar{T}'_R t_R \right]$$

not chirality
opposite chirality
of SM quark

$$\begin{aligned} Q'_L &: \left(3, 2, \tfrac{1}{6}\right) \\ T'_R &: \left(3, 1, \tfrac{2}{3}\right) \\ B'_R &: \left(3, 1, -\tfrac{1}{3}\right) . \end{aligned}$$

scattering: Xq → Q' → Xq, q=b,t, induce coupling to gluon at 1-loop

M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Phys. Lett. B 78, 443 (1978).



Explaining DAMA: WIMPless

WIMPless model (discrete symmetry)

$$V = \lambda \left[X \bar{Q}'_L q_L + X \bar{B}'_R b_R + X \bar{T}'_R t_R \right]$$

not chirality
opposite chirality
of SM quark

$$\begin{aligned} Q'_L &: \left(3, 2, \tfrac{1}{6}\right) \\ T'_R &: \left(3, 1, \tfrac{2}{3}\right) \\ B'_R &: \left(3, 1, -\tfrac{1}{3}\right) . \end{aligned}$$

Exotic 4th-generation
mirror quarks

scattering: Xq → Q' → Xq, q=b,t, induce coupling to gluon at 1-loop

M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Phys. Lett. B 78, 443 (1978).



Explaining DAMA: WIMPless

WIMPless model

m_X ~ 1-10 GeV, $m_{Q'}$~300-500 GeV, λ ~ 0.3-1, σ_{SI} in right range

J.L. Feng, J. Kumar and L.E. Strigari, PLB 670, 37 (2008)



3rd generation vs. the first two

- first two generations, tree level scattering $\Rightarrow \lambda \sim 0.03$



- third generation
  - loop level scattering, $\lambda \sim 0.3$-1, more natural
  - less constrained by FCNC

Collider Signature: exotic quarks

Y particle appears as exotic 4th generation mirror quarks Q'



Collider Signal

T'T'→ttXX, B'B' →bbXX

- differ from SUSY searches: cascade decay
- differ from usual 4th generation quark T' → Wb, B' → Wt
- appears in a general set of new physics scenarios
  - asymmetric dark matter — B. Dutta and J. Kumar, arXiv: 1012.1341
  - little Higgs with T-parity — H.C. Cheng and I. Low, JHEP 0408, 061 (2004)
  - baryon and lepton number as gauge symmetry — P. Fileviez Perez and M. B. Wise, arXiv: 1002.1754
  - ...

Constraints

- perturbativity constraints: $m_{Q'} = y_{Q'} v$, $m_{Q'} \leq 600$ GeV
- precision electroweak data: $|m_{T'}-m_{B'}| \sim 50$ GeV
- direct searches limits

B'B'→bbXX, similar to sbottom pair production with $\tilde{b} \to b\tilde{\chi}_1^0$





V. M. Abazov et al. [D0 Collaboration], 1005.2222

Constraints

- perturbativity constraints: $m_{Q'} = y_{Q'}\, v$, $m_{Q'} \leq 600$ GeV
- precision electroweak data: $|m_{T'}-m_{B'}| \sim 50$ GeV
- direct searches limits

B'B'→bbXX, similar to sbottom pair production with $\tilde{b} \to b\tilde{\chi}_1^0$



m_{sb} > 247 GeV ⇒ $m_{B'}$ > 365 (440) GeV

V. M. Abazov et al. [D0 Collaboration], 1005.2222

Constraints

B'B'→bbXX, similar to sbottom pair production with $\tilde{b} \to b\tilde{\chi}_1^0$



m_{sb}>230 GeV (95% C.L.)

T. Aaltonen et al. [CDF Collaboration], 1005.2600

Constraints: direct search

CDF, Run II, 2.5 fb^{-1}, gluino pair production, $\tilde{g} \to b\tilde{b} \quad \tilde{b} \to b\tilde{\chi}_1^0$

two or more jets, large MET, 2b-tagging

T. Aaltonen et al. [CDF Collaboration], PRL 102, 221801 (2009).





S. Su

Constraints: direct search

CDF, Run II, 2.5 fb^{-1}, gluino pair production, $\tilde{g} \to b\tilde{b} \quad \tilde{b} \to b\tilde{\chi}_1^0$

two or more jets, large MET, 2b-tagging

T. Aaltonen et al. [CDF Collaboration], PRL 102, 221801 (2009).





m_{gluino} > 340 GeV for Δm ~ 20 GeV $\Rightarrow m_{B'}$ > 370 GeV

S. Su

Constraints: direct search

CDF, Run II, 2.7 fb^{-1}, stop pair production, $\tilde{t}_1 \to b\tilde{\chi}_1^{\pm} \to b\tilde{\chi}_1^0 l\nu$

m_{st} > 150 - 185 GeV, weaker than sbottom limit





A. G. Ivanov [CDF Collaboration], arXiv:0811.0788 [hep-ex].

Simulation

MadGraph - Pythia - PGS

Signal: $T'\bar{T}' \to t^{(*)}X\bar{t}^{(*)}X \to bW^+X\bar{b}W^-X$

- hadronic channel: large cross section
  - SM backgrounds, tt, W, have MET with lepton
  - irreducible background: Z→ νν + jets
- semi-leptonic channel: isolated lepton, suppress QCD background
- purely leptonic channel: suppressed cross section

Similar analyses in the literature

- semileptonic mode, high mass, large luminosity

  T. Han, R. Mahbubani, D. G. E. Walker and L. T. E. Wang, JHEP 0905, 117 (2009)

- hadronic mode, spin and mass determination

  P. Meade and M. Reece, Phys. Rev. D 74, 015010 (2006).

Simulation

MadGraph - Pythia - PGS

Signal: $T'\bar{T}' \to t^{(*)}X\bar{t}^{(*)}X \to bW^+X\bar{b}W^-X$

- hadronic channel: large cross section
  - SM backgrounds, tt, W, have MET with lepton
  - irreducible background: Z→ νν + jets
- semi-leptonic channel: isolated lepton, suppress QCD background
- purely leptonic channel: suppressed cross section

Similar analyses in the literature

- semileptonic mode, high mass, large luminosity

  T. Han, R. Mahbubani, D. G. E. Walker and L. T. E. Wang, JHEP 0905, 117 (2009)

- hadronic mode, spin and mass determination

  P. Meade and M. Reece, Phys. Rev. D 74, 015010 (2006).

Simulation

MadGraph - Pythia - PGS

Signal: $T'\bar{T}' \to t^{(*)}X\bar{t}^{(*)}X \to bW^+X\bar{b}W^-X$

- hadronic channel: large cross section
  - SM backgrounds, tt, W, have MET with lepton
  - irreducible background: Z→ νν + jets
- semi-leptonic channel: isolated lepton, suppress QCD background
- purely leptonic channel: suppressed cross section

Similar analyses in the literature

- semileptonic mode, high mass, large luminosity

  T. Han, R. Mahbubani, D. G. E. Walker and L. T. E. Wang, JHEP 0905, 117 (2009)

- hadronic mode, spin and mass determination

  P. Meade and M. Reece, Phys. Rev. D 74, 015010 (2006).

Semileptonic channel: precuts

Signal: T'T' → ttXX → bbjj l + MET

Precuts

- one isolated electron or muon
- large MET
- large m_T^W

$$m_T^W \equiv m_T(p_T^l, \not{p}_T)$$
$$= \sqrt{2|p_T^l||\not{p}_T|\cos(\Delta\phi(p_T^l, \not{p}_T))}$$





S. Su

Semileptonic channel: precuts

Signal: T'T' → ttXX → bbjj l + MET

Precuts

- one isolated electron or muon
- large MET
- large m_T^W

$$m_T^W \equiv m_T(p_T^l, \not{p}_T)$$
$$= \sqrt{2|p_T^l||\not{p}_T|\cos(\Delta\phi(p_T^l, \not{p}_T))}$$



SM bg peak around m_W



S. Su

Semileptonic channel: precuts

- $N_{jet} \geq 4$



- second, hadronically decay W



Semileptonic channel: Tevatron

Additional cuts: MET, m_T^W, H_T

Tevatron: semileptonic

Cut	T' (300)	T' (400)	T' (500)	$t\bar{t}$	W+jets
No cut	203.2	16.33	1.11	5619	(5179)
1 μ/e, no τ	36.1	2.88	0.194	1041	(2060)
$\not{E}_T > 100$ GeV	17.7	2.00	0.157	107.2	(728.8)
$m_T^W > 100$ GeV	10.7	1.38	0.114	22.6	(36.62)
≥ 4 jets	4.81	0.64	0.062	2.6	0.30
$\lvert m_{jj} - m_W \rvert < 10$ GeV	4.13	0.51	0.049	2.2	0.19
All precuts	4.13	0.51	0.049	2.19	0.19
$m_T^W > 150$ GeV	1.93	0.325	0.036	0.62	0.035
$\not{E}_T > 150$ GeV	1.75	0.367	0.041	0.281	0.035
$H_T > 300$ GeV	1.93	0.353	0.042	1.18	0.07
$\not{E}_T > 150$, $H_T > 300$	1.04	0.279	0.037	0.056	0.017

2.4 fb

Semileptonic channel: LHC10

Additional cuts: MET, m_T^W, H_T

LHC10: semileptonic

Cut	T' (300)	T' (400)	T' (500)	$t\bar{t}$ (1 e/μ)	$t\bar{t}$ (1 τ)	$t\bar{t}$ (2 e/μ)	W+jets
No cut	14.89	3.16	0.922	66.67	43.96	10.62	(42.28)
1 μ/e, no τ	3.2	0.669	0.193	36.45	8.15	3.18	(15.74)
$\not{E}_T > 100$ GeV	1.92	0.52	0.165	5.05	2.07	0.888	(10.33)
$m_T^W > 100$ GeV	1.1	0.342	0.116	0.134	0.638	0.471	(0.235)
≥ 4 jets	0.357	0.116	0.043	0.056	0.091	0.062	0.028
$\lvert m_{jj} - m_W \rvert < 10$ GeV	0.165	0.049	0.016	0.026	0.03	0.014	0.01
All precuts	0.165	0.049	0.016	0.027	0.031	0.014	0.01

82 fb

- Additional m_T^W cut: $m_T^W > 150, 200$ GeV
- Additional $\not{E}_T$ cuts: $\not{E}_T > 150,\ 200,\ 250$ GeV.
- $H_T = \sum_{i=1}^{4} |p_T^j|_i + |p_T^l|$ cuts: $H_T > 400,\ 500$ GeV.
- Combinations of the cuts above.

Hadronic channel: precuts

Signal: T'T' → ttXX → bbjjjj + MET

Background:

- leptonic W decay with lepton missed
- tau lepton mistagged as jets
- Z→νν + jets

Precuts

- No isolated electron, muon, tau-tagged jets
- large MET
- $N_{jet} \geq 5$





Hadronic channel: precuts

- $\Delta\phi(\not{p}_T, p_T^j)$ suppress QCD bg





S. Su

Hadronic channel: precuts

- $\Delta\phi(\not{p}_T, p_T^j)$ suppress QCD bg

tau background: important for new physics with jets and MET.

after Precuts



after Precuts



S. Su

Hadronic channel: Tevatron

Additional cuts: MET, H_T , N_{jet}

Tevatron: hadronic

Cut	T' (300)	T' (400)	T' (500)	$t\bar{t}$	W+jets	Z+jets
No cut	203.24	16.33	1.11	5619.1	(5179.06)	(3030.09)
0 isolated leptons	82.88	6.97	0.499	2265.54	(1756.96)	(2545.12)
$\not{E}_T > 100$ GeV	42.86	5.28	0.422	125.93	(663.5)	(1219.22)
≥ 5 jets	22.64	3.07	0.273	22.11	3.3	2.6
$\Delta\phi$ cuts	19.0	2.74	0.245	15.8	2.8	2.2
All precuts	19	2.74	0.245	15.8	2.8	2.2

21 fb

- Additional $\not{E}_T$ cuts: $\not{E}_T > 150, 200, 250$ GeV.
- $H_T = \sum_{i=1}^{5} |p_T^j|_i$ cuts: $H_T > 300, 350, 400$ GeV.
- At least 6 jets with $|p_T^j| > 20$ GeV.
- Combinations of the cuts above.

Hadronic channel: LHC10

Additional cuts: MET, H_T , N_{jet}

LHC10: hadronic

Cut	T' (300)	T' (400)	T' (500)	$t\bar{t}$ (1 τ)	$t\bar{t}$ (1 e/μ)	$t\bar{t}$ (had)	W+jets	Z+jets
No cut	14.89	3.16	0.922	43.96	66.67	104.59	(42.28)	(18.86)
0 isolated leptons	6.75	1.5	0.45	16.88	13.11	72.29	(16.8)	(15.71)
$\not{E}_T > 100$ GeV	4.15	1.21	0.394	3.91	2.67	0.097	(11.25)	(11.48)
≥ 5 jets	1.34	0.406	0.135	0.664	0.47	0.031	0.305	0.212
$\Delta\phi$ cuts	1.19	0.374	0.125	0.56	0.41	0.01	0.265	0.187
All precuts	1.19	0.374	0.125	0.56	0.41	0.01	0.265	0.187

1.4 pb

- Additional $\not{E}_T$ cuts: $\not{E}_T > 150, 200, 250, 300$ GeV.
- $H_T = \sum_{i=1}^{5} |p_T^j|_i$ cuts: $H_T > 400, 500$ GeV.
- At least 6 jets with $|p_T^j| > 40$ GeV.
- Combinations of the cuts above.

Tevatron exclusion

- optimal cuts (after precuts)
- S/B > 0.1, more than 2 events
- Poisson statistics





Tevatron exclusion

- optimal cuts (after precuts)
- S/B > 0.1, more than 2 events
- Poisson statistics

soft decay products





Tevatron Searches (CDF)

Search for Production of Heavy Particles Decaying to Top Quarks and Invisible Particles in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

T. Aaltonen et al. [CDF Collaboration], arXiV:1103.2482



LHC 10 exclusion





LHC 10 exclusion

off-shell top





Tevatron discovery





LHC10 discovery





LHC @7TeV, multiply lumonisity by a factor of 3.

Tevatron Search

Signal: B'B' →bbXX **Bg: W(lν)jj, Z(νν) jj,WZ, ttbar**

Precuts

- no lepton
- 2 or 3 jets, with $|\eta_j|<2.5$, $P_{Tj} > 20$ GeV
- $\alpha_{j1j2} < 164^o$
- MET ≥ 40 GeV, MET/GeV > 80 - 40X$\Delta\phi_{min}$ (MET, jets)
- ≥ 2 b-jets, including leading jet
- $\Delta\phi$(MET, jets) > 0.6, A=(MET-MHT)/(MET+MHT), -0.1 < A < 0.2
- $X_{jj} = (P_{Tj1}+P_{Tj2})/HT$, $X_{jj} > 0.75$
- $H_T > 60$ GeV

Additional cuts

- P_{Tj1}, MET, X_{jj}, HT

Tevatron Reach

Preliminery results for B'B' →bbXX

- optimal cuts (after precuts)
- S/B > 0.1, more than 2 events
- Poisson statistics





S. Su

LHC7 Search

Signal: B'B' →bbXX **Bg: W(lv)jj, Z(vv) jj,WZ, ttbar**

Precuts

- no lepton
- 2 or 3 jets, with $|\eta_j|<2.5$, P_{Tj} > 100, 40 (40) GeV
- MET ≥ 80 GeV
- $f=MET/M_{eff}$, f>0.3 for 2 jets, f>0.25 for 3 jets
- $\Delta\phi$(MET, jets) > 0.2
- Sphericity S_T>0.2
- ≥ 1 b-jets, including leading jet

Additional cuts

- P_{Tj1}, MET, M_{eff}, additional btag

LHC Reach

Preliminery results for B'B' →bbXX

- optimal cuts (after precuts)
- S/B > 0.1, more than 2 events
- Poisson statistics





Conclusions

- Pair production of exotic quarks $\rightarrow$ DM + SM particles

  DM motivated, WIMPless scenario

  - natural obtain the right relic density
  - explain DAMA results: light DM, large σ_{SI}
- $T'\bar{T}' \rightarrow t\bar{t}XX$ semileptonic mode and hadronic mode
- Current CDF exclusion (4.8 fb^{-1}): 360 GeV
- Exclusion: LHC @ 10 TeV, 300 pb^{-1}
- Discovery:

  m_X<130 GeV, $m_{T'}$ < 405 GeV for Tevatron 20 fb^{-1}

  m_X<170 GeV, $m_{T'}$ < 490 GeV for LHC10 300 pb^{-1}

Conclusions

- B'B' →bbXX
  - current exclusion (5.2 fb^{-1}, D0): 440 GeV
  - Tevatron 20 fb^{-1}, exclusion, 470 GeV
  - LHC @ 7 TeV, 1 fb^{-1}, discovery: 545 GeV
- identify signal as $T'\bar{T}' \to t\bar{t}XX$,comparing with SUSY $\tilde{t}\tilde{t}^* \to t\bar{t}\tilde{\chi}_1^0\tilde{\chi}_1^0$
  - Need ILC ...
- complementary between collider studies and DM searches
  - small λ, DM searches unsuccessful
  - displaced vertex at collider