

A BASIS THEOREM FOR THE DEGENERATE AFFINE ORIENTED BRAUER-CLIFFORD SUPERCATEGORY

JONATHAN BRUNDAN, JONATHAN COMES, AND JONATHAN R. KUJAWA

ABSTRACT. We introduce the oriented Brauer-Clifford and degenerate affine oriented Brauer-Clifford supercategories. These are diagrammatically defined monoidal supercategories which provide combinatorial models for certain natural monoidal supercategories of supermodules and endosuperfunctors, respectively, for the Lie superalgebras of type Q . Our main results are basis theorems for these diagram supercategories. We also discuss connections and applications to the representation theory of the Lie superalgebra of type Q .

1. INTRODUCTION

1.1. **Overview.** Let \mathbb{k} denote a fixed ground field¹ of characteristic not two. In this paper we study certain monoidal supercategories; that is, categories in which morphisms form \mathbb{Z}_2 -graded \mathbb{k} -vector spaces, the category has a tensor product, and compositions and tensor products of morphisms are related by a graded version of the interchange law (see Section 2 for more details). While enriched monoidal categories have been the object of study for some time, it is only recently that monoidal supercategories have taken on a newfound importance thanks to the role they play in higher representation theory. To name a few examples, they appear explicitly or implicitly in the categorification of Heisenberg algebras [RS], “odd” categorifications of Kac-Moody (super)algebras (e.g. [EL, KKO1, KKO2]), the definition of super Kac-Moody categories [BE1], and in various Schur-Weyl dualities in the \mathbb{Z}_2 -graded setting (e.g. [KT]).

In this paper we introduce two monoidal supercategories. They are the *oriented Brauer-Clifford supercategory* \mathcal{OBC} and the *degenerate affine oriented Brauer-Clifford supercategory* \mathcal{AOBC} . They are defined by generators and relations. For both monoidal supercategories the generating objects are \uparrow and \downarrow . Hence, objects in both \mathcal{OBC} and \mathcal{AOBC} can be viewed as finite words in \uparrow and \downarrow (we write $\mathbb{1}$ for the unit object). For \mathcal{OBC} the generating morphisms are the three even morphisms $\smile : \mathbb{1} \rightarrow \uparrow\downarrow$, $\frown : \downarrow\uparrow \rightarrow \mathbb{1}$, $\bowtie : \uparrow\uparrow \rightarrow \uparrow\uparrow$, and one odd morphism $\updownarrow : \uparrow \rightarrow \uparrow$. For \mathcal{AOBC} the generating morphisms are the those of \mathcal{OBC} along with an even morphism $\uparrow : \uparrow \rightarrow \uparrow$. These generating morphisms are subject to an explicit list of local relations (see Definitions 3.2 and 3.7 for details). In Section 3 we explain how more complicated diagrams can be interpreted as morphisms in \mathcal{OBC} and \mathcal{AOBC} . For example, here are two diagrams which correspond to morphisms in

Date: April 19, 2018.


Research of the first author was partially supported by NSF grant DMS-1700905.

Research of the third author was partially supported by NSA grant H98230-11-1-0127.

2010 *Mathematics Subject Classification:* 17B10, 18D10.

¹More generally, one could take \mathbb{k} to be a commutative ring. See Section 6.2.

$\text{Hom}_{\mathcal{AOBC}}(\downarrow^2\uparrow^2, \downarrow^2\uparrow^3\downarrow)$:



In particular, it will be obvious that the Hom-spaces in \mathcal{OBC} and \mathcal{AOBC} are spanned by the set of all such diagrams they contain. As is often the case, the difficulty is in identifying a subset of these diagrams which form a basis. The main results of this paper are contained in Theorems 3.4 and 3.9 in which we provide a diagrammatic basis for the morphism spaces of these supercategories, and Theorem 7.1 in which we provide bases for the cyclotomic quotients of \mathcal{AOBC} .

1.2. Motivation and applications. Let us describe the motivation for these supercategories and some consequences of the aforementioned basis theorems. Let

$$\mathfrak{q} = \mathfrak{q}(n) = \left\{ \begin{pmatrix} A & B \\ B & A \end{pmatrix} \mid A, B \text{ are } n \times n \text{ matrices with entries in } \mathbb{k} \right\}.$$

Put a \mathbb{Z}_2 -grading on $\mathfrak{q} = \mathfrak{q}_0 \oplus \mathfrak{q}_1$ by setting \mathfrak{q}_0 (resp. \mathfrak{q}_1) to be the subspace of matrices with $B = 0$ (resp. $A = 0$). Then \mathfrak{q} is the Lie superalgebra of type Q , where the Lie bracket given by the graded version of the commutator bracket. See Section 4.1 for details.

The representations in type Q do not have a classical analogue. Despite the important early work done by Penkov-Serganova and others to obtain character formulas and other information (see [PS, Bru1] and references therein), the representation theory in type Q remain mysterious. For example, only very recently the structure of category \mathcal{O} for \mathfrak{q} became clear thanks to the work of Chen [Che], Cheng-Kwon-Wang [CKW], and Brundan-Davidson [BD2, BD3].

Since the enveloping superalgebra of \mathfrak{q} , $U(\mathfrak{q})$, is a Hopf superalgebra, one can consider the tensor product of \mathfrak{q} -supermodules and the duals of finite-dimensional \mathfrak{q} -supermodules. Let V denote the natural supermodule for \mathfrak{q} ; that is, column vectors of height $2n$ with the action of \mathfrak{q} given by matrix multiplication. Using the Hopf structure we can then form tensor products of V and its dual, V^* . For brevity, let us write $V^\uparrow = V$ and $V^\downarrow = V^*$ and, more generally, given a word \mathbf{a} in \uparrow and \downarrow , let $V^\mathbf{a}$ denote the tensor product of the corresponding supermodules (e.g. $V^{\uparrow\uparrow\downarrow} = V \otimes V \otimes V^*$). The full subcategory of all \mathfrak{q} -supermodules obtained in this way is a natural object of study.

Moreover, the translation superfunctors given by tensoring with V 's and V^* 's are a key tool in much of the progress made in the study of type Q representations. That is, an important role is played by the full subcategory of endosuperfunctors of the form $V^\mathbf{a} \otimes -$ as \mathbf{a} ranges over all finite words in \uparrow and \downarrow . Given the importance of these endosuperfunctors, it is of interest to understand this supercategory.

By design \mathcal{OBC} and \mathcal{AOBC} , respectively, are combinatorial models for these two supercategories. Specifically, there is a monoidal superfunctor

$$\Phi : \mathcal{OBC} \rightarrow \mathfrak{q}\text{-supermodules}$$

given on objects by $\Phi(\mathbf{a}) = V^\mathbf{a}$. When \mathbb{k} is characteristic zero, this superfunctor is full (see Theorem 4.1). That is, there is a surjective superalgebra homomorphism

$$\text{End}_{\mathcal{OBC}}(\mathbf{a}) \rightarrow \text{End}_{\mathfrak{q}(n)}(V^\mathbf{a}) \quad (1.2)$$

which, moreover, is an isomorphism whenever the length of \mathfrak{a} is less than or equal to n (see Remark 4.2).

It follows from our basis theorem that $\text{End}_{\mathcal{OBC}}(\uparrow^r)$ is isomorphic to the (finite) Sergeev superalgebra introduced in [Ser2] (see Corollary 3.5). For arbitrary \mathfrak{a} , $\text{End}_{\mathcal{OBC}}(\mathfrak{a})$ is isomorphic to the walled Brauer-Clifford superalgebra introduced by Jung-Kang [JK] (see Corollary 3.6). The fact that (1.2) is an isomorphism whenever the length of \mathfrak{a} is less than or equal to n recovers [JK, Theorem 3.5]. We should point out the definitions given in [JK] are global in nature. For example, it is not a priori clear their intricate rule defines an associative product. In contrast, our diagrammatic description for these superalgebras involves only local relations and leads to significant simplifications.

Analogously, there is a monoidal superfunctor from \mathcal{AOBC} to the supercategory of endosuperfunctors of \mathfrak{q} -supermodules, $\Psi : \mathcal{AOBC} \rightarrow \mathcal{E}nd(U(\mathfrak{q})\text{-smod})$, given on objects by $\Psi(\mathfrak{a}) = V^{\mathfrak{a}} \otimes -$. When \mathbb{k} is characteristic zero this superfunctor is faithful “asymptotically” in the sense that given any nonzero morphism in \mathcal{AOBC} , its image under Ψ is nonzero as long as n is sufficiently large. Indeed, this observation is key to proving the basis theorems. We reduce to showing that the induced map

$$\text{End}_{\mathcal{AOBC}}(\uparrow^r) \rightarrow \text{End}_{\mathcal{E}nd(U(\mathfrak{q})\text{-smod})}(V^{\otimes r} \otimes -)$$

is injective for n sufficiently large (e.g. $n \geq r$). This in turn is proven by introducing a certain $\mathfrak{q}(n)$ -supermodule M (which we call the generic Verma supermodule) and proving that the induced map of superalgebras

$$\text{End}_{\mathcal{AOBC}}(\uparrow^r) \rightarrow \text{End}_{\mathfrak{q}(n)}(V^{\otimes r} \otimes M)$$

is injective.

As an application, in Section 4.4 we use Ψ to compute a family of central elements in $U(\mathfrak{q})$. By the basis theorem $\text{End}_{\mathcal{AOBC}}(\mathbf{1})$ is known to be a polynomial ring in $\Delta_1, \Delta_3, \Delta_5, \dots$ where Δ_k is defined by (3.28). In Section 4.4 we explicitly compute the central element corresponding to $\Psi(\Delta_k)$ for each k and show that they are essentially the central elements first introduced by Sergeev [Ser1] after the application of the antipode of $U(\mathfrak{q})$ (see Proposition 4.6).

1.3. Cyclotomic quotients. Fix nonnegative integers a, b and $m_i \in \mathbb{k}$ for each $1 \leq i \leq a$. Using this data fix the polynomial of degree $\ell := 2a + b$ given by $f(t) = t^b \prod_{1 \leq i \leq a} (t^2 - m_i)$. The *cyclotomic quotient* \mathcal{OBC}^f is the supercategory defined as the quotient of \mathcal{AOBC} by the left tensor ideal generated by $f(\uparrow)$. Note that \mathcal{OBC}^f does not obviously inherit the structure of a monoidal supercategory from \mathcal{AOBC} , but it is a left module supercategory over \mathcal{AOBC} . In Section 7 we describe a basis for the Hom-spaces of \mathcal{OBC}^f (see Theorem 7.1). That this forms a basis was conjectured in a preprint version of this article (written by the second two authors). The first author provided a proof soon after the preprint appeared on the ArXiv; see Section 8. Subsequently, Gao-Rui-Son-Su posted an independent proof of Theorem 7.1 in [GRSS2]; their proof is in the spirit of our proof of Theorem 3.9.

1.4. Connection to the superalgebras of Gao-Rui-Song-Su. Shortly after the authors released the first version of this paper, Gao-Rui-Song-Su posted a preprint to the ArXiv in which they describe their independent work on affine walled Brauer-Clifford superalgebras and their cyclotomic quotients [GRSS1]. In the same spirit as this paper, they define these superalgebras by generators and

relations and provide basis theorems. The key difference is that we choose to work with the *supercategories*, whereas they work with *superalgebras*. In Section 7.2 we explain the connection between these two approaches and show their superalgebras are specializations of endomorphism superalgebras in our supercategories. To do so, we construct explicit superalgebra maps from the Gao-Rui-Song-Su superalgebras to the endomorphism superalgebras of our supercategories. Using our bases theorems, one can check that the images under those superalgebra maps of the so-called regular monomials in the superalgebras of Gao-Rui-Song-Su are bases of the appropriate endomorphism superalgebras. Whence, our basis theorems imply the linear independence of the corresponding basis theorems of Gao-Rui-Song-Su ([GRSS1, Theorem 5.15 and Theorem 6.10]). These arguments cannot merely be reversed in order to obtain our basis theorems from those in [GRSS1]. The main obstacle comes from the fact that our endomorphism superalgebras do not come equipped with nice descriptions via generators and relations. Indeed, without already having a basis theorem for the supercategory, it is difficult to extract a full system of generators and relations for the endomorphism superalgebras from the defining generators and relations for the monoidal supercategory. Hence, providing a well-defined superalgebra map from our endomorphism superalgebras to the superalgebras of Gao-Rui-Song-Su is not an easy task. In particular, we are unable to conclude Theorem 6.3 from [GRSS1, Theorem 5.14] nor Theorem 7.1 from [GRSS1, Theorem 6.10].

1.5. Future directions. There are a number of interesting questions yet to be considered. For example, in his PhD thesis, Reynolds showed that the locally finite-dimensional representations of a certain specialization of the oriented Brauer category provide a categorification of the tensor product of a highest weight representation and lowest weight representation for a Kac-Moody Lie algebra of type A [Rey]. See [Bru2] for the quantum version of this story. We expect similar results to hold for \mathcal{OBC} where the categorification is of a representation for a Kac-Moody algebra of type B. Another natural question is to extend the results of this paper from the classical to the quantum setting. There is a quantized enveloping superalgebra, $U_q(\mathfrak{q})$, which is a Hopf superalgebra and one can ask for quantum analogues of \mathcal{OBC} and \mathcal{AOBC} . The quantum walled Brauer-Clifford superalgebras were already introduced in [BGH⁺]. In a third direction, there should be representations of \mathcal{AOBC} and its cyclotomic quotients related to the representations of finite W -superalgebras of type Q and to an expected higher level mixed Schur-Weyl-Sergeev duality (see [BCNR, Section 4.6] where this is explained for type A). So far as the authors are aware, this theory has yet to be developed.

1.6. Acknowledgements. The authors are pleased to thank Nicholas Davidson and Catharina Stroppel for helpful conversations. Part of this project was completed while the second author enjoyed a visit to the Max Planck Institute in Bonn. He would like to thank the institute for providing an excellent working environment.

2. MONOIDAL SUPERCATEGORIES

In this section we give a brief introduction to monoidal supercategories following [BE2, §1]. We refer the reader to *loc. cit.* for more details and further references.

2.1. Superspaces. Let \mathbb{k} be a fixed ground field of characteristic not two. A *superspace* $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is a \mathbb{Z}_2 -graded \mathbb{k} -vector space. As we will also have \mathbb{Z} -gradings we reserve the word degree for later use and instead refer to the *parity* of an element. That is, elements of $V_{\bar{0}}$ (resp. $V_{\bar{1}}$) are said to have parity $\bar{0}$ or to be *even* (resp. parity $\bar{1}$ or *odd*). Given a homogeneous element $v \in V$ we write $|v| \in \mathbb{Z}_2$ for the parity of the element. Given two superspaces V and W , the set of all linear maps $\text{Hom}_{\mathbb{k}}(V, W)$ is naturally \mathbb{Z}_2 -graded by declaring that $f : V \rightarrow W$ has parity $\varepsilon \in \mathbb{Z}_2$ if $f(V_{\varepsilon'}) \subseteq W_{\varepsilon+\varepsilon'}$ for all $\varepsilon' \in \mathbb{Z}_2$. Let \mathbf{svect} and $\underline{\mathbf{svect}}$ denote the categories of all superspaces with $\text{Hom}_{\mathbf{svect}}(V, W) = \text{Hom}_{\mathbb{k}}(V, W)$ and $\text{Hom}_{\underline{\mathbf{svect}}}(V, W) = \text{Hom}_{\mathbb{k}}(V, W)_{\bar{0}}$.

Given superspaces V and W , the tensor product $V \otimes W$ as vector spaces is also naturally a superspace with \mathbb{Z}_2 -grading given by declaring $|v \otimes w| = |v| + |w|$ for all homogeneous $v \in V$ and $w \in W$. The tensor product of linear maps between superspaces is defined via $(f \otimes g)(v \otimes w) = (-1)^{|g||v|} f(v) \otimes g(w)$. This gives $\underline{\mathbf{svect}}$ (but *not* \mathbf{svect}) the structure of a monoidal category with $\mathbf{1} = \mathbb{k}$ (viewed as superspace concentrated in even parity). The graded flip map $v \otimes w \mapsto (-1)^{|v||w|} w \otimes v$ gives $\underline{\mathbf{svect}}$ the structure of a symmetric monoidal category. Here and elsewhere we write the formula only for homogeneous elements with the general case given by extending linearly.

2.2. Monoidal supercategories. By a *supercategory* we mean a category enriched in $\underline{\mathbf{svect}}$. Similarly, a *superfunctor* is a functor enriched in $\underline{\mathbf{svect}}$. Given two superfunctors $F, G : \mathcal{A} \rightarrow \mathcal{B}$, a *supernatural transformation* $\eta : F \rightarrow G$ consists of $\eta_{\mathbf{a}, \varepsilon} \in \text{Hom}_{\mathcal{B}}(F\mathbf{a}, G\mathbf{a})_{\varepsilon}$ for each object $\mathbf{a} \in \mathcal{A}$ and $\varepsilon \in \mathbb{Z}_2$ such that $\eta_{\mathbf{b}, \varepsilon} \circ Ff = (-1)^{\varepsilon|f|} Gf \circ \eta_{\mathbf{a}, \varepsilon}$ for every $f \in \text{Hom}_{\mathcal{A}}(\mathbf{a}, \mathbf{b})$. We will write $\eta_{\mathbf{a}} = \eta_{\mathbf{a}, \bar{0}} + \eta_{\mathbf{a}, \bar{1}} \in \text{Hom}_{\mathcal{B}}(F\mathbf{a}, G\mathbf{a})$. The space of all supernatural transformations from F to G is given the structure of a superspace by declaring a supernatural transformation η to be *even* (resp. *odd*) if $\eta_{\mathbf{a}, \bar{1}} = 0$ (resp. $\eta_{\mathbf{a}, \bar{0}} = 0$) for all objects \mathbf{a} .

Given two supercategories \mathcal{A} and \mathcal{B} , there is a supercategory $\mathcal{A} \boxtimes \mathcal{B}$ whose objects are pairs (\mathbf{a}, \mathbf{b}) of objects $\mathbf{a} \in \mathcal{A}$ and $\mathbf{b} \in \mathcal{B}$ and whose morphisms are given by the tensor product of superspaces $\text{Hom}_{\mathcal{A} \boxtimes \mathcal{B}}((\mathbf{a}, \mathbf{b}), (\mathbf{a}', \mathbf{b}')) = \text{Hom}_{\mathcal{A}}(\mathbf{a}, \mathbf{a}') \otimes \text{Hom}_{\mathcal{B}}(\mathbf{b}, \mathbf{b}')$ with composition defined using the symmetric braiding on $\underline{\mathbf{svect}}$: $(f \otimes g) \circ (h \otimes k) = (-1)^{|g||h|} (f \circ h) \otimes (g \circ k)$. This can be used to give the category \mathbf{scat} of all supercategories the structure of a monoidal category.

By a *monoidal supercategory* we mean a supercategory \mathcal{A} equipped with a superfunctor $- \otimes - : \mathcal{A} \boxtimes \mathcal{A} \rightarrow \mathcal{A}$, a unit object $\mathbf{1}$, and even supernatural isomorphisms $(- \otimes -) \otimes - \xrightarrow{\sim} - \otimes (- \otimes -)$ and $\mathbf{1} \otimes - \xrightarrow{\sim} - \xleftarrow{\sim} - \otimes \mathbf{1}$ called *coherence maps* satisfying certain axioms analogous to the ones for a monoidal category. A monoidal supercategory is called *strict* if its coherence maps are identities. A *monoidal superfunctor* between two monoidal supercategories \mathcal{A} and \mathcal{B} is a superfunctor $F : \mathcal{A} \rightarrow \mathcal{B}$ equipped with an even supernatural isomorphism $(F-) \otimes (F-) \xrightarrow{\sim} F(- \otimes -)$ and an even isomorphism $\mathbf{1}_{\mathcal{B}} \xrightarrow{\sim} F\mathbf{1}_{\mathcal{A}}$ satisfying axioms analogous to the ones for a monoidal functor.

A *braided monoidal supercategory* is a monoidal supercategory \mathcal{A} equipped with a $\underline{\mathbf{svect}}$ -enriched version of a braiding. More precisely, let $T : \mathcal{A} \boxtimes \mathcal{A} \rightarrow \mathcal{A}$ denote the superfunctor defined on objects by $(\mathbf{a}, \mathbf{b}) \mapsto \mathbf{b} \otimes \mathbf{a}$ and on morphisms by $f \otimes g \mapsto (-1)^{|f||g|} g \otimes f$. A braiding on a \mathcal{A} is a supernatural isomorphism $\gamma : - \otimes - \rightarrow T$ satisfying the usual hexagon axioms. A *symmetric monoidal supercategory* is a braided monoidal supercategory \mathcal{A} with $\gamma_{\mathbf{a}, \mathbf{b}}^{-1} = \gamma_{\mathbf{b}, \mathbf{a}}$ for all objects $\mathbf{a}, \mathbf{b} \in \mathcal{A}$.


Given a monoidal supercategory \mathcal{A} and an object $\mathfrak{a} \in \mathcal{A}$, by a *(left) dual* to \mathfrak{a} we mean an object \mathfrak{a}^* equipped with homogeneous *evaluation* and *coevaluation* morphisms $\text{ev}_{\mathfrak{a}} : \mathfrak{a}^* \otimes \mathfrak{a} \rightarrow \mathbb{1}$ and $\text{coev}_{\mathfrak{a}} : \mathbb{1} \rightarrow \mathfrak{a} \otimes \mathfrak{a}^*$, respectively, in which $\text{ev}_{\mathfrak{a}}$ and $\text{coev}_{\mathfrak{a}}$ have the same parity and satisfy the super version of the usual adjunction axioms. For example, given a finite-dimensional superspace V with homogeneous basis $\{v_i \mid i \in I\}$, then $V^* = \text{Hom}_{\mathbb{k}}(V, \mathbb{k})$ with evaluation and coevaluation given by $f \otimes v \mapsto f(v)$ and $1 \mapsto \sum_{i \in I} v_i \otimes v_i^*$ respectively, where $v_i^* \in V^*$ is defined by $v_i^*(v_j) = \delta_{i,j}$. A monoidal supercategory in which every object has a (left) dual is called *(left) rigid*.

The following examples will be relevant for what follows.

- (i) Any \mathbb{k} -linear monoidal category can be viewed as a monoidal supercategory in which all Hom-spaces are concentrated in parity $\bar{0}$. If the category is braided, symmetric braided, or rigid, then it still is as a supercategory.
- (ii) The tensor product and braiding defined in Section 2.1 give **svect** the structure of a symmetric monoidal supercategory with $\mathbb{1} = \mathbb{k}$ (viewed as a superspace concentrated in parity $\bar{0}$). The symmetric braiding $\gamma_{V,W} : V \otimes W \rightarrow W \otimes V$ is given by the graded flip map. The full subcategory of finite-dimensional superspaces is rigid.
- (iii) Given a Lie superalgebra $\mathfrak{g} = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$ over a field \mathbb{k} of characteristic not two, let \mathfrak{g} -smod denote the supercategory of all \mathfrak{g} -supermodules. That is, superspaces $M = M_{\bar{0}} \oplus M_{\bar{1}}$ with an action by \mathfrak{g} which respects the grading in the sense that $\mathfrak{g}_{\varepsilon} \cdot M_{\varepsilon'} \subseteq M_{\varepsilon + \varepsilon'}$. The tensor product $M \otimes M'$ has action given by $x.(m \otimes m') = (x.m) \otimes m' + (-1)^{|x||m|} m \otimes (x.m')$ for all homogeneous $x \in \mathfrak{g}$, $m \in M$, and $m' \in M'$ and the graded flip map provides a symmetric braiding. The unit object $\mathbb{1}$ is the ground field \mathbb{k} with trivial \mathfrak{g} -action. In this way \mathfrak{g} -smod is a symmetric monoidal supercategory. The full subcategory of finite-dimensional \mathfrak{g} -supermodules is rigid with the action given on M^* by $(x.f)(m) = -(-1)^{|x||f|} f(x.m)$.
- (iv) Given a supercategory \mathcal{A} let $\mathcal{E}nd(\mathcal{A})$ denote the supercategory whose objects are all superfunctors $\mathcal{A} \rightarrow \mathcal{A}$ with supernatural transformations as morphisms. We give $\mathcal{E}nd(\mathcal{A})$ the structure of a monoidal supercategory with $\mathbb{1} = \text{Id}$ as follows. The tensor product of two superfunctors $F, G : \mathcal{A} \rightarrow \mathcal{A}$ is defined by composition $F \otimes G = F \circ G$. Given supernatural transformations $\eta : F \rightarrow G$ and $\theta : H \rightarrow K$ we define $\eta \otimes \theta : F \circ H \rightarrow G \circ K$ by setting $(\eta \otimes \theta)_{\mathfrak{a}, \varepsilon} = \sum_{\varepsilon_1 + \varepsilon_2 = \varepsilon} \eta_{K_{\mathfrak{a}, \varepsilon_1}} \circ F\theta_{\mathfrak{a}, \varepsilon_2}$ for each object $\mathfrak{a} \in \mathcal{A}$ and $\varepsilon \in \mathbb{Z}_2$. The coherence maps are the obvious ones.

When working with monoidal supercategories it will sometimes be convenient to use the following notation. Given objects \mathfrak{a} and \mathfrak{b} in a monoidal supercategory, we write $\mathfrak{a}\mathfrak{b} := \mathfrak{a} \otimes \mathfrak{b}$. We will also write $\mathfrak{a}^r := \underbrace{\mathfrak{a} \otimes \cdots \otimes \mathfrak{a}}_{r \text{ times}}$.

2.3. String calculus. There is a well-defined string calculus for strict monoidal supercategories discussed in [BE2, §1.2]. A morphism $f : \mathfrak{a} \rightarrow \mathfrak{b}$ is drawn as



when the objects are left implicit. Notice that the convention used in this paper is to read diagrams from bottom to top. The products of morphisms $f \otimes g$ and $f \circ g$ are given by horizontal and vertical stacking respectively:

$$\begin{array}{c} | \\ \textcircled{f} \\ | \end{array} \otimes \begin{array}{c} | \\ \textcircled{g} \\ | \end{array} = \begin{array}{c} | \\ \textcircled{f} \\ | \end{array} \begin{array}{c} | \\ \textcircled{g} \\ | \end{array}, \quad \begin{array}{c} | \\ \textcircled{f} \\ | \end{array} \circ \begin{array}{c} | \\ \textcircled{g} \\ | \end{array} = \begin{array}{c} | \\ \textcircled{f} \\ | \\ \textcircled{g} \\ | \end{array}.$$

Pictures involving multiple products should be interpreted by *first composing horizontally, then composing vertically*. For example,

$$\begin{array}{c} | \\ \textcircled{f} \\ | \\ | \\ \textcircled{h} \\ | \end{array} \begin{array}{c} | \\ \textcircled{g} \\ | \\ | \\ \textcircled{k} \\ | \end{array}$$

should be interpreted as $(f \otimes g) \circ (h \otimes k)$. In general, this is *not* the same as $(f \circ h) \otimes (g \circ k)$ because of the *super-interchange law*:

$$(f \otimes g) \circ (h \otimes k) = (-1)^{|g||h|} (f \circ h) \otimes (g \circ k).$$

In terms of string calculus, the super-interchange law implies

$$\begin{array}{c} | \\ \textcircled{f} \\ | \\ | \\ \textcircled{g} \\ | \end{array} = \begin{array}{c} | \\ \textcircled{f} \\ | \\ | \\ \textcircled{g} \\ | \end{array} = (-1)^{|f||g|} \begin{array}{c} | \\ \textcircled{g} \\ | \\ | \\ \textcircled{f} \\ | \end{array}. \quad (2.1)$$

2.4. Graded and filtered superspaces. By a *graded superspace* we mean a \mathbb{k} -vector space V equipped with a decomposition $V = \bigoplus_{(i,\varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2} V_{i,\varepsilon}$. We write $V_i = V_{i,\bar{0}} \oplus V_{i,\bar{1}}$ for the elements of V that are *homogeneous of degree i* . Given two graded superspaces V and W , we write $\text{Hom}_{\mathbb{k}}(V, W)_{i,\bar{0}}$ (resp. $\text{Hom}_{\mathbb{k}}(V, W)_{i,\bar{1}}$) for the space of all \mathbb{k} -linear maps that map $V_{j,\varepsilon}$ to $W_{j+i,\varepsilon}$ (resp. $W_{j+i,\varepsilon+\bar{1}}$) for each $(j,\varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2$. We let $\underline{\mathbf{gsvec}}$ and $\underline{\mathbf{gsvec}}$ denote the supercategories of all graded superspaces with

$$\text{Hom}_{\underline{\mathbf{gsvec}}}(V, W) = \bigoplus_{(i,\varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2} \text{Hom}_{\mathbb{k}}(V, W)_{i,\varepsilon}$$

$$\text{and } \text{Hom}_{\underline{\mathbf{gsvec}}}(V, W) = \text{Hom}_{\mathbb{k}}(V, W)_{0,\bar{0}}.$$

There is a natural way to give $\underline{\mathbf{gsvec}}$ (resp. $\underline{\mathbf{gsvec}}$) the structure of a monoidal supercategory (resp. monoidal category) with

$$(V \otimes W)_{i,\varepsilon} = \bigoplus_{(j,\eta) \in \mathbb{Z} \times \mathbb{Z}_2} V_{j,\eta} \otimes W_{i-j,\varepsilon+\eta}.$$

By a *filtered superspace* we mean a superspace $V = V_{\bar{0}} \oplus V_{\bar{1}}$ equipped with a filtration $\cdots \subseteq V_{\leq i,\varepsilon} \subseteq V_{\leq i+1,\varepsilon} \subseteq \cdots$ such that $\bigcap_{(i,\varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2} V_{\leq i,\varepsilon} = 0$ and $\bigcup_{(i,\varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2} V_{\leq i,\varepsilon} = V_{\varepsilon}$ for each $\varepsilon \in \mathbb{Z}_2$. We write $V_{\leq i} = V_{\leq i,\bar{0}} \oplus V_{\leq i,\bar{1}}$ for the elements of V that are *filtered degree i* . Given two filtered superspaces V and W , we write $\text{Hom}_{\mathbb{k}}(V, W)_{\leq i,\bar{0}}$ (resp. $\text{Hom}_{\mathbb{k}}(V, W)_{\leq i,\bar{1}}$) for the space of all \mathbb{k} -linear maps that map $V_{\leq j,\varepsilon}$ to $W_{\leq j+i,\varepsilon}$ (resp. $W_{\leq j+i,\varepsilon+\bar{1}}$) for each $(j,\varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2$. We let $\underline{\mathbf{fsvec}}$ and $\underline{\mathbf{fsvec}}$ denote the supercategories of all filtered superspaces with

$$\text{Hom}_{\underline{\mathbf{fsvec}}}(V, W) = \bigoplus_{(i,\varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2} \text{Hom}_{\mathbb{k}}(V, W)_{\leq i,\varepsilon}$$

and $\mathrm{Hom}_{\mathbf{fsvec}}(V, W) = \mathrm{Hom}_{\mathbb{k}}(V, W)_{\leq 0, \bar{0}}$.

There is a natural way to give \mathbf{fsvec} (resp. \mathbf{fsvec}) the structure of a monoidal supercategory (resp. monoidal category) with

$$(V \otimes W)_{\leq i, \varepsilon} = \bigoplus_{(j, \eta) \in \mathbb{Z} \times \mathbb{Z}_2} V_{\leq j, \eta} \otimes W_{\leq i-j, \varepsilon + \eta}.$$

Every graded superspace V can be viewed as a filtered superspace by setting $V_{\leq i, \varepsilon} = \bigoplus_{j \leq i} V_{j, \varepsilon}$ for all $(i, \varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2$. On the other hand, given a filtered superspace V , we write $\mathrm{gr} V$ for the *associated graded superspace* with $(\mathrm{gr} V)_{i, \varepsilon} := V_{\leq i, \varepsilon} / V_{\leq i-1, \varepsilon}$ for each $(i, \varepsilon) \in \mathbb{Z} \times \mathbb{Z}_2$. Given filtered superspaces V and W , a map $f \in \mathrm{Hom}_{\mathbb{k}}(V, W)_{\leq i, \varepsilon}$ induces a map $\mathrm{gr}_{i, \varepsilon} f \in \mathrm{Hom}_{\mathbb{k}}(\mathrm{gr} V, \mathrm{gr} W)_{i, \varepsilon}$ in an obvious way.

2.5. Graded and filtered supercategories. By a *graded* (resp. *filtered*) *supercategory* we mean a category enriched in \mathbf{gsvec} (resp. \mathbf{fsvec}). Similarly, a *graded* (resp. *filtered*) *superfunctor* is a functor enriched in \mathbf{gsvec} (resp. \mathbf{fsvec}). By a *graded monoidal supercategory* we mean a monoidal supercategory that is graded in such a way that $f \otimes g$ is homogeneous of degree $i + j$ whenever f and g are homogeneous of degree i and j respectively. Similarly, a *filtered monoidal supercategory* is a monoidal supercategory that is filtered in such a way that $f \otimes g$ has filtered degree $i + j$ whenever f and g have filtered degree i and j respectively.

Given a filtered supercategory \mathcal{A} , the *associated graded supercategory* $\mathrm{gr} \mathcal{A}$ is the supercategory with the same objects as \mathcal{A} and with $\mathrm{Hom}_{\mathrm{gr} \mathcal{A}}(\mathbf{a}, \mathbf{b}) := \mathrm{gr} \mathrm{Hom}_{\mathcal{A}}(\mathbf{a}, \mathbf{b})$. The composition in $\mathrm{gr} \mathcal{A}$ is induced from the composition in \mathcal{A} . Similarly, given a filtered superfunctor $F : \mathcal{A} \rightarrow \mathcal{B}$ we write $\mathrm{gr} F : \mathrm{gr} \mathcal{A} \rightarrow \mathrm{gr} \mathcal{B}$ for the associated graded superfunctor defined in the obvious way.


For example, \mathbf{gsvec} and \mathbf{fsvec} are graded and filtered monoidal supercategories respectively. Note that $\mathrm{gr} \mathbf{fsvec}$ and \mathbf{gsvec} are not the same, but there is a faithful superfunctor

$$\Gamma : \mathrm{gr} \mathbf{fsvec} \rightarrow \mathbf{gsvec}$$

which maps a filtered superspace to its associated graded superspace and maps $f + \mathrm{Hom}_{\mathbb{k}}(V, W)_{\leq i-1, \varepsilon} \in \mathrm{Hom}_{\mathbb{k}}(V, W)_{\leq i, \varepsilon} / \mathrm{Hom}_{\mathbb{k}}(V, W)_{\leq i-1, \varepsilon}$ to $\mathrm{gr}_{i, \varepsilon} f$.

3. THE DEGENERATE AFFINE ORIENTED BRAUER-CLIFFORD SUPERCATEGORY

In this section we define the monoidal supercategories \mathcal{OBC} and \mathcal{AOBC} . First, however, we recall the definition of the supercategory \mathcal{OB} from [BCNR]. Our definitions will make use of the string calculus for strict monoidal supercategories (see Section 2.3). In particular, each of the supercategories mentioned above admit a diagrammatic description. The objects in each of these supercategories are \otimes -generated by two objects denoted \uparrow and \downarrow . Hence, the set of all objects can be identified with the set $\langle \uparrow, \downarrow \rangle$ of all finite words in the letters \uparrow and \downarrow . The string diagrams for these supercategories will be made with oriented strings with an upward (resp. downward) string corresponding to the object \uparrow (resp. \downarrow). For example, a diagram of the form



corresponds to a morphism $\uparrow\downarrow\uparrow\rightarrow\uparrow\downarrow\uparrow\downarrow$. We will describe classes of diagrams which give bases for the Hom-spaces of \mathcal{OBC} and \mathcal{AOBC} . In this section we will show these diagrams indeed span the appropriate Hom-spaces. Proof that the diagrams are linearly independent will be given in Section 6.

3.1. The oriented Brauer category. In [BCNR] the oriented Brauer category is defined diagrammatically and then a presentation is given in terms of generators and relations [BCNR, Theorem 1.1]. We take the latter description as our definition and view it as a supercategory concentrated in parity $\bar{0}$.

Definition 3.1. The *oriented Brauer supercategory* \mathcal{OB} is the \mathbb{k} -linear strict monoidal supercategory generated by two objects \uparrow, \downarrow and three even morphisms $\smile : \mathbb{1} \rightarrow \uparrow\downarrow$, $\frown : \downarrow\uparrow \rightarrow \mathbb{1}$, $\bowtie : \uparrow\uparrow \rightarrow \uparrow\uparrow$ subject to the following relations:

$$\begin{array}{c} \bowtie \\ \text{=} \\ \uparrow\uparrow \end{array} = \begin{array}{c} \uparrow\uparrow \\ \text{=} \\ \uparrow\uparrow \end{array}, \quad \begin{array}{c} \bowtie \\ \text{=} \\ \bowtie \end{array}, \quad (3.1)$$

$$\begin{array}{c} \uparrow \\ \text{=} \\ \uparrow \end{array}, \quad \begin{array}{c} \downarrow \\ \text{=} \\ \downarrow \end{array}, \quad \begin{array}{c} \uparrow \\ \text{=} \\ \uparrow \end{array} \text{ is invertible.} \quad (3.2)$$

Note that the last relation is the assertion that there is another distinguished generator

$$\begin{array}{c} \bowtie \\ \text{=} \\ \bowtie \end{array} : \uparrow\downarrow \rightarrow \downarrow\uparrow \text{ which is a two-sided inverse to } \begin{array}{c} \bowtie \\ \text{=} \\ \bowtie \end{array} := \begin{array}{c} \uparrow \\ \text{=} \\ \uparrow \end{array}. \quad (3.3)$$

We define rightward cups/caps and downward crossings in \mathcal{OB} as follows:

$$\begin{array}{c} \smile \\ \text{=} \\ \smile \end{array}, \quad \begin{array}{c} \frown \\ \text{=} \\ \frown \end{array}, \quad \begin{array}{c} \bowtie \\ \text{=} \\ \bowtie \end{array}. \quad (3.4)$$

An *oriented Brauer diagram with bubbles* of type $\mathbf{a} \rightarrow \mathbf{b}$ is any string diagram obtained by stacking (vertically and horizontally) the defining generators of \mathcal{OB} along with the diagrams (3.3) and (3.4) in such a way that the result can be interpreted as a morphism in $\text{Hom}_{\mathcal{OB}}(\mathbf{a}, \mathbf{b})$. For example, here are two oriented Brauer diagrams with bubbles of type $\downarrow\uparrow^4\downarrow^2 \rightarrow \uparrow^3\downarrow^2$:

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \quad (3.5)$$

The term *bubble* refers to any component of such a diagram without an endpoint. In the examples above, the left diagram has two bubbles whereas the right has none. An *oriented Brauer diagram* refers to an oriented Brauer diagram with bubbles that has no bubbles. We say that two oriented Brauer diagrams are *equivalent* if they are of the same type and one diagram can be obtained from the other by continuously deforming its strands, possibly moving them through other strands and crossings, but keeping endpoints fixed. Moreover, we say two oriented Brauer diagrams with bubbles are *equivalent* if they have the same number of bubbles and their underlying oriented Brauer diagrams (without bubbles) are equivalent. For

example, the left (resp. right) diagram in (3.5) is equivalent to the following diagram on the left (resp. right):



Of course, any morphism in \mathcal{OB} can be realized as a \mathbb{k} -linear combination of oriented Brauer diagrams with bubbles. It follows from [BCNR, Theorem 1.1] that two oriented Brauer diagrams with bubbles represent the same morphism in \mathcal{OB} if and only if they are equivalent. Moreover, the set of all equivalence classes of oriented Brauer diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ is a basis for $\text{Hom}_{\mathcal{OB}}(\mathbf{a}, \mathbf{b})$.

As explained in [BCNR, §1], \mathcal{OB} is a rigid symmetric monoidal supercategory. Briefly, the symmetric braiding on \mathcal{OB} is the obvious one given on generating objects by the crossings $\begin{array}{c} \curvearrowright \\ \times \\ \curvearrowleft \end{array}$, $\begin{array}{c} \curvearrowleft \\ \times \\ \curvearrowright \end{array}$, $\begin{array}{c} \curvearrowright \\ \times \\ \curvearrowright \end{array}$, $\begin{array}{c} \curvearrowleft \\ \times \\ \curvearrowleft \end{array}$. Moreover, the generating objects are dual to one another with evaluation and coevaluation maps given by $\begin{array}{c} \curvearrowright \\ \cup \\ \curvearrowright \end{array}$, $\begin{array}{c} \curvearrowleft \\ \cup \\ \curvearrowleft \end{array}$ and $\begin{array}{c} \curvearrowleft \\ \cup \\ \curvearrowleft \end{array}$, $\begin{array}{c} \curvearrowright \\ \cup \\ \curvearrowright \end{array}$ respectively.

3.2. The oriented Brauer-Clifford supercategory. Adjoining ‘‘Clifford generators’’ to \mathcal{OB} results in the following:

Definition 3.2. The *oriented Brauer-Clifford supercategory* \mathcal{OBC} is the \mathbb{k} -linear strict monoidal supercategory generated by two objects \uparrow, \downarrow ; three even morphisms $\begin{array}{c} \curvearrowright \\ \cup \\ \uparrow \end{array} : \mathbf{1} \rightarrow \uparrow\downarrow$, $\begin{array}{c} \curvearrowleft \\ \cup \\ \downarrow \end{array} : \downarrow\uparrow \rightarrow \mathbf{1}$, $\begin{array}{c} \curvearrowright \\ \times \\ \curvearrowright \end{array} : \uparrow\uparrow \rightarrow \uparrow\uparrow$; and one odd morphism $\begin{array}{c} \uparrow \\ \circ \\ \uparrow \end{array} : \uparrow \rightarrow \uparrow$ subject to (3.1) and (3.2), and the following relations:


$$\begin{array}{c} \uparrow \\ \circ \\ \uparrow \end{array} = \uparrow, \quad \begin{array}{c} \curvearrowright \\ \times \\ \curvearrowright \end{array} = \begin{array}{c} \curvearrowright \\ \times \\ \uparrow \end{array}, \quad \begin{array}{c} \circ \\ \cup \\ \downarrow \end{array} = 0. \quad (3.6)$$

Note that the last relation in (3.6) makes use of the rightward cup defined by (3.4).

We define the downward analogue of $\begin{array}{c} \uparrow \\ \circ \\ \uparrow \end{array}$ as follows:

$$\begin{array}{c} \downarrow \\ \circ \\ \downarrow \end{array} := \begin{array}{c} \downarrow \\ \cup \\ \downarrow \end{array} \quad (3.7)$$

There is an obvious monoidal superfunctor $\mathcal{OB} \rightarrow \mathcal{OBC}$ which allows us to view equivalence classes of oriented Brauer diagrams with bubbles as morphisms in \mathcal{OBC} . We define an *oriented Brauer-Clifford diagram* (resp. *with bubbles*) to be an oriented Brauer diagram (resp. with bubbles) with finitely many \circ 's on its segments, where *segment* refers to a connected component of the diagram obtained when all crossings and local extrema are deleted. For example, here are two oriented Brauer-Clifford diagrams of type $\downarrow\uparrow\uparrow\downarrow\downarrow \rightarrow \uparrow\downarrow\downarrow\uparrow\downarrow$:



With (3.7) in mind, we can interpret any oriented Brauer-Clifford diagram with bubbles as a morphism in \mathcal{OBC} .

Since $\uparrow\circ$ is odd, we must keep the super-interchange law in mind when sliding \circ 's along strands. For example, by (2.1) we have

$$\uparrow\circ\downarrow = \uparrow\downarrow\circ = -\uparrow\downarrow\circ.$$

For a more complicated example, in (3.8) numerous signs occur if one tries to rewrite the left-hand diagram into the right-hand diagram. For example, the \circ in the upper left corner of the right-hand diagram could be obtained by sliding the corresponding \circ in the left-hand diagram upward along its strand. In doing so it would have to pass six of the other \circ 's in the left-hand diagram. Hence this move would incur a cost of scaling by $(-1)^6$. However, the following result shows that \circ 's are allowed to freely slide along cups/caps and pass through crossings.

Proposition 3.3. *The following relations hold in \mathcal{OBC} :*

$$\uparrow\circ = \uparrow\circ, \quad \downarrow\circ = \downarrow\circ, \quad (3.9)$$

$$\circ\uparrow = \circ\uparrow, \quad \circ\downarrow = \circ\downarrow, \quad (3.10)$$

$$\uparrow\circ\downarrow = \uparrow\downarrow\circ, \quad \downarrow\circ\uparrow = \downarrow\uparrow\circ, \quad (3.11)$$

$$\uparrow\circ\downarrow = \uparrow\downarrow\circ, \quad \downarrow\circ\uparrow = \downarrow\uparrow\circ, \quad (3.12)$$

$$\uparrow\circ\downarrow = \uparrow\downarrow\circ, \quad \downarrow\circ\uparrow = \downarrow\uparrow\circ, \quad (3.13)$$

$$\uparrow\circ\downarrow = \uparrow\downarrow\circ, \quad \downarrow\circ\uparrow = \downarrow\uparrow\circ. \quad (3.14)$$

Proof. The left of (3.9) is verified below. The right of (3.9) is similar.

$$\uparrow\circ \stackrel{(3.7)}{=} \uparrow\circ = \uparrow\circ \stackrel{(3.2)}{=} \uparrow\circ.$$

Next, we prove (3.11)–(3.14). In (3.11)–(3.14) the right equality is obtained from the left by composing on top and bottom with $\uparrow\circ\downarrow$, $\downarrow\circ\uparrow$, $\uparrow\circ\downarrow$, and $\downarrow\circ\uparrow$, respectively. The left of (3.11) is one of the defining relations of \mathcal{OBC} . The left of (3.12) is verified below:

$$\uparrow\circ\downarrow \stackrel{(3.3)}{=} \uparrow\circ\downarrow \stackrel{(3.9)}{=} \uparrow\circ\downarrow \stackrel{(3.11)}{=} \uparrow\circ\downarrow \stackrel{(3.9)}{=} \uparrow\circ\downarrow \stackrel{(3.3)}{=} \uparrow\circ\downarrow.$$

Similarly, the left of (3.13) (resp. (3.14)) follows from (3.3) (resp. (3.4)) and the left of (3.11) (resp. (3.12)).

Finally, we verify the left of (3.10) below.

$$\circ\uparrow \stackrel{(3.4)}{=} \circ\uparrow \stackrel{(3.12)}{=} \circ\uparrow \stackrel{(3.9)}{=} \circ\uparrow \stackrel{(3.13)}{=} \circ\uparrow \stackrel{(3.4)}{=} \circ\uparrow.$$

The right of (3.10) is similar. □

Next, we will flesh out more of the diagrammatic nature of \mathcal{OBC} . However, before doing so we pause to point out that (3.11)–(3.14) imply the symmetric braiding on

\mathcal{OB} extends to a supernatural isomorphism for \mathcal{OBC} . In particular, \mathcal{OBC} inherits the structure of a rigid symmetric monoidal supercategory from \mathcal{OB} .

We have a downward analogue of the left relation in (3.6):

$$\begin{array}{c} \downarrow \\ \circ \\ \downarrow \\ \circ \end{array} \stackrel{(3.7)}{=} \begin{array}{c} \downarrow \\ \circ \\ \downarrow \\ \circ \end{array} \stackrel{(2.1)}{=} \begin{array}{c} \downarrow \\ \circ \\ \downarrow \\ \circ \end{array} \stackrel{(3.2)}{=} \begin{array}{c} \downarrow \\ \circ \\ \downarrow \\ \circ \end{array} \stackrel{(3.6)}{=} \begin{array}{c} \downarrow \\ \circ \\ \downarrow \\ \circ \end{array} \stackrel{(3.2)}{=} \begin{array}{c} \downarrow \\ \circ \\ \downarrow \\ \circ \end{array}. \quad (3.15)$$

In particular, it follows that every oriented Brauer-Clifford diagram with bubbles can be reduced, up to sign, to a diagram with the same underlying oriented Brauer diagram and at most one \circ on each strand. For example, the two diagrams in (3.8) represent the same morphism in \mathcal{OBC} .

Additionally, we have an even analogue of the right relation in (3.6). Indeed,

$$\begin{array}{c} \circ \\ \circ \end{array} \stackrel{(3.6)}{=} \begin{array}{c} \circ \\ \circ \end{array} \stackrel{(3.9)}{=} \begin{array}{c} \circ \\ \circ \end{array} \stackrel{(2.1)}{=} \begin{array}{c} \circ \\ \circ \end{array} \stackrel{(3.10)}{=} \begin{array}{c} \circ \\ \circ \end{array} \stackrel{(3.6)}{=} \begin{array}{c} \circ \\ \circ \end{array}.$$

Since $\text{char } \mathbb{k} \neq 2$, the computation above implies

$$\begin{array}{c} \circ \\ \circ \end{array} = 0. \quad (3.16)$$

It follows that any oriented Brauer-Clifford diagram with at least one bubble reduces to zero in \mathcal{OBC} . Hence, the Hom-spaces in \mathcal{OBC} are spanned by oriented Brauer-Clifford diagrams (without bubbles) that have at most one \circ on each strand. Next, we refine this spanning set:

We say that an oriented Brauer-Clifford diagram is *normally ordered* if

- ◇ it has at most one \circ on each strand; all \circ 's are on outward-pointing boundary segments (i.e. segments which intersect the boundary at a point that is directed out of the picture);
- ◇ all \circ 's which occur on upwardly oriented segments are positioned at the same height; similarly, all \circ 's which occur on downwardly oriented segments are positioned at the same height.

For example, in (3.8) the diagram on the right is normally ordered and the one on the left is not. We say that two normally ordered oriented Brauer-Clifford diagrams are *equivalent* if their underlying oriented Brauer diagrams are equivalent (see Section 3.1) and their corresponding strands have the same number of \circ 's on them. Note that two equivalent normally ordered oriented Brauer-Clifford diagrams correspond to the same morphism in \mathcal{OBC} . Moreover, it follows from the discussion above that the Hom-spaces in \mathcal{OBC} are spanned by normally ordered oriented Brauer-Clifford diagrams. In fact, we have:

Theorem 3.4. *For any $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$ the superspace $\text{Hom}_{\mathcal{OBC}}(\mathbf{a}, \mathbf{b})$ has basis given by equivalence classes of normally ordered oriented Brauer-Clifford diagrams of type $\mathbf{a} \rightarrow \mathbf{b}$.*

It is possible to give a straightforward combinatorial proof of Theorem 3.4. However, we omit such a proof since Theorem 3.4 is a consequence of our basis theorem for \mathcal{AOBC} as explained in Section 3.4.

Meanwhile, let us point the following consequence of Theorem 3.4. For $r \geq 1$, let C_r denote the Clifford algebra generated by c_1, \dots, c_r , subject to the relations $c_i^2 = 1$ and $c_i c_j = -c_j c_i$ for $i \neq j$. We view C_r as a superalgebra by declaring the generators to be odd. Let $\mathbb{k}\Sigma_r$ denote the group algebra of the symmetric group

on r letters viewed as a superalgebra concentrated in parity $\bar{0}$. Then the (finite) *Sergeev superalgebra* is

$$\text{Ser}_r = C_r \otimes \mathbb{k}\Sigma_r \quad (3.17)$$

as a superspace with $C_r \otimes 1 \cong C_r$ and $1 \otimes \mathbb{k}\Sigma_r \cong \mathbb{k}\Sigma_r$ as subsuperalgebras and with mixed relation $wc_i = c_{w(i)}w$ for $i = 1, \dots, r$ and all $w \in \Sigma_r$.

Corollary 3.5. *For $r \geq 1$,*

$$\text{End}_{\mathcal{OBC}}(\uparrow^r) \cong \text{Ser}_r.$$

Proof. The Sergeev superalgebra is generated by $s_1, \dots, s_{r-1}, c_1, \dots, c_r$ subject to $s_i^2 = 1$, $s_i s_j = s_j s_i$ if $|i - j| > 1$, $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, $c_i^2 = 1$, $c_i c_j = -c_j c_i$ if $i \neq j$, and $s_i c_i = c_{i+1} s_i$ for all admissible $1 \leq i, j \leq r$. From this it follows that there is a well defined superalgebra map

$$\varphi : \text{Ser}_r \rightarrow \text{End}_{\mathcal{OBC}}(\uparrow^r)$$

given by

$$\begin{aligned} \varphi(s_i) &= \begin{array}{c} \uparrow^{r-i-1} \quad \uparrow^{i-1} \\ \quad \quad \quad \times \\ \uparrow^{r-i} \quad \uparrow^{i-1} \end{array}, \\ \varphi(c_i) &= \begin{array}{c} \uparrow^{r-i} \quad \uparrow^{i-1} \\ \quad \quad \quad \circ \\ \uparrow^{r-i} \quad \uparrow^{i-1} \end{array}. \end{aligned}$$

Using Theorem 3.4 it is straightforward to see that φ is an isomorphism. In particular, a direct count shows that there are $2^r(r!)$ normally ordered oriented Brauer diagrams of type $\uparrow^r \rightarrow \uparrow^r$ which matches the dimension of Ser_r . \square

More generally, given nonnegative integers r, s , let $BC_{r,s}$ denote the *walled Brauer-Clifford superalgebra*. This superalgebra was introduced in [JK] where it is denoted $\tilde{B}_{r,s}$ and called the *walled Brauer superalgebra*. Our notation and terminology is chosen so as to be consistent with that of [BGH⁺]. In the next result we assume \mathbb{k} contains $\sqrt{-1}$.

Corollary 3.6. *Let \mathbf{a} be a word consisting of r \uparrow 's and s \downarrow 's. Then*

$$\text{End}_{\mathcal{OBC}}(\mathbf{a}) \cong BC_{r,s}$$

as superalgebras.

Proof. We first note that if \mathbf{a} and \mathbf{b} are two words with r \uparrow 's and s \downarrow 's, then

$$\text{End}_{\mathcal{OBC}}(\mathbf{a}) \cong \text{End}_{\mathcal{OBC}}(\mathbf{b}).$$

The isomorphism is given by applying suitable symmetric braidings for \mathcal{OBC} . Consequently, assume $\mathbf{a} = \downarrow^s \uparrow^r$. By [JK, Theorem 5.1], $BC_{r,s}$ is generated by even generators $s_1, \dots, s_{r-1}, s_{r+1}, \dots, s_{r+s-1}, e_{r,r+1}$ and odd generators c_1, \dots, c_{r+s} subject to an explicit set of relations. By checking relations we see that there is a superalgebra map $\alpha : BC_{r,s} \rightarrow \text{End}_{\mathcal{OBC}}(\downarrow^s \uparrow^r)$ given by

$$\begin{aligned} \alpha(s_i) &= \begin{cases} \begin{array}{c} \downarrow^s \uparrow^{r-i-1} \quad \uparrow^{i-1} \\ \quad \quad \quad \times \\ \downarrow^{r+s-i-1} \quad \uparrow^{i-1} \end{array}, & i = 1, \dots, r-1; \\ \begin{array}{c} \downarrow^s \quad \uparrow^{r-1} \\ \quad \quad \quad \times \\ \downarrow^{r+s-i-1} \quad \uparrow^{i-1} \end{array}, & i = r+1, \dots, r+s-1; \end{cases}, \\ \alpha(e_{r,r+1}) &= \begin{array}{c} \downarrow^{s-1} \quad \uparrow^{r-1} \\ \quad \quad \quad \curvearrowright \\ \downarrow^{s-1} \quad \uparrow^{r-1} \end{array}, \end{aligned}$$

$$\alpha(c_i) = \begin{cases} \sqrt{-1} \begin{array}{c} \downarrow \uparrow \uparrow \downarrow \\ \downarrow \uparrow \downarrow \uparrow \\ \downarrow \uparrow \downarrow \uparrow \end{array}, & i = 1, \dots, r; \\ \sqrt{-1} \begin{array}{c} \downarrow \uparrow \uparrow \downarrow \\ \downarrow \uparrow \downarrow \uparrow \\ \downarrow \uparrow \downarrow \uparrow \end{array}, & i = r + 1, \dots, r + s. \end{cases}$$

As for the previous theorem one can use Theorem 3.4 to verify that α is an isomorphism. We leave the details to the reader. \square

3.3. The supercategory \mathcal{AOBC} . We now introduce an affine version of \mathcal{OBC} .


Definition 3.7. The *degenerate affine oriented Brauer-Clifford supercategory \mathcal{AOBC}* is the \mathbb{k} -linear strict monoidal supercategory generated by two objects \uparrow, \downarrow ; four even morphisms $\curvearrowright : \mathbb{1} \rightarrow \uparrow\downarrow, \curvearrowleft : \downarrow\uparrow \rightarrow \mathbb{1}, \times : \uparrow\uparrow \rightarrow \uparrow\uparrow, \uparrow \circlearrowleft : \uparrow \rightarrow \uparrow$; and one odd morphism $\uparrow \circlearrowright : \uparrow \rightarrow \uparrow$ subject to (3.1), (3.2), (3.6), and the following relations:

$$\begin{array}{c} \uparrow \circlearrowright \\ \bullet \end{array} = - \begin{array}{c} \uparrow \circlearrowleft \\ \bullet \end{array}, \quad \begin{array}{c} \times \\ \bullet \end{array} - \begin{array}{c} \times \\ \bullet \end{array} = \begin{array}{c} \uparrow \\ \bullet \end{array} \begin{array}{c} \uparrow \\ \bullet \end{array} - \begin{array}{c} \uparrow \\ \bullet \end{array} \begin{array}{c} \uparrow \\ \bullet \end{array}. \quad (3.18)$$

We define the downward analogue of $\uparrow \circlearrowright$ as follows:

$$\begin{array}{c} \downarrow \circlearrowright \\ \bullet \end{array} := \begin{array}{c} \downarrow \\ \bullet \end{array} \begin{array}{c} \downarrow \\ \bullet \end{array} \quad (3.19)$$

We define a *dotted oriented Brauer-Clifford diagram (resp. with bubbles)* to be an oriented Brauer diagram (resp. with bubbles) with finitely many \circ 's and \bullet 's on its segments. For example, (1.1) shows two dotted oriented Brauer-Clifford diagrams with bubbles of type $\downarrow^2 \uparrow^2 \rightarrow \downarrow^2 \uparrow^3 \downarrow$. Given a nonnegative integer k , we will draw a \bullet labeled by k to denote k \bullet 's on a strand; that is, the vertical composition of k $\uparrow \circlearrowright$'s or k $\downarrow \circlearrowright$'s. For example, the diagram on the right of (1.1) could have been drawn as



We can interpret any dotted oriented Brauer-Clifford diagram with bubbles as a morphism in \mathcal{AOBC} , and the Hom-spaces in \mathcal{AOBC} are certainly spanned by all dotted oriented Brauer-Clifford diagrams with bubbles. For all $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$ and each $k \in \mathbb{Z}$ we let $\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k, \bar{0}}$ (resp. $\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k, \bar{1}}$) denote the \mathbb{k} -span of all dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ having at most k \bullet 's and an even (resp. odd) number of \circ 's. This gives \mathcal{AOBC} the structure of a filtered monoidal supercategory (see Section 2.5). Given a dotted oriented Brauer-Clifford diagram with bubbles, d , we will write $\text{deg}(d)$ for the number of \bullet 's appearing in d . For example, $\text{deg}(d) = 7$ when d is either of the diagrams in (1.1).

Definition 3.8. A dotted oriented Brauer-Clifford diagram with bubbles is *normally ordered* if

- ◇ removing all bubbles and all \bullet 's results in a normally ordered oriented Brauer-Clifford diagram;
- ◇ each bubble has zero \circ 's and an odd number of \bullet 's, are crossing-free, counterclockwise, and there are no other strands shielding it from the rightmost edge of the picture;

- ◇ each \bullet is either on a bubble or on an inward-pointing boundary segment;
- ◇ whenever a \bullet and a \circ appear on a segment that is both inward and outward-pointing, the \circ appears ahead of the \bullet in the direction of the orientation.

For example, in (1.1) the diagram on the right is normally ordered and the one on the left is not. We say that two normally ordered dotted oriented Brauer-Clifford diagrams with bubbles are *equivalent* if their underlying oriented Brauer diagrams with bubbles are equivalent and their corresponding strands have the same number of \circ 's and \bullet 's. We can now state our main result:

Theorem 3.9. *For any $\mathfrak{a}, \mathfrak{b} \in \langle \uparrow, \downarrow \rangle$ the superspace $\text{Hom}_{\mathcal{AOBC}}(\mathfrak{a}, \mathfrak{b})$ has basis given by equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathfrak{a} \rightarrow \mathfrak{b}$.*

3.4. Consequences of Theorem 3.9 for \mathcal{OBC} . For this subsection we assume Theorem 3.9. Let $\mathcal{OB}(0)$ denote the quotient of \mathcal{OB} obtained by requiring (3.16). Then the Hom-spaces in $\mathcal{OB}(0)$ have a basis given by equivalence classes of oriented Brauer diagrams *without bubbles*. Moreover, there are obvious monoidal superfunctors

$$\mathcal{OB}(0) \rightarrow \mathcal{OBC} \rightarrow \mathcal{AOBC}. \quad (3.20)$$

We saw in Section 3.2 that the Hom-spaces in \mathcal{OBC} are spanned by normally ordered oriented Brauer-Clifford diagrams. It follows from Theorem 3.9 that the image of the equivalence classes of normally ordered oriented Brauer-Clifford diagrams under (3.20) are linearly independent in \mathcal{AOBC} , whence they are linearly independent in \mathcal{OBC} . Theorem 3.4 follows. Moreover, it follows that the monoidal superfunctors in (3.20) are both faithful.

3.5. Normally ordered diagrams span. In Section 3.2 we showed that normally ordered oriented Brauer-Clifford diagrams span the Hom-spaces in \mathcal{OBC} . The goal of this subsection is to show that normally ordered dotted oriented Brauer-Clifford diagrams with bubbles span the Hom-spaces in \mathcal{AOBC} . We start with an analogue of Proposition 3.3 for \mathcal{AOBC} . In the proof, we will make use of Proposition 3.3 without reference.

Proposition 3.10. *The following relations hold in \mathcal{AOBC} :*

$$\begin{array}{c} \uparrow \bullet \curvearrowright = \uparrow \bullet \curvearrowright, \\ \uparrow \bullet \curvearrowright = \uparrow \bullet \curvearrowright, \end{array} \quad \begin{array}{c} \bullet \curvearrowright = \bullet \curvearrowright, \\ \bullet \curvearrowright = \bullet \curvearrowright, \end{array} \quad (3.21)$$

$$\begin{array}{c} \uparrow \bullet \curvearrowright = \uparrow \bullet \curvearrowright, \\ \uparrow \bullet \curvearrowright = \uparrow \bullet \curvearrowright, \end{array} \quad \begin{array}{c} \bullet \curvearrowright = \bullet \curvearrowright, \\ \bullet \curvearrowright = \bullet \curvearrowright, \end{array} \quad (3.22)$$

$$\begin{array}{c} \uparrow \bullet \times = \uparrow \bullet \times + \uparrow \uparrow - \circ \circ, \\ \uparrow \bullet \times = \uparrow \bullet \times + \uparrow \uparrow - \circ \circ, \end{array} \quad \begin{array}{c} \bullet \times = \bullet \times - \uparrow \uparrow - \circ \circ, \\ \bullet \times = \bullet \times - \uparrow \uparrow - \circ \circ, \end{array} \quad (3.23)$$

$$\begin{array}{c} \uparrow \bullet \times = \uparrow \bullet \times + \uparrow \uparrow + \circ \circ, \\ \uparrow \bullet \times = \uparrow \bullet \times + \uparrow \uparrow + \circ \circ, \end{array} \quad \begin{array}{c} \bullet \times = \bullet \times - \uparrow \uparrow - \circ \circ, \\ \bullet \times = \bullet \times - \uparrow \uparrow - \circ \circ, \end{array} \quad (3.24)$$

$$\begin{array}{c} \uparrow \bullet \times = \uparrow \bullet \times - \uparrow \uparrow + \circ \circ, \\ \uparrow \bullet \times = \uparrow \bullet \times - \uparrow \uparrow + \circ \circ, \end{array} \quad \begin{array}{c} \bullet \times = \bullet \times + \uparrow \uparrow - \circ \circ, \\ \bullet \times = \bullet \times + \uparrow \uparrow - \circ \circ, \end{array} \quad (3.25)$$

$$\begin{array}{c} \uparrow \bullet \times = \uparrow \bullet \times + \uparrow \uparrow + \circ \circ, \\ \uparrow \bullet \times = \uparrow \bullet \times + \uparrow \uparrow + \circ \circ, \end{array} \quad \begin{array}{c} \bullet \times = \bullet \times - \uparrow \uparrow + \circ \circ, \\ \bullet \times = \bullet \times - \uparrow \uparrow + \circ \circ. \end{array} \quad (3.26)$$

Proof. The proof of (3.21) is similar to that of (3.9). In (3.23), (3.24), (3.25), and (3.26) the right equality is obtained from the left by composing on top and bottom

with , , , and  respectively. The left of (3.23) is one of the defining relations of \mathcal{AOCB} . The left of (3.24) is verified below:

$$\begin{array}{c}
 \begin{array}{c} \text{crossing with dot on top strand} \\ \text{(3.3)} \\ \text{crossing with dot on bottom strand} \end{array} \stackrel{(3.21)}{=} \begin{array}{c} \text{crossing with dot on left strand} \\ \text{(3.21)} \\ \text{crossing with dot on right strand} \end{array} \stackrel{(3.23)}{=} \begin{array}{c} \text{crossing with dot on top strand} \\ \text{(3.23)} \\ \text{crossing with dot on bottom strand} \end{array} + \begin{array}{c} \text{crossing with dot on left strand} \\ \text{(3.23)} \\ \text{crossing with dot on right strand} \end{array} - \begin{array}{c} \text{crossing with dot on top strand} \\ \text{(3.23)} \\ \text{crossing with dot on bottom strand} \end{array} \\
 \begin{array}{c} \text{crossing with dot on top strand} \\ \text{(3.21)} \\ \text{crossing with dot on bottom strand} \end{array} \stackrel{(2.1)}{=} \begin{array}{c} \text{crossing with dot on left strand} \\ \text{(3.21)} \\ \text{crossing with dot on right strand} \end{array} + \begin{array}{c} \text{crossing with dot on top strand} \\ \text{(3.21)} \\ \text{crossing with dot on bottom strand} \end{array} + \begin{array}{c} \text{crossing with dot on left strand} \\ \text{(3.21)} \\ \text{crossing with dot on right strand} \end{array} \stackrel{(3.3)}{=} \begin{array}{c} \text{crossing with dot on top strand} \\ \text{(3.3)} \\ \text{crossing with dot on bottom strand} \end{array} + \begin{array}{c} \text{crossing with dot on left strand} \\ \text{(3.3)} \\ \text{crossing with dot on right strand} \end{array} + \begin{array}{c} \text{crossing with dot on top strand} \\ \text{(3.3)} \\ \text{crossing with dot on bottom strand} \end{array} .
 \end{array}$$

Proofs for the left equalities in (3.25) and (3.26) are similar. Finally, we verify the right of (3.22) below.

$$\begin{array}{c}
 \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.4)} \\ \text{cup with dot on bottom strand} \end{array} \stackrel{(3.24)}{=} \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.24)} \\ \text{cup with dot on bottom strand} \end{array} - \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.24)} \\ \text{cup with dot on bottom strand} \end{array} - \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.24)} \\ \text{cup with dot on bottom strand} \end{array} \stackrel{(3.6)}{=} \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.16)} \\ \text{cup with dot on bottom strand} \end{array} \\
 \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.21)} \\ \text{cup with dot on bottom strand} \end{array} \stackrel{(3.25)}{=} \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.25)} \\ \text{cup with dot on bottom strand} \end{array} + \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.25)} \\ \text{cup with dot on bottom strand} \end{array} - \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.25)} \\ \text{cup with dot on bottom strand} \end{array} \stackrel{(3.6)}{=} \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.16)} \\ \text{cup with dot on bottom strand} \end{array} \stackrel{(3.4)}{=} \begin{array}{c} \text{cup with dot on top strand} \\ \text{(3.4)} \\ \text{cup with dot on bottom strand} \end{array} .
 \end{array}$$

The left of (3.22) is similar. □

Note that (3.9)-(3.10) and (3.21)-(3.22) allow us to draw \circ 's and \bullet 's on local extrema of strands in dotted oriented Brauer-Clifford diagrams with bubbles without ambiguity. We will do so whenever convenient.

We have a downward analogue of the left relation in (3.18):

$$\begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} = \begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} = \begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} = \begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} = - \begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} = - \begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} = - \begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} = - \begin{array}{c} \text{strand with dot on top} \\ \text{(3.27)} \\ \text{strand with dot on bottom} \end{array} . \quad (3.27)$$

Before proving that normally ordered dotted oriented Brauer-Clifford diagrams with bubbles span arbitrary Hom-spaces in \mathcal{AOCB} , we first consider diagrams of type $\mathbb{1} \rightarrow \mathbb{1}$. In this case we must show the superalgebra $\text{End}_{\mathcal{AOCB}}(\mathbb{1})$ is generated by $\Delta_1, \Delta_3, \Delta_5, \dots$ where

$$\Delta_k = \begin{array}{c} \text{bubble with dot on top strand} \\ \text{(3.28)} \\ \text{bubble with dot on bottom strand} \end{array} . \quad (3.28)$$

First, note the defining relations of \mathcal{AOCB} (and the similar relations on non-upward pointed strands implied by these) along with Propositions 3.3 and 3.10 can be used to express any dotted oriented Brauer-Clifford diagram with bubbles of type $\mathbb{1} \rightarrow \mathbb{1}$ in terms of crossing-free unnested bubbles. These relations allow one to pull apart nested bubbles, pass \circ 's through crossings, cups, and caps as needed, and \bullet 's through crossings at the cost of additional diagrams with fewer \bullet 's. Repeated moves of this sort can be used to rewrite any diagram of type $\mathbb{1} \rightarrow \mathbb{1}$ into the claimed form. For example,

$$\begin{array}{c} \text{bubble with dot on top strand} \\ \text{(3.28)} \\ \text{bubble with dot on bottom strand} \end{array} = \begin{array}{c} \text{bubble with dot on top strand} \\ \text{(3.28)} \\ \text{bubble with dot on bottom strand} \end{array} - \begin{array}{c} \text{bubble with dot on top strand} \\ \text{(3.28)} \\ \text{bubble with dot on bottom strand} \end{array} - \begin{array}{c} \text{bubble with dot on top strand} \\ \text{(3.28)} \\ \text{bubble with dot on bottom strand} \end{array} - \begin{array}{c} \text{bubble with dot on top strand} \\ \text{(3.28)} \\ \text{bubble with dot on bottom strand} \end{array} .$$

$$\begin{aligned}
 &= \text{Diagram 1} - 2 \text{Diagram 2} \\
 &= \text{Diagram 3} + \text{Diagram 4} + \text{Diagram 5} - 2 \text{Diagram 6} \\
 &= \text{Diagram 7} + \text{Diagram 8} + \text{Diagram 9} - 2 \text{Diagram 10} .
 \end{aligned}$$

Moreover, we can express any clockwise bubble in terms of counterclockwise ones using computations similar to the following:

$$\text{Clockwise bubble} = \text{Diagram 1} \stackrel{(3.25)}{=} \text{Diagram 2} - \text{Diagram 3} + \text{Diagram 4} \stackrel{(3.6)}{=} \text{Diagram 5} \stackrel{(3.16)}{=} \text{Diagram 6} = \text{Diagram 7} .$$

Remark 3.11. (Compare with [BCNR, Remark 1.3]) Set $\Delta_{-1} := \mathbf{1}$, $\Delta'_{-1} := -\mathbf{1}$, and

$$\Delta'_k := k \text{Diagram 8}$$

for all integers $k \geq 0$. The computation above shows $\Delta_{-1}\Delta'_1 + \Delta_1\Delta'_{-1} = 0$. A similar computation can be used to show

$$\sum_{0 \leq i \leq \frac{k+1}{2}} \Delta_{2i-1}\Delta'_{k-2i} = 0$$

whenever k is a nonnegative odd integer.

Now, using (3.6), (3.15), (3.18), and (3.27) we can reduce the number of \circ 's on any bubble to zero or one. Finally, the following proposition shows that the only nonzero counterclockwise bubbles with at most one \circ are $\Delta_1, \Delta_3, \Delta_5, \dots$, whence they generate $\text{End}_{\mathcal{AOBC}}(\mathbf{1})$.

Proposition 3.12. *The following relations hold in \mathcal{AOBC} for any nonnegative integer k :*

$$\text{Diagram 8} \text{ with } k \text{ } \circ \text{'s} = 0, \quad \text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} = 0. \tag{3.29}$$

Proof. The left relation when $k = 0$ follows from the right of (3.6). If $k > 0$ the left relation follows from the calculation below since $\text{char } \mathbb{k} \neq 2$:

$$\text{Diagram 8} \text{ with } k \text{ } \circ \text{'s} \stackrel{(3.21)}{=} \text{Diagram 8} \text{ with } k-1 \text{ } \circ \text{'s} \stackrel{(3.27)}{=} - \text{Diagram 8} \text{ with } k-1 \text{ } \circ \text{'s} \stackrel{(3.22)}{=} - \text{Diagram 8} \text{ with } k \text{ } \circ \text{'s} .$$

The right relation follows from the calculation below:

$$\begin{aligned}
 &\text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} \stackrel{(3.6)}{=} \text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} \stackrel{(3.9)}{=} \text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} \stackrel{(2.1)}{=} - \text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} \\
 &\stackrel{(3.10)}{=} - \text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} \stackrel{(3.18)}{=} -(-1)^{2k} \text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} \stackrel{(3.6)}{=} - \text{Diagram 8} \text{ with } 2k \text{ } \circ \text{'s} .
 \end{aligned}$$

□

The following lemma implies that normally ordered oriented Brauer-Clifford diagrams with bubbles span the Hom-spaces of \mathcal{AOBC} .

Lemma 3.13. *For any $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$ the superspace $\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k}$ is equal to the \mathbb{k} -span of all equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ with at most k \bullet 's.*

Proof. Let d denote a dotted oriented Brauer-Clifford diagram with bubbles with $\deg(d) \leq k$. Let d' denote the diagram obtained from d as follows. First, remove pairs of \circ 's appearing on the same strand until each strand has at most one \circ . Next, freely slide each bubble to the right of the picture and redraw them so that they are crossing-free and counterclockwise, without changing the number of \circ 's and \bullet 's appearing on each bubble. Finally, on every non-bubble strand freely slide all \circ 's (resp. \bullet 's) until they lie on an outward-pointing (resp. inward-pointing) segment. For example, if d is the diagram on the left of (1.1), then d' is the diagram on the right. It follows from Proposition 3.12 that d' is either zero or normally ordered with $\deg(d') = \deg(d)$. Moreover, by Propositions 3.3 and 3.10 along with (3.6), (3.15), and (3.27) we have $d = \pm d' + d''$ where d'' is a linear combination of dotted oriented Brauer-Clifford diagrams with bubbles each of which having fewer than $\deg(d)$ \bullet 's. The result now follows by inducting on $\deg(d)$. \square

4. CONNECTION TO REPRESENTATIONS OF LIE SUPERALGEBRAS OF TYPE Q

We next explain how the supercategories OBC and \mathcal{AOBC} are connected with the representations of the Lie superalgebras of type Q . In what follows we assume that $\sqrt{-1}$ is an element of \mathbb{k} . This is only for convenience. In particular, the arguments in Section 6.2 imply the basis theorems hold in general.

4.1. The Lie superalgebra \mathfrak{q} . Fix a \mathbb{k} -superspace $V = V_{\bar{0}} \oplus V_{\bar{1}}$ with $\dim_{\mathbb{k}}(V_{\bar{0}}) = \dim_{\mathbb{k}}(V_{\bar{1}}) = n$. Fix a homogeneous basis $v_1, \dots, v_n, v_{\bar{1}}, \dots, v_{\bar{n}}$ with $|v_i| = \bar{0}$ and $|v_{\bar{i}}| = \bar{1}$ for $i = 1, \dots, n$. We write I for the index set $\{1, \dots, n, \bar{1}, \dots, \bar{n}\}$ and I_0 for the index set $\{1, \dots, n\}$. We adopt the convention that $\bar{\bar{i}} = i$ for all $i \in I$. Let $c : V \rightarrow V$ be the odd linear map given by $c(v_i) = (-1)^{|v_i|} \sqrt{-1} v_{\bar{i}}$ for all $i \in I$.

The vector space of all linear endomorphisms of V , $\mathfrak{gl}(V)$, is naturally \mathbb{Z}_2 -graded as in Section 2.1. Furthermore, $\mathfrak{gl}(V)$ is a Lie superalgebra under the graded commutator bracket; this, by definition, is given by $[x, y] = xy - (-1)^{|x||y|}yx$ for all homogeneous $x, y \in \mathfrak{gl}(V)$. For $i, j \in I$ we write $e_{i,j} \in \mathfrak{gl}(V)$ for the linear map $e_{i,j}(v_k) = \delta_{j,k} v_i$. These are the matrix units and they form a homogeneous basis for $\mathfrak{gl}(V)$ with $|e_{i,j}| = |v_i| + |v_j|$.

By definition $\mathfrak{q}(V)$ is the Lie subsuperalgebra of $\mathfrak{gl}(V)$ given by

$$\mathfrak{q}(V) = \{x \in \mathfrak{gl}(V) \mid [x, c] = 0\}.$$

Then $\mathfrak{q}(V)$ has a homogenous basis given by $e_{i,j}^{\bar{0}} := e_{i,j} + e_{\bar{i},\bar{j}}$ and $e_{i,j}^{\bar{1}} := e_{\bar{i},j} + e_{i,\bar{j}}$ for $1 \leq i, j \leq n$. Set $\tilde{e}_{i,j}^{\bar{0}} := e_{i,j} - e_{\bar{i},\bar{j}}$ and $\tilde{e}_{i,j}^{\bar{1}} := e_{\bar{i},j} - e_{i,\bar{j}}$ for $1 \leq i, j \leq n$. These are homogeneous elements of $\mathfrak{gl}(V)$ and, together with our basis for $\mathfrak{q}(V)$, provide a homogeneous basis for $\mathfrak{gl}(V)$. Note that $|\tilde{e}_{i,j}^{\varepsilon}| = |e_{i,j}^{\varepsilon}| = \varepsilon$ for all $1 \leq i, j \leq n$ and $\varepsilon \in \mathbb{Z}_2$.

Let $U(\mathfrak{q})$ denote the universal enveloping superalgebra of the Lie superalgebra $\mathfrak{q} = \mathfrak{q}(V)$. The superalgebra $U(\mathfrak{q})$ has a homogeneous PBW basis given by all ordered monomials in the elements $(e_{i,j}^{\varepsilon})^r$ (with $1 \leq i, j \leq n$, and $r \geq 1$ if $\varepsilon = \bar{0}$ or

$r = 0, 1$ if $\varepsilon = \bar{1}$). Set $U(\mathfrak{h})$, $U(\mathfrak{n})$, and $U(\mathfrak{n}^-)$, respectively, to be the subsuperalgebras generated by $\{e_{i,i}^\varepsilon \mid i = 1, \dots, n, \varepsilon \in \mathbb{Z}_2\}$, $\{e_{i,j}^\varepsilon \mid 1 \leq i < j \leq n, \varepsilon \in \mathbb{Z}_2\}$, and $\{e_{i,j}^\varepsilon \mid 1 \leq j < i \leq n, \varepsilon \in \mathbb{Z}_2\}$. The PBW basis implies that there is a triangular decomposition $U(\mathfrak{q}) = U(\mathfrak{n}^-) \otimes U(\mathfrak{h}) \otimes U(\mathfrak{n})$.

A $U(\mathfrak{q})$ -supermodule is a \mathbb{k} -superspace $M = M_{\bar{0}} \oplus M_{\bar{1}}$ with an action by $U(\mathfrak{q})$ which respects the \mathbb{Z}_2 -grading in that $U(\mathfrak{q})_\varepsilon M_{\varepsilon'} \subseteq M_{\varepsilon+\varepsilon'}$ for all $\varepsilon, \varepsilon' \in \mathbb{Z}_2$. In particular, the superspace V defined above is naturally a $U(\mathfrak{q})$ -supermodule. A supermodule homomorphism is a \mathbb{k} -linear map $f : M \rightarrow N$ which satisfies $f(am) = (-1)^{|f||a|}af(m)$ for all homogeneous $a \in U(\mathfrak{q})$ and $m \in M$. Note that homomorphisms are not assumed to preserve parity. However, Hom-spaces are naturally \mathbb{Z}_2 -graded as in Section 2.1. Let $U(\mathfrak{q})$ -smod denote the supercategory of all $U(\mathfrak{q})$ -supermodules. We will write $\text{Hom}_{U(\mathfrak{q})}(M, M') = \text{Hom}_{U(\mathfrak{q})\text{-smod}}(M, M')$.

The \mathbb{k} -superalgebra $U(\mathfrak{q})$ is a Hopf superalgebra. In particular, given $U(\mathfrak{q})$ -supermodules M and M' , the action of a homogeneous $x \in \mathfrak{q} \subseteq U(\mathfrak{q})$ on $M \otimes M'$ is given by $x.(m \otimes m') = (x.m) \otimes m' + (-1)^{|x||m|}m \otimes (x.m')$ for homogeneous $m \in M$ and $m' \in M'$. The unit object is given by viewing \mathbb{k} as a superspace concentrated in parity $\bar{0}$ and with trivial $U(\mathfrak{q})$ action. The symmetric braiding is given by the graded flip map. Thus $U(\mathfrak{q})$ -smod is a symmetric braided monoidal supercategory.

The antipode $\sigma : U(\mathfrak{q}) \rightarrow U(\mathfrak{q})$ is given by $\sigma(x) = -x$ for $x \in \mathfrak{q}$. Using the antipode each finite-dimensional $U(\mathfrak{q})$ -supermodule M has a dual given by $M^* = \text{Hom}_{\mathbb{k}}(M, \mathbb{k})$ with the evaluation and coevaluation maps given by the same formulas as for superspaces (see Section 2.1). In particular, V is finite-dimensional and so admits a dual, V^* .

4.2. Mixed Schur-Weyl-Sergeev duality. As Theorem 3.4 and Corollary 3.5 will be used to prove the following results, let us point out that this section is not used in the proof of Theorems 3.4 and 3.9 and Corollary 3.5.

There is a monoidal superfunctor

$$\Phi : \mathcal{OBC} \rightarrow U(\mathfrak{q})\text{-smod}$$

mapping the objects \uparrow, \downarrow to the superspaces V, V^* respectively, and defined on morphisms by

$$\begin{aligned} \Phi \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) : \mathbb{1} \rightarrow V \otimes V^*, & \quad \mathbb{1} \mapsto \sum_{i \in I} v_i \otimes v_i^*, \\ \Phi \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) : V^* \otimes V \rightarrow \mathbb{1}, & \quad f \otimes v \mapsto f(v), \\ \Phi \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) : V \otimes V \rightarrow V \otimes V, & \quad u \otimes v \mapsto (-1)^{|u||v|}v \otimes u, \\ \Phi \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) : V \rightarrow V, & \quad v \mapsto c(v). \end{aligned}$$

Indeed, a direct check confirms that Φ respects the defining relations of \mathcal{OBC} . Given $\mathbf{a} \in \langle \uparrow, \downarrow \rangle$, we write $V^{\mathbf{a}} := \Phi(\mathbf{a})$. For example, $V^{\uparrow\downarrow\uparrow} = V \otimes V^* \otimes V$.

Theorem 4.1. *If the characteristic of the ground field \mathbb{k} is zero, then Φ is full.*

Proof. We are required to show


$$\Phi : \text{Hom}_{\mathcal{OBC}}(\mathbf{a}, \mathbf{b}) \rightarrow \text{Hom}_{U(\mathfrak{q})}(V^{\mathbf{a}}, V^{\mathbf{b}}) \quad (4.1)$$

is surjective for all $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$. Suppose \mathbf{a} (resp. \mathbf{b}) consists of r_1 (resp. r'_1) \uparrow 's and r_2 (resp. r'_2) \downarrow 's. Acting by the central element $\sum_{i=1}^n e_{i,i}^{\bar{0}}$ one sees that $\text{Hom}_{U(\mathfrak{q})}(V^{\mathbf{a}}, V^{\mathbf{b}}) = 0$ unless $r_1 + r'_2 = r'_1 + r_2$.


In the nontrivial case, set $r = r_1 + r'_2 = r'_1 + r_2$ and consider the following:

$$\begin{array}{ccccc}
 \mathrm{Hom}_{\mathcal{OBC}}(\mathbf{a}, \mathbf{b}) & \longrightarrow & \mathrm{Hom}_{\mathcal{OBC}}(\downarrow^{r_2} \uparrow^{r_1}, \uparrow^{r'_1} \downarrow^{r'_2}) & \longrightarrow & \mathrm{End}_{\mathcal{OBC}}(\uparrow^r) \\
 \Phi \downarrow & & \Phi \downarrow & & \Phi \downarrow \\
 \mathrm{Hom}_{U(\mathfrak{q})}(V^{\mathbf{a}}, V^{\mathbf{b}}) & \longrightarrow & \mathrm{Hom}_{U(\mathfrak{q})}(V^{\downarrow^{r_2} \uparrow^{r_1}}, V^{\uparrow^{r'_1} \downarrow^{r'_2}}) & \longrightarrow & \mathrm{End}_{U(\mathfrak{q})}(V^{\otimes r}).
 \end{array} \tag{4.2}$$

The horizontal maps are all isomorphisms of superspaces. Indeed, the left horizontal maps are given by the symmetric braidings on \mathcal{OBC} and $U(\mathfrak{q})$ -smod. The right horizontal maps are the \mathbb{k} -linear isomorphisms that hold in any monoidal supercategory with duals. In particular, the top right horizontal map is the \mathbb{k} -linear isomorphism given on diagrams by



with inverse mapping



Since the monoidal superfunctor Φ respects the symmetric braidings and duality, the diagram given in (4.2) commutes. Thus, surjectivity of (4.1) follows from the surjectivity of the right vertical map in (4.2). However, composing the right vertical map in (4.2) with the isomorphism φ from Corollary 3.5 gives the superalgebra map $\mathrm{Ser}_r \rightarrow \mathrm{End}_{U(\mathfrak{q})}(V^{\otimes r})$ from Schur-Weyl-Sergeev duality. When \mathbb{k} has characteristic zero this is known to be surjective by [Ser2] (see also [CW, Section 3.4]). \square

Remark 4.2. When \mathbb{k} has characteristic zero, Schur-Weyl-Sergeev duality also implies the right vertical map in (4.2) is injective whenever $r \leq n$. It follows that (4.1) is an isomorphism whenever the average length of the words \mathbf{a} and \mathbf{b} is less than or equal to n . In particular, Φ prescribes an isomorphism of superalgebras $\mathrm{End}_{\mathcal{OBC}}(\mathbf{a}) \cong \mathrm{End}_{U(\mathfrak{q})}(V^{\mathbf{a}})$ whenever the length of \mathbf{a} is less than or equal to n . Coupled with Corollary 3.6 this recovers [JK, Theorem 3.5].

Remark 4.3. If \mathbb{k} has positive characteristic, then one can replace $U(\mathfrak{q})$ with the superalgebra of distributions for the supergroup $Q(n)$ and again have the superfunctor Φ . Moreover, the above argument for the fullness of Φ goes through modulo the statement that the map $\mathrm{Ser}_r \rightarrow \mathrm{End}_{U(\mathfrak{q})}(V^{\otimes r})$ from Schur-Weyl-Sergeev duality is surjective. For given r it can be deduced from [BK2] that this map is surjective whenever $n \geq r$ or the characteristic of \mathbb{k} is greater than r . As far as the authors are aware, surjectivity is not known in general. It is reasonable to expect it to hold

(and, hence, the fullness of Φ) under mild conditions on \mathbb{k} (c.f. [BD1]). Similar remarks apply to injectivity.

4.3. The monoidal superfunctor $\Psi : \mathcal{AOBC} \rightarrow \mathcal{E}nd(U(\mathfrak{q})\text{-smod})$. Let $U(\mathfrak{gl})$ (resp. $U(\mathfrak{q})$) denote the enveloping superalgebra of $\mathfrak{gl}(V)$ (resp. $\mathfrak{q}(V)$). Using the bases given in Section 4.1 we can naturally view $U(\mathfrak{q})$ as Hopf subsuperalgebra of $U(\mathfrak{gl})$.

Let $\Omega \in U(\mathfrak{gl}) \otimes U(\mathfrak{q})$ be the Casimir element given by

$$\Omega = \sum_{1 \leq i, j \leq n} \tilde{e}_{i,j}^{\bar{0}} \otimes e_{j,i}^{\bar{0}} + \tilde{e}_{i,j}^{\bar{1}} \otimes e_{j,i}^{\bar{1}}. \quad (4.3)$$

Given a $U(\mathfrak{gl})$ -supermodule W and a $U(\mathfrak{q})$ -supermodule M , we have an even linear map $W \otimes M \rightarrow W \otimes M$ given by

$$\Omega.(w \otimes m) = \sum_{1 \leq i, j \leq n} \tilde{e}_{i,j}^{\bar{0}}.w \otimes e_{j,i}^{\bar{0}}.m + (-1)^{|w|} \tilde{e}_{i,j}^{\bar{1}}.w \otimes e_{j,i}^{\bar{1}}.m,$$

for all homogeneous $w \in W$ and $m \in M$. By restriction W is a $U(\mathfrak{q})$ -supermodule and so $W \otimes M$ is a $U(\mathfrak{q})$ -supermodule via its coproduct. The action of Ω defines an even $U(\mathfrak{q})$ -supermodule homomorphism by [HKS, Theorem 7.4.1] (there it is assumed that $\mathbb{k} = \mathbb{C}$ but the calculations do not depend on this fact). Alternatively, one can use the odd invariant bilinear form given by the supertrace on \mathfrak{q} to define the so-called odd Casimir element of $U(\mathfrak{q}) \otimes U(\mathfrak{q})$ which by standard arguments commutes with the image of the coproduct. In turn, since Ω equals the product of the odd Casimir with $\sqrt{-1}c \otimes 1$ it necessarily defines a supermodule homomorphism. See [BD2, proof of Lemma 3.1] for details.

Theorem 4.4. *There is a monoidal superfunctor $\Psi : \mathcal{AOBC} \rightarrow \mathcal{E}nd(U(\mathfrak{q})\text{-smod})$ by mapping the objects \uparrow, \downarrow to the endofunctors $V \otimes -, V^* \otimes -,$ respectively, and on morphisms by*

$$\begin{aligned} \Psi \left(\begin{array}{c} \curvearrowright \end{array} \right) : \text{Id} \rightarrow V \otimes V^* \otimes -, & \quad m \mapsto \sum_{i \in I} v_i \otimes v_i^* \otimes m, \\ \Psi \left(\begin{array}{c} \curvearrowleft \end{array} \right) : V^* \otimes V \otimes - \rightarrow \text{Id}, & \quad f \otimes v \otimes m \mapsto f(v)m, \\ \Psi \left(\begin{array}{c} \times \end{array} \right) : V \otimes V \otimes - \rightarrow V \otimes V \otimes -, & \quad u \otimes v \otimes m \mapsto (-1)^{|u||v|} v \otimes u \otimes m, \\ \Psi \left(\begin{array}{c} \uparrow \bullet \end{array} \right) : V \otimes - \rightarrow V \otimes -, & \quad v \otimes m \mapsto \Omega(v \otimes m), \\ \Psi \left(\begin{array}{c} \uparrow \circ \end{array} \right) : V \otimes - \rightarrow V \otimes -, & \quad v \otimes m \mapsto c(v) \otimes m. \end{aligned}$$

Proof. To show the existence of the superfunctor requires that we verify the defining relations of \mathcal{AOBC} . The first three supernatural transformations are given by maps which are the coevaluation, evaluation, and braiding, respectively, in the supercategory \mathbf{svect} . From this it follows that (3.1) and (3.2) are satisfied. A direct calculation verifies that (3.6) is also satisfied. The relations in (3.18) follow from the verification of [HKS, (3.1.4) and (3.1.5)], keeping in mind the authors chose to assume that Clifford elements square to minus one and to tensor by V on the right. Our different choices impact the signs which appear in formulas but otherwise have no effect. \square

4.4. Bubbles and central elements of $U(\mathfrak{q})$. As is well known (e.g. [CM, Proposition 46]), the supernatural transformations from the identity superfunctor to itself identify as a superalgebra with $Z(\mathfrak{q})$, the supercenter of $U(\mathfrak{q})$. In particular, using the notation for supernatural transformations set in Section 2.2, $\Psi(\Delta_k)_{U(\mathfrak{q})} : U(\mathfrak{q}) \rightarrow U(\mathfrak{q})$ is a supermodule homomorphism and $z_k := \Psi(\Delta_k)_{U(\mathfrak{q})}(1)$ lies in $Z(\mathfrak{q})$. In this section we compute these central elements.

To do so requires further notation. For any $k \geq 1$ and $\varepsilon = (\varepsilon_k, \dots, \varepsilon_1) \in \mathbb{Z}_2^k$, define $|\varepsilon| = \varepsilon_k + \dots + \varepsilon_1$. Furthermore, define $\text{sgn}(\varepsilon) = \pm 1$ recursively by

$$\text{sgn}(\varepsilon_k, \dots, \varepsilon_1) = \begin{cases} (-1)^{(\varepsilon_k + \bar{1})|(\varepsilon_{k-1}, \dots, \varepsilon_1)|} \text{sgn}(\varepsilon_{k-1}, \dots, \varepsilon_1), & \text{if } k > 1; \\ 1, & \text{if } k = 1. \end{cases}$$

Theorem 4.5. *Let k be a positive odd integer, then the central element of $U(\mathfrak{q})$ determined by the even supernatural transformation $\Psi(\Delta_k)$ is*

$$z_k = 2 \sum_{\substack{(i_k, \dots, i_1) \in I_0^k, \\ \varepsilon = (\varepsilon_k, \dots, \varepsilon_1) \in \mathbb{Z}_2^k, \\ \text{with } |\varepsilon| = \bar{0}}} \text{sgn}(\varepsilon) e_{i_{k-1}, i_k}^{\varepsilon_k} e_{i_{k-2}, i_{k-1}}^{\varepsilon_{k-1}} \cdots e_{i_2, i_3}^{\varepsilon_3} e_{i_1, i_2}^{\varepsilon_2} e_{i_k, i_1}^{\varepsilon_1}.$$

Proof. We compute $\Psi(\Delta_k)_{U(\mathfrak{q})}(1) = (\text{ev}_V \otimes 1) \circ (1 \otimes \Omega^k) \circ (\gamma_{V, V^*} \otimes 1) \circ (\text{coev}_V \otimes 1)(1)$. First observe that

$$(\gamma_{V, V^*} \otimes 1) \circ (\text{coev}_V \otimes 1)(1) = \sum_{i \in I} (-1)^{|v_i|} v_i^* \otimes v_i \otimes 1 = (1 - c_{2,1}) \left(\sum_{i_0 \in I_0} v_{i_0}^* \otimes v_{i_0} \otimes 1 \right), \quad (4.4)$$

where $c_{2,1} := \Psi \left(\begin{array}{c} \updownarrow \\ \updownarrow \end{array} \right)_{U(\mathfrak{q})} \in \text{End}_{U(\mathfrak{q})}(V^* \otimes V \otimes U(\mathfrak{q}))$.

To continue it is helpful to introduce some notation to simplify formulas. Given $\mathbf{i} = (i_k, \dots, i_0) \in I_0^{k+1}$ and $\varepsilon = (\varepsilon_k, \dots, \varepsilon_1) \in \mathbb{Z}_2^k$, for short let $e_{\mathbf{i}}^{\varepsilon} \in U(\mathfrak{q})$ be given by

$$e_{\mathbf{i}}^{\varepsilon} = e_{i_{k-1}, i_k}^{\varepsilon_k} e_{i_{k-2}, i_{k-1}}^{\varepsilon_{k-1}} \cdots e_{i_1, i_2}^{\varepsilon_2} e_{i_0, i_1}^{\varepsilon_1}.$$

Given $\varepsilon \in \mathbb{Z}_2^k$ and $i \in I_0$ we write

$$v_i^{|\varepsilon|} = \begin{cases} v_i, & \text{if } |\varepsilon| = \bar{0}; \\ v_{\bar{i}}, & \text{if } |\varepsilon| = \bar{1}. \end{cases}$$

A straightforward induction on $k \geq 1$ proves that for each fixed $i_0 \in I_0$

$$(1 \otimes \Omega)^k \cdot v_{i_0}^* \otimes v_{i_0} \otimes 1 = \sum \text{sgn}(\varepsilon) v_{i_0}^* \otimes v_{i_k}^{|\varepsilon|} \otimes e_{\mathbf{i}}^{\varepsilon},$$

where the sum is over all $\varepsilon = (\varepsilon_k, \dots, \varepsilon_1) \in \mathbb{Z}_2^k$ and $\mathbf{i} = (i_k, \dots, i_1, i_0) \in I_0^{k+1}$. Combining this formula with (4.4) and using $(1 \otimes \Omega) \circ c_{2,1} = -c_{2,1} \circ (1 \otimes \Omega)$ and $(\text{ev}_V \otimes 1) \circ c_{2,1} = \text{ev}_V \otimes 1$ yields

$$\begin{aligned} \Psi(\Delta_k)_{U(\mathfrak{q})}(1) &= (\text{ev}_V \otimes 1) \circ (1 - (-1)^k c_{2,1}) \left(\sum \text{sgn}(\varepsilon) v_{i_0}^* \otimes v_{i_k}^{|\varepsilon|} \otimes e_{\mathbf{i}}^{\varepsilon} \right) \\ &= \begin{cases} 0, & \text{if } k \text{ is even;} \\ 2 \sum \text{sgn}(\varepsilon) e_{\mathbf{i}}^{\varepsilon}, & \text{if } k \text{ is odd.} \end{cases} \end{aligned}$$

The upper sum is over all $\mathbf{i} = (i_k, \dots, i_1, i_0) \in I_0^{k+1}$, and $\varepsilon = (\varepsilon_k, \dots, \varepsilon_1) \in \mathbb{Z}_2^k$. The lower sum is over all $\mathbf{i} \in I_0^{k+1}$ with $i_0 = i_k$ and all $\varepsilon \in \mathbb{Z}_2^k$ with $|\varepsilon| = \bar{0}$. This proves the stated result. \square

For each integer $t \geq 0$, Sergeev defined an explicit element $S_t \in Z(\mathfrak{q})$. As we do not need Sergeev's elements we do not reproduce their definition here. The interested reader can find it in [Ser1] or [BK3, Section 8]. For completeness' sake we explain how Sergeev's elements relate to those described above. Recall that $\sigma : U(\mathfrak{q}) \rightarrow U(\mathfrak{q})$ is the antipode of $U(\mathfrak{q})$ (see Section 4.1).

Proposition 4.6. *For each integer $t \geq 1$,*

$$z_{2t-1} = -2\sigma(S_t).$$

Proof. By expanding the recursive formula for S_t we see that it is a sum with coefficients of ± 1 over precisely the same set of monomials as given by the formula for z_{2t-1} in Theorem 4.5, except that they are in reverse order. That is, $\sigma(S_t)$ and z_{2t-1} are sums over precisely the same set of monomials. All that remains is to verify that the sign in front of each monomial agrees. This is a straightforward check, keeping in mind that since σ is a superalgebra anti-involution, if $x_1, \dots, x_k \in \mathfrak{q}$, then $\sigma(x_1 \cdots x_k) = (-1)^\delta x_k \cdots x_1$, where $\delta = k + \sum_{1 \leq r < s \leq k} |x_r||x_s|$. It is also helpful in comparing signs to verify that a closed formula for $\text{sgn}(\varepsilon_k, \dots, \varepsilon_1)$ is given by

$$\text{sgn}(\varepsilon_k, \dots, \varepsilon_1) = (-1)^{\varepsilon_{k-1} + \varepsilon_{k-3} + \cdots + \varepsilon_p + \sum_{1 \leq r < s \leq k} \varepsilon_r \varepsilon_s},$$

where $p = 1$ if k is even and $p = 2$ if k is odd. \square

Remark 4.7. Assume \mathbb{k} has characteristic zero. By our basis theorem for \mathcal{AOBC} the set $\{\Delta_1, \Delta_3, \Delta_5, \dots\}$ is algebraically independent (see also Remark 5.5). Moreover, Sergeev's elements are known to generate $Z(\mathfrak{q})$ for every $n \geq 1$ by [Ser1]. Therefore Ψ defines a surjective homomorphism $\text{End}_{\mathcal{AOBC}}(\mathbb{1}) \rightarrow \text{End}_{\mathcal{E}nd(U(\mathfrak{q})\text{-smod})}(\text{Id})$.

5. THE GENERIC VERMA SUPERMODULE

We now introduce the generic Verma supermodule. This supermodule will play a key role in proving the basis theorem for \mathcal{AOBC} . Analogous modules exist for Lie (super)algebras of other types. Our approach could be used to give a different proof of the basis theorem for \mathcal{AOB} in type A and to obtain similar results in other types (e.g., see [BDE⁺]).

5.1. The polynomial ring. Let $U(\mathfrak{h}_{\bar{0}})$ denote the subsuperalgebra of $U(\mathfrak{h})$ generated by $\{h_i \mid i = 1, \dots, n\}$, where for brevity we set $h_i = e_{i,i}^{\bar{0}}$ for $i = 1, \dots, n$. We put the usual \mathbb{Z} -grading on $U(\mathfrak{h}_{\bar{0}})$ by putting each h_i in degree 1. Since $\mathfrak{h}_{\bar{0}}$ is an abelian Lie superalgebra concentrated in parity zero, $U(\mathfrak{h}_{\bar{0}})$ is nothing more than a polynomial ring on generators h_n, \dots, h_1 . We refine the \mathbb{Z} -grading on $U(\mathfrak{h}_{\bar{0}})$ by also putting a graded lexicographic order on the monomials with the convention that $h_n > \cdots > h_1$; that is, $h_n^{r_n} \cdots h_1^{r_1} < h_n^{s_n} \cdots h_1^{s_1}$ if and only if either $\sum_i r_i < \sum_i s_i$ or $\sum_i r_i = \sum_i s_i$ and if $t = 1, \dots, n$ is maximal with $r_t \neq s_t$, then $r_t < s_t$.

5.2. The generic Verma supermodule. Let $U(\mathfrak{b})$ denote the subsuperalgebra of $U(\mathfrak{q})$ generated by $U(\mathfrak{h})$ and $U(\mathfrak{n})$. By the PBW theorem we have $U(\mathfrak{b}) = U(\mathfrak{h}) \otimes U(\mathfrak{n})$ as superspaces. Since the span of the monomials of positive degree, $U(\mathfrak{n})_+$, is an ideal of $U(\mathfrak{b})$ with $U(\mathfrak{b})/U(\mathfrak{n})_+ \cong U(\mathfrak{h})$, we can and will view $U(\mathfrak{h})$ as a $U(\mathfrak{b})$ -supermodule by inflation. Define the *generic Verma supermodule* to be the $U(\mathfrak{q})$ -supermodule

$$M = U(\mathfrak{q}) \otimes_{U(\mathfrak{b})} U(\mathfrak{h}) \cong U(\mathfrak{q}) \otimes_{U(\mathfrak{b})} (U(\mathfrak{h}) \otimes_{U(\mathfrak{h}_{\bar{0}})} U(\mathfrak{h}_{\bar{0}})).$$

We will write $\hat{u} := 1 \otimes 1 \otimes 1 \in M$ for the ‘‘highest weight vector’’. While we do not need this fact, note that if W is a $U(\mathfrak{q})$ -supermodule and $w \in W$ is a homogeneous weight vector for $U(\mathfrak{h}_0)$ such that $e_{i,j}^\varepsilon w = 0$ for all $1 \leq i < j \leq n$ and all $\varepsilon \in \mathbb{Z}_2$, then there is a unique $U(\mathfrak{q})$ -supermodule homomorphism $M \rightarrow W$ such that $\hat{u} \mapsto w$.

For brevity, set $N = n(n-1)/2$ and let $\{f_1, \dots, f_N\}$ (resp. $\{\bar{f}_1, \dots, \bar{f}_N\}$) be any basis for \mathfrak{n}_0^- (resp. \mathfrak{n}_1^-) such that $\{e_{i,j}^0 \mid 1 \leq j < i \leq n\} = \{f_1, \dots, f_N\}$ and $\{e_{i,j}^1 \mid 1 \leq j < i \leq n\} = \{\bar{f}_1, \dots, \bar{f}_N\}$. In particular, a PBW basis for $U(\mathfrak{n}^-)$ is given by the monomials $f_1^{a_1} \dots f_N^{a_N} \bar{f}_1^{b_1} \dots \bar{f}_N^{b_N}$, where $a_1, \dots, a_N \in \mathbb{Z}_{\geq 0}$ and $b_1, \dots, b_N \in \{0, 1\}$. By the PBW theorem M is a free right $U(\mathfrak{h}_0)$ -supermodule on basis

$$f_1^{a_1} \dots f_N^{a_N} \bar{f}_1^{b_1} \dots \bar{f}_N^{b_N} \otimes \bar{h}_1^{c_1} \dots \bar{h}_n^{c_n} \otimes 1 \quad (5.1)$$

where each $a_k \in \mathbb{Z}_{\geq 0}$ and $b_k, c_k \in \{0, 1\}$. In particular, the superspace M inherits a \mathbb{Z} -grading (hence a \mathbb{Z} -filtration) from the right action of $U(\mathfrak{h}_0)$ where for $t \in \mathbb{Z}$, M_t is the span of $f_1^{a_1} \dots f_N^{a_N} \bar{f}_1^{b_1} \dots \bar{f}_N^{b_N} \otimes \bar{h}_1^{c_1} \dots \bar{h}_n^{c_n} \otimes U(\mathfrak{h}_0)_t$ for all $a_k \in \mathbb{Z}_{\geq 0}$ and $b_k, c_k \in \{0, 1\}$. More generally, for any $\mathfrak{a} \in \langle \uparrow, \downarrow \rangle$ the superspace $V^{\mathfrak{a}} \otimes M$ is a free right $U(\mathfrak{h}_0)$ -supermodule and is similarly a graded (hence filtered) superspace.

Lemma 5.1. *Suppose $1 \leq i, j \leq n$ and $\varepsilon \in \mathbb{Z}_2$. Left multiplication by $e_{i,j}^\varepsilon$ prescribes a map $M \rightarrow M$ that is*

- (1) *homogeneous degree 0 when $i > j$ and,*
- (2) *filtered degree 1 when $i \leq j$.*

Proof. Since M is a $(U(\mathfrak{q}), U(\mathfrak{h}_0))$ -bisupermodule with the grading coming from the right action by $U(\mathfrak{h}_0)$, it suffices to show that the result of acting on (5.1) on the left by $e_{i,j}^\varepsilon$ is (1) in M_0 when $i > j$; and (2) in $M_{\leq 1}$ when $i \leq j$.

Part (1) follows from the fact that $e_{i,j}^\varepsilon$ is an element of the subsuperalgebra $U(\mathfrak{n}^-)$ whenever $i > j$. That is, since monomials of the form $f_1^{a_1} \dots f_N^{a_N} \bar{f}_1^{b_1} \dots \bar{f}_N^{b_N}$ are a basis for $U(\mathfrak{n}^-)$ by the PBW theorem, after acting by $e_{i,j}^\varepsilon$ in $U(\mathfrak{n}^-)$ one can rewrite the result as a sum of elements which lie in M_0 .

For part (2) one can argue by induction on

$$a_1 + b_1 + \dots + a_N + b_N. \quad (5.2)$$

Namely, act by $e_{i,j}^\varepsilon$ on (5.1) and then use the commutator formulas $U(\mathfrak{q})$ given in [DW, Sections 2 and 3] to rewrite the expression into a sum of terms with smaller (5.2). The base case when this sum equals zero also follows by a calculation using the commutator formulas of [DW]. \square

Lemma 5.2. *For any $\mathfrak{a} \in \langle \uparrow, \downarrow \rangle$, the map $V \otimes (V^{\mathfrak{a}} \otimes M) \rightarrow V \otimes (V^{\mathfrak{a}} \otimes M)$ given by the action of Ω is filtered degree 1.*

Proof. This follows from the definition of the Casimir (4.3) and Lemma 5.1. \square

Using the previous lemma we can show that the composition of Ψ followed by evaluation at M defines a filtered monoidal superfunctor

$$\Psi_M : \mathcal{AOBC} \rightarrow \mathfrak{fsvec}.$$

Indeed, it is straightforward to see that $\Psi_M(d)$ is filtered degree 0 whenever d is an oriented Brauer-Clifford diagram. The fact that $\Psi_M(d)$ is a filtered map with filtered degree $\deg(d)$ whenever d is a dotted oriented Brauer-Clifford diagram with bubbles follows from Lemma 5.2.

5.3. Some Ψ_M calculations. In this subsection we prove several lemmas concerning the superfunctor Ψ_M , which will be used in Section 6 to prove our basis theorem for \mathcal{AOBC} .

Lemma 5.3. *Whenever $1 \leq i \leq n$, the degree 1 components of $\Psi_M(\uparrow_{\bullet})(v_i \otimes \hat{u})$ and $\Psi_M(\uparrow_{\bullet})(v_{\bar{i}} \otimes \hat{u})$ are $v_i \otimes \hat{u}h_i$ and $-v_{\bar{i}} \otimes \hat{u}h_i$ respectively.*

Proof. Note that $e_{i,j}^{\varepsilon} \hat{u}$ has degree 0 unless $\varepsilon = 0$ and $i = j$, in which case we have $e_{i,i}^{\varepsilon} \hat{u} = h_i \hat{u} = \hat{u}h_i$. Thus, using (4.3) we see the degree 1 components of $\Psi_M(\uparrow_{\bullet})(v_i \otimes \hat{u})$ and $\Psi_M(\uparrow_{\bullet})(v_{\bar{i}} \otimes \hat{u})$ are as given by the lemma. \square

Lemma 5.4. *The degree k component of $\Psi_M(\Delta_k)(\hat{u})$ is $2\hat{u}(h_n^k + \cdots + h_1^k)$ whenever k is odd.*

Proof. In the following computation of $\Psi_M(\Delta_k)(\hat{u})$ we list only the top degree terms:

$$\begin{aligned} \hat{u} &\mapsto \sum_{1 \leq i \leq n} v_i \otimes v_i^* \otimes \hat{u} + v_{\bar{i}} \otimes v_{\bar{i}}^* \otimes \hat{u} && \text{(apply } \Psi_M(\text{cup})) \\ &\mapsto \sum_{1 \leq i \leq n} v_i^* \otimes v_i \otimes \hat{u} - v_{\bar{i}}^* \otimes v_{\bar{i}} \otimes \hat{u} && \text{(apply } \Psi_M(\text{cross})) \\ &\mapsto \sum_{1 \leq i \leq n} v_i^* \otimes v_i \otimes \hat{u}h_i^k - (-1)^k v_{\bar{i}}^* \otimes v_{\bar{i}} \otimes \hat{u}h_i^k + \cdots && \text{(apply } \Psi_M(\downarrow_{\bullet k}) \text{ and} \\ &&& \text{use Lemma 5.3)} \\ &\mapsto \sum_{1 \leq i \leq n} \hat{u}h_i^k - (-1)^k \hat{u}h_i^k + \cdots && \text{(apply } \Psi_M(\text{cup})) \end{aligned}$$

Now, assume k is odd to get $\Psi_M(\Delta_k)(\hat{u}) = 2\hat{u}(h_n^k + \cdots + h_1^k) + \cdots$. \square

Remark 5.5. Assume \mathbb{k} has characteristic zero. By taking n sufficiently large, the previous lemma along with the fact the first n power sums in h_n, \dots, h_1 are algebraically independent can be used to show $\Delta_1, \Delta_3, \Delta_5, \dots$ are algebraically independent.

The next two lemmas concern $\Psi_M(d)$ for certain diagrams of the form $d : \uparrow^r \rightarrow \uparrow^r$. It will be convenient to let $x_k : \uparrow^r \rightarrow \uparrow^r$ denote the diagram obtained from the identity diagram by placing a single \bullet on the k th strand from the right. We will also let $(i, j) : \uparrow^r \rightarrow \uparrow^r$ denote the crossing of the i th and j th strands whenever $1 \leq i, j \leq r$. For example, if $r = 8$ then

$$x_3 = \begin{array}{c} \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \\ \bullet \end{array} \quad \text{and} \quad (4, 1) = \begin{array}{c} \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \\ \text{crossing} \end{array}.$$

Lemma 5.6. *The degree 1 component of $\Psi_M(x_k)(v_r \otimes \cdots \otimes v_1 \otimes \hat{u})$ is $v_r \otimes \cdots \otimes v_1 \otimes \hat{u}h_k$.*

Proof. First, $x_k - (k, 1)x_1(k, 1)$ is a linear combination of oriented Brauer-Clifford diagrams whenever $k > 1$. Hence, since Ψ_M is filtered and oriented Brauer-Clifford diagrams are degree 0, the degree 1 components of $\Psi_M(x_k)(v_r \otimes \cdots \otimes v_1 \otimes \hat{u})$ and $\Psi_M((k, 1)x_1(k, 1))(v_r \otimes \cdots \otimes v_1 \otimes \hat{u})$ are equal. Thus, it suffices to prove the case $k = 1$, which follows from Lemma 5.3. \square

Let us fix the following notation for the remainder of the paper. Given a dotted oriented Brauer-Clifford diagram d , let $\text{undot}(d)$ denote the oriented Brauer-Clifford diagram obtained from d by removing all \bullet 's. For example, if

$$d = \begin{array}{c} \circlearrowleft \quad \circlearrowleft \quad \circlearrowleft \quad \circlearrowleft \quad \circlearrowleft \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \circlearrowright \quad \circlearrowright \quad \circlearrowright \quad \circlearrowright \quad \circlearrowright \end{array}, \quad \text{then } \text{undot}(d) = \begin{array}{c} \circlearrowleft \quad \circlearrowleft \quad \circlearrowleft \quad \circlearrowleft \quad \circlearrowleft \\ \circlearrowright \quad \circlearrowright \quad \circlearrowright \quad \circlearrowright \quad \circlearrowright \end{array}. \quad (5.3)$$

Now, suppose $d : \uparrow^r \rightarrow \uparrow^r$ is a normally ordered dotted oriented Brauer-Clifford diagram (as in Definition 3.8) without bubbles. We let $\beta_k(d)$ denote the number of \bullet 's on the k th strand of d , where we count strands right-to-left according to their position on the bottom boundary of d . In particular, we have

$$d = \text{undot}(d) \circ x_r^{\beta_r(d)} \circ \dots \circ x_1^{\beta_1(d)}.$$

Finally, given $g \in \text{End}_{\mathcal{OBC}}(\uparrow^r)$ we let $v(g) \in V^{\otimes r}$ denote the image of $v_r \otimes \dots \otimes v_1$ under $\Phi(g)$. For example, when d as in (5.3) we have

$$v(\text{undot}(d)) = -\sqrt{-1}v_4 \otimes v_3 \otimes v_6 \otimes v_1 \otimes v_5 \otimes v_2.$$

With this notation in mind, the following result follows immediately from Lemma 5.6.

Lemma 5.7. *For any normally ordered dotted oriented Brauer-Clifford diagram $d : \uparrow^r \rightarrow \uparrow^r$ without bubbles, the top degree component of $\Psi_M(d)(v_r \otimes \dots \otimes v_1 \otimes \hat{u})$ is $v(\text{undot}(d)) \otimes \hat{u} h_r^{\beta_r(d)} \dots h_1^{\beta_1(d)}$.*

6. PROOF OF THE MAIN RESULT

We can now prove the key special case of the main result. Namely, the normally ordered dotted oriented Brauer-Clifford diagrams with bubbles provide a basis for $\text{End}_{\mathcal{AOCBC}}(\uparrow^r)$.

Theorem 6.1. *Assume \mathbb{k} has characteristic zero. Then the set of equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\uparrow^r \rightarrow \uparrow^r$ form a basis for $\text{End}_{\mathcal{AOCBC}}(\uparrow^r)$.*

Proof. By Lemma 3.13 the proposed basis spans $\text{End}_{\mathcal{AOCBC}}(\uparrow^r)$. Toward showing linear independence, note that any linear combination of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\uparrow^r \rightarrow \uparrow^r$ can be written in the form

$$\sum_d f_d(\Delta_1, \Delta_3, \Delta_5, \dots) d, \quad (6.1)$$

where the sum is over all normally ordered dotted oriented Brauer-Clifford diagrams $d : \uparrow^r \rightarrow \uparrow^r$ (without bubbles); and where the f_d 's are polynomials in countably many variables, only finitely many of which are nonzero. Set $B = \{d \mid f_d \neq 0\}$.

We will show (6.1) is nonzero whenever $B \neq \emptyset$ (completing the proof of the theorem) by showing its image under Ψ_M is nonzero whenever n is sufficiently large. In turn, this will follow from the fact that, when we choose n sufficiently large to ensure the relevant power sums are algebraically independent,

$$f_d(h_n + \dots + h_1, h_n^3 + \dots + h_1^3, h_n^5 + \dots + h_1^5, \dots) \neq 0 \quad (6.2)$$

for any $d \in B$. Recall from Section 5.1 that we have a graded lexicographic ordering on the monomials of $U(\mathfrak{h})$. For the rest of the proof we assume n is large enough so that for each $d \in B$ the leading monomial of the symmetric polynomial (6.2) with respect to this ordering does not contain any of h_r, \dots, h_1 .

Given $d \in B$, it follows from Lemma 5.4 that the top degree component of $\Psi_M(f_d(\Delta_1, \Delta_3, \dots))(v_r \otimes \cdots \otimes v_1 \otimes \hat{u})$ is of the form $v_r \otimes \cdots \otimes v_1 \otimes \hat{u}g_d(h_n, \dots, h_1)$ where $g_d(h_n, \dots, h_1)$ is some homogeneous symmetric polynomial. Fix $d_0 \in B$ with

$$\deg d_0 + \deg g_{d_0} \geq \deg d + \deg g_d \text{ for all } d \in B. \quad (6.3)$$

Set

$$B_0 = \{d \in B \mid \text{undot}(d) = \text{undot}(d_0) \text{ and } \deg d_0 + \deg g_{d_0} = \deg d + \deg g_d\}.$$

It follows from (6.3) and Lemma 5.7 that the top degree component of the image of $v_r \otimes \cdots \otimes v_1 \otimes \hat{u}$ under $\Psi_M(\sum_d f_d(\Delta_1, \Delta_3, \Delta_5, \dots)d)$ is of the form

$$\sum_{d \in B_0} v(\text{undot}(d_0)) \otimes \hat{u}g_d(h_n, \dots, h_1)h_r^{\beta_r(d)} \cdots h_1^{\beta_1(d)} + w, \quad (6.4)$$

where w lies in the $U(\mathfrak{h})$ -span of the basis elements of the form $v(d) \otimes u$ with $v(d) \neq v(\text{undot}(d_0))$. Recall that, since n was chosen sufficiently large, the elements h_r, \dots, h_1 do not appear in leading monomials of each $g_d(h_n, \dots, h_1)$. Therefore, since a diagram $d \in B_0$ is completely determined by $\beta_1(d), \dots, \beta_r(d)$, it follows that the leading monomials of $g_d(h_n, \dots, h_1)h_r^{\beta_r(d)} \cdots h_1^{\beta_1(d)}$ for $d \in B_0$ are pairwise distinct. Thus (6.4) is nonzero, which implies (6.1) is nonzero, as desired. \square

6.1. Proof of Theorem 3.9. It is straightforward to see that the validity of Theorem 3.9 when \mathbb{k} has characteristic zero is equivalent to the following lemma.

Lemma 6.2. *Assume \mathbb{k} has characteristic zero. Then for any $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$ the superspace $\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k}$ has basis given by equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ with at most k \bullet 's.*

Proof. It follows from Lemma 3.13 that the proposed basis spans $\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k}$. In particular, $\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k}$ is finite-dimensional over \mathbb{k} . Hence, it suffices to show the proposed basis has size $\dim_{\mathbb{k}} \text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k}$. Now, suppose \mathbf{a} (resp. \mathbf{b}) consists of r_1 (resp. r'_1) \uparrow 's and r_2 (resp. r'_2) \downarrow 's. If $r_1 + r'_2 \neq r'_1 + r_2$ then there are no oriented Brauer-Clifford diagrams of type $\mathbf{a} \rightarrow \mathbf{b}$, whence $\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b}) = 0$. Thus, we may assume $r_1 + r'_2 = r'_1 + r_2 =: r$. In this case we have \mathbb{k} -linear isomorphisms

$$\text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k} \rightarrow \text{Hom}_{\mathcal{AOBC}}(\downarrow^{r_2} \uparrow^{r_1}, \uparrow^{r'_1} \downarrow^{r'_2})_{\leq k} \rightarrow \text{End}_{\mathcal{AOBC}}(\uparrow^r)_{\leq k}$$

defined on diagrams in the same manner as the top horizontal maps in (4.2). In particular, $\dim_{\mathbb{k}} \text{Hom}_{\mathcal{AOBC}}(\mathbf{a}, \mathbf{b})_{\leq k} = \dim_{\mathbb{k}} \text{End}_{\mathcal{AOBC}}(\uparrow^r)_{\leq k}$. On the other hand, there are precisely r strands in any dotted oriented Brauer-Clifford diagram of type $\mathbf{a} \rightarrow \mathbf{b}$. It follows that there are the same number of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles with at most k \bullet 's of type $\mathbf{a} \rightarrow \mathbf{b}$ as there are of type $\uparrow^r \rightarrow \uparrow^r$. Thus, the result follows from Theorem 6.1. \square

6.2. The positive characteristic case. We now explain how to deduce Theorem 3.9 when \mathbb{k} has positive characteristic and, more generally, is an arbitrary graded commutative \mathbb{Z}_2 -graded ring of characteristic not two. We first observe that the definitions given in Section 2.2 work equally well if \mathbb{k} is replaced with an arbitrary graded commutative \mathbb{Z}_2 -graded ring, R , and \mathbb{k} -superspaces are replaced with \mathbb{Z}_2 -graded R -modules. We refer to these as (monoidal) R -supercategories. We define \mathcal{AOBC}_R to be the monoidal R -supercategory given by the same generators and

relations used in Definition 3.7. For example, we have the integral form $\mathcal{AOBC}_{\mathbb{Z}}$ of the degenerate affine oriented Brauer-Clifford supercategory.

With the above in mind we have the following integral version of Theorem 3.9.

Theorem 6.3. *For any $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$ the \mathbb{Z} -supermodule $\text{Hom}_{\mathcal{AOBC}_{\mathbb{Z}}}(\mathbf{a}, \mathbf{b})$ is a free \mathbb{Z} -supermodule with basis given by equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$.*

Proof. Since the relations for \mathcal{AOBC} involve only integral coefficients, the arguments given in Section 3.5 apply to $\mathcal{AOBC}_{\mathbb{Z}}$ and so the normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ span $\text{Hom}_{\mathcal{AOBC}_{\mathbb{Z}}}(\mathbf{a}, \mathbf{b})$ as a \mathbb{Z} -supermodule. On the other hand, consider a finite sum

$$\sum_d f_d d, \quad (6.5)$$

where the sum is over normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ and where the f_d 's are integers. There is an obvious superfunctor $\mathcal{AOBC}_{\mathbb{Z}} \rightarrow \mathcal{AOBC}_{\mathbb{C}}$ which can be applied to (6.5) and linear independence follows from Theorem 3.9. \square

Let \mathbb{k} be a graded commutative \mathbb{Z}_2 -graded ring. If $\mathcal{C}_{\mathbb{Z}}$ denotes a (monoidal) \mathbb{Z} -supercategory, then by base change one can define a (monoidal) \mathbb{k} -supercategory $\mathcal{C}_{\mathbb{Z}} \otimes \mathbb{k}$. Namely, the objects of $\mathcal{C}_{\mathbb{Z}} \otimes \mathbb{k}$ are the objects of $\mathcal{C}_{\mathbb{Z}}$ and the morphisms are

$$\text{Hom}_{\mathcal{C}_{\mathbb{Z}} \otimes \mathbb{k}}(\mathbf{a}, \mathbf{b}) = \text{Hom}_{\mathcal{C}_{\mathbb{Z}}}(\mathbf{a}, \mathbf{b}) \otimes_{\mathbb{Z}} \mathbb{k}.$$

Composition, the monoidal structure, etc., are extended to $\mathcal{C}_{\mathbb{Z}} \otimes \mathbb{k}$ by linearity. There are obvious mutually inverse superfunctors which provide an isomorphism of monoidal supercategories between $\mathcal{AOBC}_{\mathbb{Z}} \otimes \mathbb{k}$ and $\mathcal{AOBC}_{\mathbb{k}}$. The previous theorem and base change immediately implies the following result.

Corollary 6.4. *Let \mathbb{k} be an arbitrary graded commutative \mathbb{Z}_2 -graded ring of characteristic not two. For any $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$ the \mathbb{k} -supermodule $\text{Hom}_{\mathcal{AOBC}_{\mathbb{k}}}(\mathbf{a}, \mathbf{b})$ is a free \mathbb{k} -supermodule with basis given by equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$.*

We mention one other application of our basis theorem.

Corollary 6.5. *Let \mathbb{k} be a field of characteristic not two. The subsuperalgebra of $\text{End}_{\mathcal{AOBC}}(\uparrow^r)$ consisting of linear combinations of dotted oriented Brauer-Clifford diagrams without bubbles is isomorphic to the degenerate affine Sergeev superalgebra² ASerg_r introduced in [Naz, Section 3].*

Proof. The superalgebra ASerg_r has a presentation with even generators $s_1, \dots, s_{r-1}, x_1, \dots, x_r$, and odd generators c_1, \dots, c_r subject to the relations (for all admissible i, j):

- (1) $s_i^2 = 1, s_i s_j = s_j s_i$ when $|i - j| > 1, s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$;
- (2) $c_i^2 = 1, c_i c_j = -c_j c_i$ when $i \neq j$;
- (3) $x_i x_j = x_j x_i$;
- (4) $c_i x_i = -x_i c_i, c_i x_j = x_j c_i$ when $i \neq j$;
- (5) $s_i x_i = x_{i+1} s_i - 1 - c_i c_{i+1}$.

²Also known as the degenerate affine Hecke-Clifford superalgebra.

By checking relations we see that the following defines a superalgebra homomorphism $\nu : \text{ASerg}_r \rightarrow \text{End}_{\mathcal{AOBC}}(\uparrow^r)$.

$$\nu(s_i) = \begin{array}{c} \uparrow^{r-i-1} \quad \uparrow^{i-1} \\ \diagdown \quad \diagup \\ \uparrow \quad \uparrow \end{array}, \quad \nu(c_i) = \begin{array}{c} \uparrow^{r-i} \quad \uparrow^{i-1} \\ \uparrow \quad \uparrow \\ \circ \quad \uparrow \end{array}, \quad \nu(x_i) = \begin{array}{c} \uparrow^{r-i} \quad \uparrow^{i-1} \\ \uparrow \quad \uparrow \\ \bullet \quad \uparrow \end{array}.$$

Take note that this map follows our convention of numbering strands from right-to-left. The image of this map is the subsuperalgebra of $\text{End}_{\mathcal{AOBC}}(\uparrow^r)$ spanned by the dotted oriented Brauer-Clifford diagrams without bubbles. From Corollary 6.4 and the PBW-type basis for ASerg_r given in [Kle, Theorem 14.2.2] one can verify that this map is an isomorphism onto its image. \square

7. CYCLOTOMIC QUOTIENTS

Fix $a, b \in \mathbb{Z}_{\geq 0}$ and $m_i \in \mathbb{k}$ for each $1 \leq i \leq a$. Let $f(t) = t^b \prod_{1 \leq i \leq a} (t^2 - m_i)$, $\ell = 2a + b$, and \mathcal{OBC}^f be as in Section 1.3.

7.1. Bases for cyclotomic quotients. Since \mathcal{OBC}^f is a quotient of \mathcal{AOBC} , we may interpret any dotted oriented Brauer-Clifford diagram with bubbles as a morphism in \mathcal{OBC}^f .

Theorem 7.1. *For any $\mathbf{a}, \mathbf{b} \in \langle \uparrow, \downarrow \rangle$ the superspace $\text{Hom}_{\mathcal{OBC}^f}(\mathbf{a}, \mathbf{b})$ has basis given by equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ with fewer than ℓ \bullet 's on each strand.*

A full proof of this theorem can be found in Section 8. Note it is easy to show the proposed basis in Theorem 7.1 spans the appropriate Hom-space. Indeed, any dotted oriented Brauer-Clifford diagram with bubbles having ℓ \bullet 's on one of its strands can be realized as a linear combination of diagrams with fewer total \bullet 's by using Proposition 3.10 to slide those ℓ \bullet 's to the right of the picture, and then reducing as prescribed by f . For example, the following holds in \mathcal{OBC}^f when $f(t) = t$:

7.2. Connection to the superalgebras of Gao-Rui-Song-Su. In this subsection we explain how to recover the affine and cyclotomic walled Brauer-Clifford superalgebras from [GRSS1] from our supercategories. The discussion here parallels the analogous one in [BCNR, Section 5.5].

First, \mathcal{AOBC} can be viewed as a $\mathbb{k}[\Delta_1, \Delta_3, \dots]$ -linear supercategory with the action of each Δ_k given by tensoring on the *right*: $h\Delta_k := h \otimes \Delta_k$. Given $\delta_1, \delta_2, \dots \in \mathbb{k}$ we let $\mathcal{AOBC}(\delta_1, \delta_2, \dots)$ denote the supercategory obtained by specializing each Δ_{2k-1} at δ_k . In other words, $\mathcal{AOBC}(\delta_1, \delta_2, \dots) := \mathcal{AOBC} \otimes_{\mathbb{k}[\Delta_1, \Delta_3, \dots]} \mathbb{k}$ viewing \mathbb{k} as a $\mathbb{k}[\Delta_1, \Delta_3, \dots]$ -module with Δ_{2k-1} acting as δ_k . It follows from Theorem 3.9 that the superspace $\text{Hom}_{\mathcal{AOBC}(\delta_1, \delta_2, \dots)}(\mathbf{a}, \mathbf{b})$ has basis consisting of all equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams (without bubbles) of type $\mathbf{a} \rightarrow \mathbf{b}$. We have a similar specialization for level ℓ cyclotomic quotients. Namely, let us write $f(t) = \sum_{i=0}^{\ell} a_i t^i$. Now, fix $\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor} \in \mathbb{k}$, and define δ_k recursively for $k > \lfloor \ell/2 \rfloor$ by

$$\delta_k = - \sum_{1 \leq j \leq \lfloor \ell/2 \rfloor} a_{\ell-2j} \delta_{k-j}. \tag{7.1}$$

Then the specialization $\mathcal{OBC}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor}) := \mathcal{OBC}^f \otimes_{\mathbb{k}[\Delta_1, \Delta_3, \dots]} \mathbb{k}$ is well defined. By Theorem 7.1 the superspace $\text{Hom}_{\mathcal{OBC}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor})}(\mathbf{a}, \mathbf{b})$ has basis consisting of all equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams (without bubbles) of type $\mathbf{a} \rightarrow \mathbf{b}$ with fewer than ℓ \bullet 's on each strand. For the remainder of this section we will write $ABC_{s,r}(\delta_1, \delta_2, \dots)$ and $BC_{s,r}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor})$ for the endomorphism algebras of the object $\downarrow^s \uparrow^r$ in the supercategories $\mathcal{AOCB}(\delta_1, \delta_2, \dots)$ and $\mathcal{OBC}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor})$, respectively.

Let $BC_{r,s}^{\text{aff}}$ denote the affine walled Brauer-Clifford superalgebra defined in [GRSS1, Definition 3.1]. This superalgebra is defined via odd generators c_i ($1 \leq i \leq r$), \bar{c}_j ($1 \leq j \leq s$); even generators e_1, x_1, \bar{x}_1, s_i ($1 \leq i < r$), \bar{s}_j ($1 \leq j < s$); and even central generators $\omega_{2k+1}, \bar{\omega}_k$ ($k \in \mathbb{Z}_{>0}$) subject to a long list of relations. It is an exercise in checking those relations to see that there is a well-defined superalgebra map $BC_{r,s}^{\text{aff}} \rightarrow ABC_{s,r}(\delta_1, \delta_2, \dots)$ defined by

$$\begin{array}{ll}
c_i \mapsto \sqrt{-1} \downarrow \uparrow^{s-r-i} \uparrow^{i-1} & \bar{c}_j \mapsto \sqrt{-1} \downarrow^{s-j} \downarrow^{j-1} \uparrow^r \\
x_1 \mapsto - \downarrow \uparrow^{s-r-1} \uparrow \bullet & \bar{x}_1 \mapsto \downarrow^{s-1} \uparrow^r \bullet \\
s_i \mapsto \downarrow \uparrow^{s-r-i-1} \uparrow^{i-1} & \bar{s}_j \mapsto \downarrow^{s-j-1} \uparrow^{j-1} \uparrow^r \\
e_1 \mapsto \downarrow^{s-1} \uparrow^{r-1} & \bar{\omega}_{2k} \mapsto 0 \\
\omega_{2k+1} \mapsto -\delta_{k+1} & \bar{\omega}_{2k-1} \mapsto \delta'_k
\end{array}$$

where δ'_k is defined recursively by $\delta_k - \delta'_k = \sum_{0 < i < \lfloor k/2 \rfloor} \delta_i \delta'_{k-i}$ (compare with Remark 3.11). This map factors through the quotient $\widetilde{BC}_{r,s}$ of $BC_{r,s}^{\text{aff}}$ by the additional relations $\omega_{2k-1} = -\delta_k$, $\bar{\omega}_{2k} = 0$, $\bar{\omega}_{2k-1} = \delta'_k$ for all $k \in \mathbb{Z}_{>0}$ which is precisely the specialized superalgebra in [GRSS1, Theorem 5.15]. Using our basis theorem one can easily check that the spanning set for $\widetilde{BC}_{r,s}$ described in [GRSS1, Definition 3.15 and Corollary 3.16] maps to a basis for $ABC_{s,r}(\delta_1, \delta_2, \dots)$. Hence, $\widetilde{BC}_{r,s} \cong ABC_{s,r}(\delta_1, \delta_2, \dots)$. Note that this also gives a different proof of the linear independence in [GRSS1, Theorem 5.15].

A similar discussion applies to cyclotomic quotients. Again, fix $\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor} \in \mathbb{k}$, and define δ_k for $k > \lfloor \ell/2 \rfloor$ by (7.1). In [GRSS1, Definition 3.14] the cyclotomic walled Brauer-Clifford superalgebra $BC_{\ell,r,s}$ is defined as the quotient of $\widetilde{BC}_{r,s}$ by the additional relations $f(x_1) = g(\bar{x}_1) = 0$ where $g(t)$ is another monic degree ℓ polynomial satisfying certain conditions. One can check that those conditions imply $f(x_1)$ and $g(\bar{x}_1)$ are mapped to zero under the composition of the isomorphism $BC_{r,s}^{\text{aff}} \rightarrow ABC_{s,r}(\delta_1, \delta_2, \dots)$ from the previous paragraph with the quotient map $ABC_{s,r}(\delta_1, \delta_2, \dots) \rightarrow BC_{s,r}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor})$. Hence, that composition factors through $BC_{\ell,r,s}$ to induce a surjection $BC_{\ell,r,s} \rightarrow BC_{s,r}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor})$. Now, using our basis theorem one can check that the spanning set for $BC_{\ell,r,s}$ from [GRSS1, Definition 3.15 and Corollary 3.16] maps to a basis for $BC_{s,r}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor})$. Hence, $BC_{\ell,r,s} \cong BC_{s,r}^f(\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor})$.

8. THE CYCLOTOMIC BASIS THEOREM

In this section we provide a proof of Theorem 7.1. We actually consider some slightly more general cyclotomic quotients than the ones defined so far, whose definition is similar in spirit to Rouquier's deformed cyclotomic quotients of quiver Hecke algebras from [Rou, §4.4]. Let \mathbb{K} be some commutative \mathbb{k} -algebra and consider the base-changed \mathbb{K} -linear monoidal supercategory $\mathcal{AOBC}_{\mathbb{K}} := \mathcal{AOBC} \otimes_{\mathbb{k}} \mathbb{K}$.

Fix $\ell \geq 0$ and monic polynomials

$$f(u) = z_0 u^\ell + z_1 u^{\ell-2} + z_2 u^{\ell-4} + \cdots, \quad (8.1)$$

$$f'(u) = z'_0 u^\ell + z'_1 u^{\ell-2} + z'_2 u^{\ell-4} + \cdots \quad (8.2)$$

in $\mathbb{K}[u]$. So $z_0 = z'_0 = 1$, and all powers of u in these polynomials are even or odd according to the parity of ℓ . Define the power series

$$\delta(u) = \delta_0 + \delta_1 u^{-1} + \delta_2 u^{-2} + \cdots, \quad (8.3)$$

$$\delta'(u) = \delta'_0 + \delta'_1 u^{-1} + \delta'_2 u^{-2} + \cdots \quad (8.4)$$

in $\mathbb{K}[[u^{-1}]]$ from

$$\delta(u^2) := f'(u)/f(u), \quad (8.5)$$

$$\delta'(u^2) := -f(u)/f'(u). \quad (8.6)$$

Note $\delta_0 = 1$ but $\delta'_0 = -1$. Computing the coefficients of $u^{\ell-2r}$ in $f'(u) = f(u)\delta(u^2)$ gives

$$\sum_{s=0}^r z_s \delta_{r-s} = z'_r \text{ for } r = 0, \dots, \lfloor \ell/2 \rfloor, \quad (8.7)$$

$$\sum_{s=0}^{\lfloor \ell/2 \rfloor} z_s \delta_{r-s} = 0 \text{ for } r > \lfloor \ell/2 \rfloor. \quad (8.8)$$

Let Sym be the algebra of symmetric functions over \mathbb{K} , viewed as a purely even superalgebra. Denote the elementary and complete symmetric functions by e_r and h_r as usual; in particular, $e_0 = h_0 = 1$. Working in $\text{Sym}[[u^{-1}]]$, we set

$$e(u) := e_0 + u^{-1}e_1 + u^{-2}e_2 + \cdots, \quad (8.9)$$

$$h(u) := h_0 + u^{-1}h_1 + u^{-2}h_2 + \cdots, \quad (8.10)$$

and recall the fundamental identity $e(u)h(-u) = 1$. By Theorem 3.9 and Remark 3.11, there is a well-defined superalgebra isomorphism

$$\beta : \text{Sym} \xrightarrow{\sim} \text{End}_{\mathcal{AOBC}_{\mathbb{K}}}(\mathbb{1}), \quad h_r \mapsto (-1)^r \text{⦿}_{2r-1}, \quad e_r \mapsto - \text{⦿}_{2r-1}. \quad (8.11)$$

So that this also makes sense in the case $r = 0$, it is natural to adopt the convention that $\text{⦿}_{-1} := 1_{\mathbb{1}}$ and $-1 \text{⦿}_{-1} := -1_{\mathbb{1}}$.

Lemma 8.1. *The \mathbb{K} -linear left tensor ideal $\mathcal{I}_{f,f'}$ of $\mathcal{AOBC}_{\mathbb{K}}$ generated by*

$$\left\{ f \left(\text{⦿}_{\uparrow} \right), \text{⦿}_{2r-1} - \delta_r 1_{\mathbb{1}} \mid r = 1, \dots, \lfloor \ell/2 \rfloor \right\} \quad (8.12)$$

is generated equivalently by

$$\left\{ f' \left(\text{⦿}_{\downarrow} \right), \text{⦿}_{2r-1} - \delta'_r 1_{\mathbb{1}} \mid r = 1, \dots, \lfloor \ell/2 \rfloor \right\}. \quad (8.13)$$

Moreover, $\mathcal{I}_{f,f'}$ contains $\text{⦿}_{2r-1} - \delta_r 1_{\mathbb{1}}$ and $\text{⦿}_{2r-1} - \delta'_r 1_{\mathbb{1}}$ for all $r \geq 0$.

Proof. This is similar to Lemma 1.8 of [Bru3]. We first show by induction on r that $\mathcal{I}_{f,f'}$ contains $\circlearrowleft^{2r-1} - \delta_r 1_{\mathbb{1}}$ for all $r \geq 0$. This is immediate from the definitions for $r \leq \lfloor \ell/2 \rfloor$, so assume that $r > \lfloor \ell/2 \rfloor$. Since $2r - 1 \geq \ell$, we get from $f \left(\uparrow \right) \in \mathcal{I}_{f,f'}$ that $\sum_{s=0}^{\lfloor \ell/2 \rfloor} z_s \circlearrowleft^{2(r-s)-1} \in \mathcal{I}_{f,f'}$ too. Now the following verifies the induction step:

$$\circlearrowleft^{2r-1} - \delta_r 1_{\mathbb{1}} \stackrel{(8.8)}{=} \circlearrowleft^{2r-1} + \sum_{s=1}^{\lfloor \ell/2 \rfloor} z_s \delta_{r-s} 1_{\mathbb{1}} \equiv \sum_{s=0}^{\lfloor \ell/2 \rfloor} z_s \circlearrowleft^{2(r-s)-1} \equiv 0 \pmod{\mathcal{I}_{f,f'}}.$$

Hence, recalling (8.11), we have that $\beta(h(-u^2)) \equiv \delta(u^2) 1_{\mathbb{1}} \pmod{\mathcal{I}_{f,f'}}$. Since $e(u) = h(-u)^{-1}$ and $\delta'(u) = -\delta(u)^{-1}$, it follows that $\beta(e(u^2)) \equiv -\delta'(u^2) 1_{\mathbb{1}} \pmod{\mathcal{I}_{f,f'}}$. This shows that $\circlearrowleft^{2r-1} - \delta'_r 1_{\mathbb{1}} \in \mathcal{I}_{f,f'}$ for all $r \geq 0$. Now we can show that $f' \left(\downarrow \right) \in \mathcal{I}_{f,f'}$:

$$\begin{aligned} f' \left(\downarrow \right) &= \sum_{r=0}^{\lfloor \ell/2 \rfloor} z'_r \downarrow^{\ell-2r} \stackrel{(8.7)}{=} \sum_{r=0}^{\lfloor \ell/2 \rfloor} \sum_{s=0}^r z_s \delta_{r-s} \downarrow^{\ell-2r} \equiv \sum_{s=0}^{\lfloor \ell/2 \rfloor} z_s \sum_{r=s}^{\lfloor \ell/2 \rfloor} \downarrow^{\ell-2r} \circlearrowleft^{2(r-s)-1} \\ &= \sum_{s=0}^{\lfloor \ell/2 \rfloor} z_s \downarrow^{\ell-2s} \equiv 0 \pmod{\mathcal{I}_{f,f'}}, \end{aligned}$$

where for the last equality we have used Proposition 3.10 repeatedly to pull the $\ell - 2s$ dots on the right curl down past the crossing, plus Proposition 3.12 to see many of the dotted bubbles produced are zero. Now we have shown that the left tensor ideal generated by (8.12) contains (8.13). A similar argument shows that the left tensor ideal generated by (8.13) contains (8.12), completing the proof of the lemma. \square

Definition 8.2. Define the *cyclotomic oriented Brauer-Clifford supercategory* associated to the polynomials f and f' fixed above to be the \mathbb{K} -linear supercategory $\mathcal{OBC}^{f,f'}$ that is the quotient of $\mathcal{AOBC}_{\mathbb{K}}$ by the \mathbb{K} -linear left tensor ideal $\mathcal{I}_{f,f'}$ from Lemma 8.1.

Our goal is to establish a basis theorem for the morphism spaces in $\mathcal{OBC}^{f,f'}$. As we will explain fully later on, the cyclotomic oriented Brauer-Clifford supercategory \mathcal{OBC}^f is a special case, so that Theorem 7.1 will follow from this more general result.

Continuing to work over \mathbb{K} , recall from Corollary 6.5 there is a \mathbb{K} -superalgebra homomorphism

$$\alpha : \text{ASerg}_n \rightarrow \text{End}_{\mathcal{AOBC}_{\mathbb{K}}}(\uparrow^{\otimes n}) \tag{8.14}$$

sending x_i to the \bullet on the i th strand, c_i to the \circ on the i th strand, and s_i to the crossing of the i th and $(i+1)$ th strands (numbering strands $1, \dots, n$ from right to left). By Theorem 3.9, the map

$$\alpha \otimes \beta : \text{ASerg}_n \otimes_{\mathbb{K}} \text{Sym} \rightarrow \text{End}_{\mathcal{AOBC}_{\mathbb{K}}}(\uparrow^{\otimes n}) \tag{8.15}$$

is a superalgebra isomorphism. Let Serg_n^f be the *cyclotomic Sergeev superalgebra* from [BK1, Section 3-e], namely, the quotient of ASerg_n by the two-sided ideal generated by $f(x_1)$. Composing α with the canonical quotient map $\Pi : \text{End}_{\mathcal{AOBC}_{\mathbb{K}}}(\uparrow^{\otimes n}) \rightarrow \text{End}_{\mathcal{OBC}^{f,f'}}(\uparrow^{\otimes n})$ gives a well-defined \mathbb{K} -superalgebra homomorphism

$$\gamma : \text{Serg}_n^f \rightarrow \text{End}_{\mathcal{OBC}^{f,f'}}(\uparrow^{\otimes n}). \tag{8.16}$$

The following is the key to all our subsequent arguments.

Lemma 8.3. γ is an isomorphism.

Proof. Let $\pi : \text{ASerg}_n \otimes_{\mathbb{K}} \text{Sym} \rightarrow \text{Serg}_n^f$ be the \mathbb{K} -superalgebra homomorphism that sends $a \otimes 1$ to the canonical image of a in Serg_n^f , and $1 \otimes h_r$ to $(-1)^r \delta_r$. Note that $\ker \pi$ is $I \otimes \text{Sym} + \text{ASerg}_n \otimes J$ where I is the two-sided ideal of ASerg_n generated by $f(x_1)$ and J is the two-sided ideal of Sym generated by $h_r - (-1)^r \delta_r$ for $r \geq 1$. Also let $\Pi : \text{End}_{\mathcal{AOBC}_{\mathbb{K}}}(\uparrow^{\otimes n}) \rightarrow \text{End}_{\mathcal{OBC}^{f,f'}}(\uparrow^{\otimes n})$ be the canonical quotient map as above. By directly checking it on generators, one sees that the following diagram commutes:

$$\begin{array}{ccc} \text{ASerg}_n \otimes_{\mathbb{K}} \text{Sym} & \xrightarrow[\alpha \otimes \beta]{\sim} & \text{End}_{\mathcal{AOBC}_{\mathbb{K}}}(\uparrow^{\otimes n}) \\ \pi \downarrow & & \downarrow \Pi \\ \text{Serg}_n^f & \xrightarrow{\gamma} & \text{End}_{\mathcal{OBC}^{f,f'}}(\uparrow^{\otimes n}). \end{array}$$

It follows immediately that γ is surjective. Moreover, the injectivity of γ follows if we can show that $(\alpha \otimes \beta)^{-1}(\ker \Pi) \subseteq \ker \pi$.

By the definition of $\mathcal{OBC}^{f,f'}$, $\ker \Pi$ is the subspace of $\text{End}_{\mathcal{AOBC}_{\mathbb{K}}}(\uparrow^{\otimes n})$ defined by the left tensor ideal $\mathcal{I}_{f,f'}$. This means that any element of $\ker \Pi$ is a \mathbb{K} -linear combination of morphisms $\theta : \uparrow^{\otimes n} \rightarrow \uparrow^{\otimes n}$ in $\mathcal{AOBC}_{\mathbb{K}}$ of the form

$$\theta = \sigma \circ (\lambda \otimes \rho) \circ \tau = \begin{array}{c} \begin{array}{c} \uparrow \dots \uparrow \\ \boxed{\sigma} \\ \boxed{\lambda} \quad \boxed{\rho} \\ \boxed{\tau} \\ \uparrow \dots \uparrow \end{array} \end{array}.$$

where ρ is one of the generating morphisms $f(\uparrow)$ or $\beta(h_r) - (-1)^r \delta_r$ for $\mathcal{I}_{f,f'}$, and σ, τ, λ are any other morphisms so that the compositions make sense. Thus, we must show that the inverse image under $\alpha \otimes \beta$ of such a morphism $\theta = \sigma \circ (\lambda \otimes \rho) \circ \tau$ lies in $\ker \pi = I \otimes \text{Sym} + \text{ASerg}_n \otimes J$.

If $\rho = \beta(h_r) - (-1)^r \delta_r$ for some r then $(\alpha \otimes \beta)^{-1}(\theta)$ obviously lies in $\text{ASerg}_n \otimes J$. Instead, suppose that $\rho = f(\uparrow)$. Using the relations established earlier in the paper (especially Proposition 3.10 and Proposition 3.12), we “straighten” the diagram θ leaving the ρ -coupon on the right edge fixed, to rewrite it as a \mathbb{K} -linear combination of morphisms of the following two types:

- (I) $\begin{array}{c} \begin{array}{c} \uparrow \dots \uparrow \\ \boxed{\sigma'} \\ \dots \\ \boxed{\rho} \\ \boxed{\tau'} \\ \uparrow \dots \uparrow \end{array} \otimes \delta$ for $\sigma', \tau' \in \text{Im } \alpha$ and $\delta \in \text{Im } \beta$;
- (II) $\begin{array}{c} \begin{array}{c} \uparrow \dots \uparrow \\ \boxed{\lambda'} \\ \dots \\ \boxed{\rho} \end{array} \otimes \delta$ for $\lambda' \in \text{Im}(\alpha), \delta \in \text{Im}(\beta)$ and $r > \lfloor \ell/2 \rfloor$.

These morphisms arise when the ρ -coupon ends up on a propagating strand (type I) or on a dotted bubble (type II) after straightening. The inverse image under $\alpha \otimes \beta$ of a type I morphism lies in $I \otimes \text{Sym}$. The inverse image under $\alpha \otimes \beta$ of a

Assuming that is the case, the object \mathbf{a} is a “shuffle” of the tensor $\downarrow^{\otimes n} \otimes \uparrow^{\otimes n}$ already treated. Consider the minimal length permutation of tensor factors taking \mathbf{a} to $\downarrow^{\otimes n} \otimes \uparrow^{\otimes n}$. There is corresponding isomorphism $w : \mathbf{a} \xrightarrow{\sim} \downarrow^{\otimes n} \otimes \uparrow^{\otimes n}$ in $\mathcal{OBC}^{f,f'}$ obtained by composing various rightward crossings. Hence, we get another \mathbb{K} -module isomorphism

$$\mathrm{Hom}_{\mathcal{OBC}^{f,f'}}(\downarrow^{\otimes n} \otimes \uparrow^{\otimes n}, \mathbb{1}) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{OBC}^{f,f'}}(\mathbf{a}, \mathbb{1}), \quad \theta \mapsto w \circ \theta.$$

Applying this isomorphism to the basis for $\mathrm{Hom}_{\mathcal{OBC}^{f,f'}}(\downarrow^{\otimes n} \otimes \uparrow^{\otimes n}, \mathbb{1})$ already obtained at the end of the previous paragraph, we obtain a basis for $\mathrm{Hom}_{\mathcal{OBC}^{f,f'}}(\mathbf{a}, \mathbb{1})$. It is not quite the same as the basis of normally ordered diagrams we are after, but from it we can slide \bullet 's and \circ 's along strands to obtain the desired basis (up to some signs), modulo diagrams with strictly fewer \bullet 's, just like in the opening paragraph of this proof. This means that the transition matrix between the basis in hand and the basis in mind is unitriangular when suitably ordered, which is all that is needed to complete the proof. \square

Finally we explain the connection to the cyclotomic quotients \mathcal{OBC}^f . For this, we specialize to the case that the monic polynomial $f(u)$ from (8.1) has its coefficients in the original ground field \mathbb{k} . Also let

$$f'(u) := u^\ell + z'_1 u^{\ell-2} + \dots \in \mathbb{K}[u] \quad (8.17)$$

where $\mathbb{K} := \mathbb{k}[z'_1, \dots, z'_{\lfloor \ell/2 \rfloor}]$ for *indeterminates* $z'_1, \dots, z'_{\lfloor \ell/2 \rfloor}$. Define $\delta(u), \delta'(u)$ from (8.5) and (8.6). By the identity (8.7), the coefficients $\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor}$ of $\delta(u)$ are related to the indeterminates $z'_1, \dots, z'_{\lfloor \ell/2 \rfloor}$ by a unitriangular transition matrix. Thus, \mathbb{K} is also freely generated by $\delta_1, \dots, \delta_{\lfloor \ell/2 \rfloor}$. In $\mathcal{OBC}^{f,f'}$, the scalar δ_r acts in the same way as tensoring on the right with the counterclockwise bubble with $2r - 1$ \bullet 's. So we get the following corollary immediately from Theorem 8.4.

Corollary 8.5. *Assume $f(u) \in \mathbb{k}[u]$ and $f'(u) \in \mathbb{K}[u]$ as in (8.17). For objects \mathbf{a}, \mathbf{b} in $\mathcal{OBC}^{f,f'}$, the morphism space $\mathrm{Hom}_{\mathcal{OBC}^{f,f'}}(\mathbf{a}, \mathbf{b})$ has basis as a vector space over \mathbb{k} given by equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ having fewer than ℓ \bullet 's on each strand.*

Continuing with $f(u) \in \mathbb{k}[u]$, the \mathbb{k} -linear supercategory \mathcal{OBC}^f is the quotient of \mathcal{AOBC} by the \mathbb{k} -linear left tensor ideal generated by $f \left(\begin{array}{c} \uparrow \\ \bullet \end{array} \right)$. The composition of the natural \mathbb{k} -linear superfunctor $\mathcal{AOBC} \rightarrow \mathcal{AOBC}_{\mathbb{k}}$ followed by the quotient functor $\mathcal{AOBC}_{\mathbb{k}} \rightarrow \mathcal{OBC}^{f,f'}$ factors through \mathcal{OBC}^f to induce a \mathbb{k} -linear superfunctor

$$\mathcal{OBC}^f \rightarrow \mathcal{OBC}^{f,f'}. \quad (8.18)$$

This is the identity on objects. On morphisms, it is noted already in Section 7.1 that $\mathrm{Hom}_{\mathcal{OBC}^f}(\mathbf{a}, \mathbf{b})$ is spanned as a vector space over \mathbb{k} by the equivalence classes of normally ordered dotted oriented Brauer-Clifford diagrams with bubbles of type $\mathbf{a} \rightarrow \mathbf{b}$ having fewer than ℓ \bullet 's on each strand; the proof is the same argument as in the first paragraph of the proof of Theorem 8.4. The images of these spanning morphisms under the superfunctor (8.18) give a basis for $\mathrm{Hom}_{\mathcal{OBC}^{f,f'}}(\mathbf{a}, \mathbf{b})$ thanks to Corollary 8.5. Hence, they are already linearly independent in $\mathrm{Hom}_{\mathcal{OBC}^f}(\mathbf{a}, \mathbf{b})$, and (8.18) is an isomorphism on morphism spaces. We have proved:

Corollary 8.6. *The \mathbb{k} -linear superfunctor (8.18) is an isomorphism.*

Corollaries 8.5 and 8.6 together imply Theorem 7.1.

REFERENCES

- [BCNR] J. Brundan, J. Comes, D. Nash, and A. Reynolds. A basis theorem for the affine oriented Brauer category and its cyclotomic quotients. *Quantum Topol.*, 8(1):75–112, 2017.
- [BD1] D. Benson and S. Doty. Schur-Weyl duality over finite fields. *Arch. Math. (Basel)*, 93(5):425–435, 2009.
- [BD2] J. Brundan and N. Davidson. Type A blocks of super category \mathcal{O} . *J. Algebra*, 473:447–480, 2017.
- [BD3] J. Brundan and N. Davidson. Type C blocks of super category \mathcal{O} . *ArXiv e-prints*, February 2017.
- [BDE⁺] M. Balagovic, Z. Daugherty, I. Entova-Aizenbud, I. Halacheva, J. Hennig, M. S. Im, G. Letzter, E. Norton, V. Serganova, and C. Stroppel. The affine VW supercategory. *ArXiv e-prints*, January 2018.
- [BE1] J. Brundan and A. P. Ellis. Super Kac-Moody 2-categories. *ArXiv e-prints*, January 2017.
- [BE2] J. Brundan and A. P. Ellis. Monoidal supercategories. *Comm. Math. Phys.*, 351:1045–1089, May 2017.
- [BGH⁺] G. Benkart, N. Guay, J. Hye Jung, S.-J. Kang, and S. Wilcox. Quantum walled Brauer-Clifford superalgebras. *ArXiv e-prints*, April 2014.
- [BK1] J. Brundan and A. Kleshchev. Hecke-Clifford superalgebras, crystals of type $A_{2l}^{(2)}$ and modular branching rules for \hat{S}_n . *Represent. Theory*, 5:317–403, 2001.
- [BK2] J. Brundan and A. Kleshchev. Projective representations of symmetric groups via Sergeev duality. *Math. Z.*, 239(1):27–68, 2002.
- [BK3] J. Brundan and A. Kleshchev. Modular representations of the supergroup $Q(n)$. I. *J. Algebra*, 260(1):64–98, 2003. Special issue celebrating the 80th birthday of Robert Steinberg.
- [Bru1] J. Brundan. Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra $q(n)$. *Adv. Math.*, 182(1):28–77, 2004.
- [Bru2] J. Brundan. Representations of oriented skein categories. *ArXiv e-prints*, December 2017.
- [Bru3] J. Brundan. On the definition of Heisenberg category. *ArXiv e-prints*, September 2017.
- [Che] C.-W. Chen. Reduction method for representations of queer Lie superalgebras. *J. Math. Phys.*, 57(5):051703, 12, 2016.
- [CKW] S.-J. Cheng, J.-H. Kwon, and W. Wang. Character formulae for queer Lie superalgebras and canonical bases of types A/C . *Comm. Math. Phys.*, 352(3):1091–1119, 2017.
- [CM] K. Coulembier and V. Mazorchuk. The G-centre and gradable derived equivalences. *ArXiv e-prints*, March 2017.
- [CW] S.-J. Cheng and W. Wang. *Dualities and representations of Lie superalgebras*, volume 144 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.
- [DW] J. Du and J. Wan. Presenting queer Schur superalgebras. *Int. Math. Res. Not. IMRN*, (8):2210–2272, 2015.
- [EL] A. P. Ellis and A. D. Lauda. An odd categorification of $U_q(\mathfrak{sl}_2)$. *Quantum Topol.*, 7(2):329–433, 2016.
- [GRSS1] M. Gao, H. Rui, L. Song, and Y. Su. Affine walled Brauer-Clifford superalgebras. *ArXiv e-prints*, August 2017.
- [GRSS2] M. Gao, H. Rui, L. Song, and Y. Su. A proof of Comes-Kujawa’s conjecture. *ArXiv e-prints*, January 2018.
- [HKS] D. Hill, J. R. Kujawa, and J. Sussan. Degenerate affine Hecke-Clifford algebras and type Q Lie superalgebras. *Math. Z.*, 268(3-4):1091–1158, 2011.
- [JK] J. Jung and S. Kang. Mixed Schur-Weyl-Sergeev duality for queer Lie superalgebras. *J. Algebra*, 399:516–545, 2014.
- [KKO1] S.-J. Kang, M. Kashiwara, and S.-j. Oh. Supercategorification of quantum Kac-Moody algebras. *Adv. Math.*, 242:116–162, 2013.
- [KKO2] S.-J. Kang, M. Kashiwara, and S.-j. Oh. Supercategorification of quantum Kac-Moody algebras II. *Adv. Math.*, 265:169–240, 2014.
- [Kle] A. Kleshchev. *Linear and projective representations of symmetric groups*, volume 163 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2005.
- [KT] J. R. Kujawa and B. Tharp. The marked Brauer category. *J. London Mathematical Society*, 95:393–413, 2017.

- [Naz] M. Nazarov. Young's symmetrizers for projective representations of the symmetric group. *Adv. Math.*, 127(2):190–257, 1997.
- [PS] I. Penkov and V. Serganova. Characters of irreducible G -modules and cohomology of G/P for the Lie supergroup $G = Q(N)$. *J. Math. Sci. (New York)*, 84(5):1382–1412, 1997. Algebraic geometry, 7.
- [Rey] A. Reynolds. *Representations of the Oriented Brauer Category*. PhD thesis, University of Oregon, Eugene, Oregon, June 2015.
- [Rou] R. Rouquier. Quiver hecke algebras and 2-lie algebras. *Algebra Colloq.*, 19:359–410, 2012.
- [RS] D. Rosso and A. Savage. A general approach to Heisenberg categorification via wreath product algebras. *Math. Z.*, 286(1-2):603–655, 2017.
- [Ser1] A. N. Sergeev. The centre of enveloping algebra for Lie superalgebra $Q(n, \mathbf{C})$. *Lett. Math. Phys.*, 7(3):177–179, 1983.
- [Ser2] A. N. Sergeev. Tensor algebra of the identity representation as a module over the Lie superalgebras $Gl(n, m)$ and $Q(n)$. *Mat. Sb. (N.S.)*, 123(165)(3):422–430, 1984.

J.B.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON
E-mail address: brundan@uoregon.edu

J.C.: DEPARTMENT OF MATHEMATICS & PHYSICAL SCIENCES, THE COLLEGE OF IDAHO
E-mail address: jonnycomes@gmail.com

J.K.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA
E-mail address: kujawa@math.ou.edu