

**CAPACITIES IN WIENER SPACE,  
QUASI-SURE LOWER FUNCTIONS,  
AND KOLMOGOROV'S  $\varepsilon$ -ENTROPY**

DAVAR KHOSHNEVISAN, DAVID A. LEVIN, AND PEDRO J. MÉNDEZ-HERNÁNDEZ

ABSTRACT. We propose a set-indexed family of capacities  $\{\text{cap}_G\}_{G \subseteq \mathbf{R}_+}$  on the classical Wiener space  $C(\mathbf{R}_+)$ . This family interpolates between the Wiener measure  $(\text{cap}_{\{0\}})$  on  $C(\mathbf{R}_+)$  and the standard capacity  $(\text{cap}_{\mathbf{R}_+})$  on Wiener space. We then apply our capacities to characterize all quasi-sure lower functions in  $C(\mathbf{R}_+)$ . In order to do this we derive the following capacity estimate (Theorem 2.3) which may be of independent interest: There exists a constant  $a > 1$  such that for all  $r > 0$ ,

$$\frac{1}{a} K_G(r^6) e^{-\pi^2/(8r^2)} \leq \text{cap}_G\{f^* \leq r\} \leq a K_G(r^6) e^{-\pi^2/(8r^2)}.$$

Here,  $K_G$  denotes the Kolmogorov  $\varepsilon$ -entropy of  $G$ , and  $f^* := \sup_{[0,1]} |f|$ .

1. INTRODUCTION

Let  $C(\mathbf{R}_+)$  denote the collection of all continuous functions  $f : \mathbf{R}_+ \rightarrow \mathbf{R}$ . We endow  $C(\mathbf{R}_+)$  with its usual topology of uniform convergence on compacts as well as the corresponding Borel  $\sigma$ -algebra  $\mathcal{B}$ . In keeping with the literature, elements of  $\mathcal{B}$  are called *events*.

Denote by  $\mu$  the Wiener measure on  $(C(\mathbf{R}_+), \mathcal{B})$ . Recall that an event  $\Lambda$  is said to hold almost surely [a.s.] if  $\mu(\Lambda) = 1$ .

Next we define  $U := \{U_s\}_{s \geq 0}$  to be the *Ornstein–Uhlenbeck process* on  $C(\mathbf{R}_+)$ . The process  $U$  is characterized by the following requirements:

- (1) It is a stationary infinite-dimensional diffusion with value in  $C(\mathbf{R}_+)$ ;
- (2) Its invariant measure is  $\mu$ . This implies that for any fixed  $s \geq 0$ ,  $\{U_s(t)\}_{t \geq 0}$  is a standard linear Brownian motion.
- (3) For any given  $t \geq 0$ ,  $\{U_s(t)\}_{s \geq 0}$  is a standard Ornstein–Uhlenbeck process on  $\mathbf{R}$ ; i.e., it satisfies the stochastic differential equation,

$$(1.1) \quad dU_s(t) = -U_s(t) ds + \sqrt{2} dX_s \quad \forall s \geq 0,$$

where  $X$  is a Brownian motion.

Following P. Malliavin (1979), we say that an event  $\Lambda$  holds *quasi-surely* [q.s.] if

$$(1.2) \quad \mathbb{P} \{U_s \in \Lambda \text{ for all } s \geq 0\} = 1.$$

Because  $t \mapsto U_s(t)$  is a Brownian motion, any event  $\Lambda$  that holds q.s. also holds a.s. The converse is not always true. For example, define  $\Lambda_0$  to be the collection of all

---

*Date:* September 25, 2004.

*1991 Mathematics Subject Classification.* 60J45, 60J65, 28C20.

*Key words and phrases.* Capacity in Wiener space, lower functions, Kolmogorov entropy.

The research of D. Kh. is partially supported by a grant from the NSF.

functions  $f \in C(\mathbf{R}_+)$  that satisfy  $f(1) \neq 0$  (Fukushima, 1984). Evidently,  $\Lambda_0$  holds a.s. because with probability one Brownian motion at time one is not at the origin. On the other hand,  $\Lambda_0$  does not hold q.s. because  $\{U_s(1)\}_{s \geq 0}$  is point-recurrent. So the chances are 100% that  $U_s(1) = 0$  for some  $s \geq 0$ .

Despite the preceding disclaimer, a number of interesting classical events of full Wiener measure do hold q.s. A notable example is a theorem of M. Fukushima (1984). We can state it, somewhat informally, as follows:

(1.3) The Law of the Iterated Logarithm (LIL) of Khintchine (1933) holds q.s.

It might help to recall Khintchine's theorem: For  $\mu$ -every  $f \in C(\mathbf{R}_+)$ ,

$$(1.4) \quad \limsup_{t \rightarrow \infty} \frac{f(t)}{\sqrt{2t \ln \ln t}} = 1.$$

Thus we are led to the precise formulation of (1.3): With probability one, the continuous function  $f := U_s$  satisfies (1.4), simultaneously for all  $s \geq 0$ .

For another example consider "the other LIL" which was discovered by K. L. Chung (1948). Chung's LIL states that for  $\mu$ -almost every  $f \in C(\mathbf{R}_+)$ ,

$$(1.5) \quad \liminf_{t \rightarrow \infty} \frac{\sup_{u \in [0, t]} |f(u)|}{\sqrt{t / \ln \ln t}} = \frac{\pi}{\sqrt{8}}.$$

Fukushima's method can be adapted to prove that

(1.6) Chung's LIL holds q.s.

To be more precise: With probability one, the continuous function  $f := U_s$  satisfies (1.5) simultaneously for all  $s \geq 0$ .

T. S. Mountford (1992) has derived the quasi-sure integral test corresponding to (1.3). One of the initial aims of this article was to complement Mountford's theorem by finding a precise quasi-sure integral test for (1.6). Before presenting this work, let us introduce the notion of "relative capacity."

For all Borel sets  $G \subseteq \mathbf{R}_+$  and  $\Lambda \in C(\mathbf{R}_+)$  define

$$(1.7) \quad \text{cap}_G(\Lambda) := \int_0^\infty \mathbb{P} \{U_s \in \Lambda \text{ for some } s \in G \cap [0, \sigma]\} e^{-\sigma} d\sigma.$$

We think of  $\text{cap}_G(\Lambda)$  as the *capacity of  $\Lambda$  relative to the coordinates in  $G$* . The special case  $\text{cap}_{\mathbf{R}_+}$  is well known and well studied (Fukushima, 1984);  $\text{cap}_{\mathbf{R}_+}$  is called *the capacity on Wiener space*. According to (1.2), an event  $\Lambda$  holds q.s. iff its complement has zero  $\text{cap}_{\mathbf{R}_+}$ -capacity.

The case where  $G := \{s\}$  is a singleton is even better studied because of the simple fact that  $\text{cap}_{\{s\}}$  is a multiple of the Wiener measure. Thus,  $G \mapsto \text{cap}_G(\Lambda)$  interpolates from the Wiener measure ( $G = \{0\}$ ) to the standard capacity on Wiener space ( $G = \mathbf{R}_+$ ). This "interpolation" property was announced in the Abstract.

Now let  $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  be decreasing and measurable, and define

$$(1.8) \quad \mathcal{L}(H) := \left\{ f \in C(\mathbf{R}_+) : \liminf_{t \rightarrow \infty} \left[ \sup_{u \in [0, t]} |f(u)| - H(t)\sqrt{t} \right] > 0 \right\}.$$

A decreasing measurable function  $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  is called an *a.s.-lower function* if  $\mathcal{L}(H)$  holds a.s.; i.e.,  $\mu$ -almost every  $f \in C(\mathbf{R}_+)$  is in  $\mathcal{L}(H)$ . Likewise,  $H$  is

called a *q.s.-lower function* if  $\mathcal{L}(H)$  holds q.s. [The literature actually calls the function  $t \mapsto H(t)\sqrt{t}$  an a.s.[q.s]-lower function if  $\mathcal{L}(H)$  holds a.s.[q.s.], but we find our parameterization here convenient.]

To understand the utility of these definitions better, consider the special case that  $H(t) = \sqrt{c/\ln \ln t}$  for a fixed  $c > 0$  ( $t \geq 0$ ). In this case, Chung's LIL (1.5) states that  $\mathcal{L}(H)$  holds a.s. if  $c < \pi/\sqrt{8}$ ; its complement holds a.s. if  $c > \pi/\sqrt{8}$ . In fact, a precise P-a.s. integral test is known (Chung, 1948); see Corollary 1.3 below.

We aim to characterize exactly when  $(\mathcal{L}(H))^{\mathbb{G}}$  has positive  $\text{cap}_G$ -capacity. Define  $K_G$  to be the *Kolmogorov  $\varepsilon$ -entropy* of  $G$  (Dudley, 1973; Tihomirov, 1963); i.e., for any  $\varepsilon > 0$ ,  $k = K_E(\varepsilon)$  is the maximal number of points  $x_1, \dots, x_k \in E$  such that whenever  $i \neq j$ ,  $|x_i - x_j| \geq \varepsilon$ .

**Theorem 1.1.** *Choose and fix a decreasing measurable function  $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ , and a bounded Borel set  $G \subset \mathbf{R}_+$ . Then,  $\text{cap}_G((\mathcal{L}(H))^{\mathbb{G}}) = 0$  if and only if there exists a decomposition  $G = \cup_{n=1}^{\infty} G_n$  in terms of closed sets  $\{G_n\}_{n=1}^{\infty}$ , such that*

$$(1.9) \quad \int_1^{\infty} \frac{K_{G_n}(H^6(s))}{sH^2(s)} \exp\left(-\frac{\pi^2}{8H^2(s)}\right) ds < \infty \quad \forall n \geq 1.$$

Theorem 1.1 yields the following definite refinement of (1.5).

**Corollary 1.2.** *Choose and fix a decreasing measurable function  $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ . Then,  $\mathcal{L}(H)$  holds q.s. if and only if*

$$(1.10) \quad \int_1^{\infty} \exp\left(-\frac{\pi^2}{8H^2(s)}\right) \frac{ds}{sH^8(s)} < \infty.$$

Theorem 1.1 also contains the original almost-sure integral test of Chung (1948). To prove this, simply plug  $G = \{u\}$  in Theorem 1.1. Then,  $K_{\{u\} \cap J}(\varepsilon)$  is one if  $u \in J$  and zero otherwise. Thus we obtain the following.

**Corollary 1.3** (Chung (1948)). *Choose and fix a decreasing measurable function  $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ . Then  $\mathcal{L}(H)$  holds a.s. if and only if*

$$(1.11) \quad \int_1^{\infty} \exp\left(-\frac{\pi^2}{8H^2(s)}\right) \frac{ds}{sH^2(s)} < \infty.$$

To put the preceding in perspective define

$$(1.12) \quad H_{\nu}(t) := \frac{\pi}{\sqrt{8(\ln_+ \ln_+ t + \nu \ln_+ \ln_+ \ln_+ t)}} \quad \forall t, \nu > 0.$$

[ $1/0 := \infty$ ] Then, we can deduce from Corollaries 1.2 and 1.3 that  $\mathcal{L}(H_{\nu})$  occurs q.s. iff  $\nu > 5$ , whereas  $\mathcal{L}(H_{\nu})$  occurs a.s. iff  $\nu > 2$ . In particular,  $\mathcal{L}(H_{\nu})$  occurs a.s. but not q.s. if  $\nu \in [2, 5)$ . The following is another interesting consequence of Theorem 1.1.

**Corollary 1.4.** *Let  $G \subseteq [0, 1]$  be a non-random Borel set. Then,*

$$(1.13) \quad \begin{aligned} \dim_{\varphi} G > \frac{\nu - 2}{3} &\implies \text{cap}_G\left((\mathcal{L}(H_{\nu}))^{\mathbb{G}}\right) > 0, \text{ whereas} \\ \dim_{\varphi} G < \frac{\nu - 2}{3} &\implies \text{cap}_G\left((\mathcal{L}(H_{\nu}))^{\mathbb{G}}\right) = 0. \end{aligned}$$

Here,  $\dim_{\varphi} G$  denotes the *packing dimension* (Mattila, 1995) of the set  $G$ .

Throughout this paper, uninteresting constants are denoted by  $a, b, \alpha, A$ , etc. Their values may change from line to line.

**Acknowledgements.** We wish to thank Professor Zhan Shi for generously sharing with us the English translation of Lifshits and Shi (2003).

## 2. BROWNIAN SHEET, AND CAPACITY IN WIENER SPACE

We will be working with a special construction of the process  $U$ . This construction is due to D. Williams (Meyer, 1982, Appendix).

Let  $B := \{B(s, t)\}_{s, t \geq 0}$  denote a two-parameter Brownian sheet. This means that  $B$  is a centered, continuous, Gaussian process with

$$(2.1) \quad \text{Cov}(B(s, t), B(s', t')) = \min(s, s') \times \min(t, t') \quad \forall s, s', t, t' \geq 0.$$

The Ornstein–Uhlenbeck process  $U = \{U_s\}_{s \geq 0}$  on  $C(\mathbf{R}_+)$  is precisely the infinite-dimensional process that is defined by

$$(2.2) \quad U_s(t) = \frac{B(e^s, t)}{e^{s/2}} \quad \forall s, t \geq 0.$$

Indeed, one can check directly that  $U$  is a  $C(\mathbf{R}_+)$ -valued, stationary, symmetric diffusion. And for every  $t \geq 0$ ,  $\{U_s(t)\}_{s \geq 0}$  solves the stochastic differential equation (1.1) of the Ornstein–Uhlenbeck type. Furthermore, the invariant measure of  $U$  is the Wiener measure.

The following well-known result is a useful localization tool.

**Lemma 2.1.** *For all bounded Borel sets  $G \subseteq \mathbf{R}_+$  and  $\Lambda \in \mathcal{B}$ ,  $\text{cap}_G(\Lambda) > 0$  iff with positive probability there exists  $s \in G$  such that  $U_s \in \Lambda$ .*

*Remark 2.2.* The previous lemma continues to hold even when  $G$  is unbounded.

*Proof.* Without loss of much generality, we may—and will—assume that  $G \subseteq [0, q]$  for some  $q > 0$ . Let  $p_G(\Lambda)$  denote the probability that there exists  $s \in G$  such that  $U_s \in \Lambda$ . Evidently,  $\text{cap}_G(\Lambda) \leq p_G(\Lambda)$ . Furthermore,  $\text{cap}_G(\Lambda) = \int_0^q \mathbb{P}\{\exists s \in G \cap [0, \tau] : U_s \in \Lambda\} e^{-\tau} d\tau + e^{-q} p_G(\Lambda)$ , whence the bounds,

$$(2.3) \quad e^{-q} p_G(\Lambda) \leq \text{cap}_G(\Lambda) \leq p_G(\Lambda).$$

The lemma follows. □

Define

$$(2.4) \quad f^* := \sup_{u \in [0, 1]} |f(u)| \quad \forall f \in C(\mathbf{R}_+).$$

The following is the main step in the proof of Theorem 1.1. It was announced earlier in the Abstract.

**Theorem 2.3.** *There exists  $a > 1$  such that for all  $r \in (0, 1)$  and all Borel sets  $G \subseteq [0, 1]$ ,*

$$(2.5) \quad \frac{1}{a} \mathbb{K}_G(r^6) e^{-\pi^2/(8r^2)} \leq \text{cap}_G \{f^* \leq r\} \leq a \mathbb{K}_G(r^6) e^{-\pi^2/(8r^2)}.$$

*Remark 2.4.* The constant  $a$  depends on  $G$  only through the fact that  $G$  is a subset of  $[0, 1]$ . Therefore, there exists  $a > 1$  such that simultaneously for all Borel sets  $F, G \subseteq [0, 1]$ ,

$$(2.6) \quad \frac{1}{a} \frac{\mathbb{K}_F(r^6)}{\mathbb{K}_G(r^6)} \leq \frac{\text{cap}_F \{f^* \leq r\}}{\text{cap}_G \{f^* \leq r\}} \leq a \frac{\mathbb{K}_F(r^6)}{\mathbb{K}_G(r^6)} \quad \forall r \in (0, 1).$$

*Remark 2.5.* It turns out that for any fixed  $\varepsilon > 0$ ,  $\text{cap}_{\mathbf{R}_+}$  and  $\text{cap}_{[0,\varepsilon]}$  are equivalent. To prove this, we can assume without loss of generality that  $\varepsilon \in (0, 1)$ . [This is because  $\varepsilon \mapsto \text{cap}_{[0,\varepsilon]}(\Lambda)$  is increasing.] Now, on one hand,  $\text{cap}_{[0,\varepsilon]}(\Lambda) \leq \text{cap}_{\mathbf{R}_+}(\Lambda)$ . On the other hand,

$$(2.7) \quad \begin{aligned} \text{cap}_{\mathbf{R}_+}(\Lambda) &\leq \int_0^\infty \sum_{0 \leq j \leq \sigma/\varepsilon} \mathbb{P} \{ \exists s \in [j\varepsilon, (j+1)\varepsilon] : U_s \in \Lambda \} e^{-\sigma} d\sigma \\ &\leq \mathbb{P} \{ \exists s \in [0, \varepsilon] : U_s \in \Lambda \} \int_0^\infty \frac{\sigma+1}{\varepsilon} e^{-\sigma} d\sigma, \end{aligned}$$

by stationarity. In the notation of Lemma 2.1, the last term is  $(2/\varepsilon)p_{[0,\varepsilon]}(\Lambda) \leq (2e/\varepsilon)\text{cap}_{[0,\varepsilon]}(\Lambda)$ ; cf. (2.3). Thus,

$$(2.8) \quad \frac{\varepsilon}{2e} \text{cap}_{\mathbf{R}_+}(\Lambda) \leq \text{cap}_{[0,\varepsilon]}(\Lambda) \leq \text{cap}_{\mathbf{R}_+}(\Lambda) \quad \forall \Lambda \in \mathcal{B}.$$

This proves amply the claimed equivalence of  $\text{cap}_{[0,\varepsilon]}$  and  $\text{cap}_{\mathbf{R}_+}$ .

According to the eigenfunction expansion of Chung (1948),

$$(2.9) \quad \mu \{ f^* \leq r \} \sim \frac{4}{\pi} e^{-\pi^2/(8r^2)} \quad (r \rightarrow 0).$$

Therefore, thanks to (2.3), Theorem 2.3 is equivalent to our next result.

**Theorem 2.6.** *Recall that  $U_s^* = \sup_{t \in [0,1]} |U_s(t)|$  [eq. (2.4)]. Then, there exists a constant  $a > 1$  such that for all  $r \in (0, 1)$  and all Borel sets  $G \subseteq [0, 1]$ ,*

$$(2.10) \quad \frac{1}{a} \text{K}_G(r^6) \mu \{ f^* \leq r \} \leq \mathbb{P} \left\{ \inf_{s \in G} U_s^* \leq r \right\} \leq a \text{K}_G(r^6) \mu \{ f^* \leq r \}.$$

We will derive this particular reformulation of Theorem 2.3. The following result plays a key role in our analysis.

**Proposition 2.7** (Lifshits and Shi (2003, Proposition 2.1)). *Let  $\{X_t\}_{t \geq 0}$  denote planar Brownian motion. For every  $r > 0$  and  $\lambda \in (0, 1]$  define*

$$(2.11) \quad \mathcal{D}_\lambda^r = \left\{ (x, y) \in \mathbf{R}^2 : |x| \leq r, \left| x\sqrt{1-\lambda} + y\sqrt{\lambda} \right| \leq r \right\}.$$

*Then there exists an  $a \in (0, 1/2)$  such that for all  $r > 0$  and  $\lambda \in (0, 1]$ ,*

$$(2.12) \quad \mathbb{P} \{ X_t \in \mathcal{D}_\lambda^r \quad \forall t \in [0, 1] \} \leq \frac{1}{a} \mu \{ f^* \leq r \} e^{-a\lambda^{1/3}/r^2}.$$

**Lemma 2.8.** *There exists a constant  $a \in (0, 1)$  such that for all  $1 \geq S > s > 0$ ,*

$$(2.13) \quad \mathbb{P} \{ U_s^* \leq r, U_S^* \leq r \} \leq \frac{1}{a} \mu \{ f^* \leq r \} e^{-a(S-s)^{1/3}/r^2} \quad \forall r \in (0, 1).$$

*Proof.* Define  $\lambda = 1 - e^{-(S-s)}$ . Then owing to (2.2) we can write

$$(2.14) \quad U_S(t) = U_s(t)\sqrt{1-\lambda} + \frac{B(e^S, t) - B(e^s, t)}{\sqrt{e^S - e^s}} \sqrt{\lambda} := U_s(t)\sqrt{1-\lambda} + V(t)\sqrt{\lambda}.$$

By the Markov properties of the Brownian sheet,  $X_t := (U_s(t), V(t))$  defines a planar Brownian motion. Moreover,  $\mathbb{P} \{ U_s^* \leq r, U_S^* \leq r \} = \mathbb{P} \{ X_t \in \mathcal{D}_\lambda^r, \forall t \in [0, 1] \}$ . By Taylor's expansion,  $1 - e^{-x} \geq (x/2)$  ( $x \in [0, 1]$ ). Therefore, Proposition 2.7 completes the proof.  $\square$

*Proof of Theorem 2.6: Lower Bound.* Let  $k = K_G(r^6)$ , and choose maximal Kolmogorov points  $s(1) < \dots < s(k)$  such that  $s(i+1) - s(i) \geq r^6$ . Evidently, whenever  $j > i$  we have  $s(j) - s(i) \geq (j-i)r^6$ . Now define

$$(2.15) \quad N_r = \sum_{i=1}^k \mathbf{1}_{\{U_{s(i)}^* \leq r\}}.$$

According to Lemma 2.8,

$$(2.16) \quad \begin{aligned} \mathbb{E}[N_r^2] &= k\mu\{f^* \leq r\} + 2 \sum_{i=1}^{k-1} \sum_{j=i+1}^k \mathbb{P}\{U_{s(i)}^* \leq r, U_{s(j)}^* \leq r\} \\ &\leq k\mu\{f^* \leq r\} + \frac{2}{a}\mu\{f^* \leq r\} \sum_{i=1}^{k-1} \sum_{j=i+1}^k \exp\left(-\frac{a(s(j) - s(i))^{1/3}}{r^2}\right) \\ &\leq k\mu\{f^* \leq r\} + \frac{2}{a}\mu\{f^* \leq r\} \sum_{i=1}^{k-1} \sum_{j=i+1}^k \exp\left(-a(j-i)^{1/3}\right) \\ &\leq Ak\mu\{f^* \leq r\}. \end{aligned}$$

Note that  $A$  is a positive and finite constant that does not depend on  $r$ . Also note that  $\mathbb{E}[N_r] = k\mu\{f^* \leq r\}$ . This and the Paley–Zygmund inequality (Khoshnevisan, 2002, Lemma 1.4.1, p. 72) together reveal that

$$(2.17) \quad \mathbb{P}\left\{\inf_{s \in G} U_s^* \leq r\right\} \geq \mathbb{P}\{N_r > 0\} \geq \frac{(\mathbb{E}[N_r])^2}{\mathbb{E}[N_r^2]} \geq \frac{k}{A}\mu\{f^* \leq r\}.$$

The definition of  $k$  implies the lower bound in Theorem 2.6.  $\square$

Before proving the upper bound of Theorem 2.6 in complete generality, we first derive the following weak form:

**Proposition 2.9.** *There exists a finite constant  $a > 1$  such that for all  $r \in (0, 1)$ ,  $\mathbb{P}\{\inf_{s \in [0, r^6]} U_s^* \leq r\} \leq a\mu\{f^* \leq r\}$ .*

*Proof.* Recall (2.15), and define

$$(2.18) \quad L(s; r) = \int_0^s \mathbf{1}_{\{U_\nu^* \leq r\}} d\nu \quad \forall s, r > 0.$$

Let  $\mathcal{F} := \{\mathcal{F}_s\}_{s \geq 0}$  denote the augmented filtration generated by the infinite-dimensional process  $\{U_s\}_{s \geq 0}$ . The latter process is Markov with respect to  $\mathcal{F}$ . Moreover,

$$(2.19) \quad \mathbb{E}[L(2r^6; r + r^3) \mid \mathcal{F}_s] \geq \int_s^{2r^6} \mathbb{P}\{U_\nu^* \leq r + r^3 \mid \mathcal{F}_s\} d\nu \cdot \mathbf{1}_{\{U_s^* \leq r\}}.$$

As in (2.14), if  $\nu > s$  are fixed, then we can write

$$(2.20) \quad \begin{aligned} U_\nu(t) &= U_s(t)e^{-(\nu-s)/2} + \frac{B(e^\nu, t) - B(e^s, t)}{\sqrt{e^\nu - e^s}} \sqrt{1 - e^{-(\nu-s)}} \\ &:= U_s(t)e^{-(\nu-s)/2} + V(t)\sqrt{1 - e^{-(\nu-s)}}. \end{aligned}$$

We emphasize, once again, that  $(U_s, V)$  is a planar Brownian motion. In addition,  $V$  is independent of  $\mathcal{F}_s$ , and  $U_\nu^* \leq U_s^* + V^* \sqrt{1 - \exp\{-(\nu - s)\}}$ . Consequently, as long as  $0 \leq s \leq r^6$  and  $s < \nu < 2r^6$ ,

$$(2.21) \quad U_\nu^* \leq U_s^* + \frac{r^3}{\sqrt{2}} V^*.$$

[We have used the inequality  $1 - e^{-z} \leq z/2$  valid for all  $z \in (0, 1)$ .] Therefore, for all  $0 \leq s \leq r^6$ ,

$$(2.22) \quad \begin{aligned} M(s) &= \mathbb{E} [L(2r^6; r + r^3) \mid \mathcal{F}_s] \\ &\geq \int_s^{2r^6} \mathbb{P} \{V^* \leq \sqrt{2}\} d\nu \cdot \mathbf{1}_{\{U_s^* \leq r\}} \\ &= \mu \{f^* \leq \sqrt{2}\} (2r^6 - s) \cdot \mathbf{1}_{\{U_s^* \leq r\}} \\ &\geq \mu \{f^* \leq \sqrt{2}\} r^6 \cdot \mathbf{1}_{\{U_s^* \leq r\}}. \end{aligned}$$

Because  $\{M(s)\}_{s \geq 0}$  is a martingale, we can apply Doob's maximal inequality to obtain the following:

$$(2.23) \quad \begin{aligned} \mathbb{P} \left\{ \inf_{s \in [0, r^6]} U_s^* \leq r \right\} &\leq \mathbb{P} \left\{ \sup_{s \in [0, r^6]} M(s) \geq \mu \{f^* \leq \sqrt{2}\} r^6 \right\} \\ &\leq \frac{\mathbb{E} [L(2r^6; r + r^3)]}{\mu \{f^* \leq \sqrt{2}\} r^6} = \frac{2\mu \{f^* \leq r + r^3\}}{\mu \{f^* \leq \sqrt{2}\}}. \end{aligned}$$

Thanks to (2.9),

$$(2.24) \quad \frac{\mu \{f^* \leq r + r^3\}}{\mu \{f^* \leq r\}} \sim \exp \left( -\frac{\pi^2}{8} \left[ \frac{1}{(r + r^3)^2} - \frac{1}{r^2} \right] \right) \rightarrow e^{\pi^2/4}. \quad (r \rightarrow 0).$$

Thus, the left-hand side is bounded ( $r \in (0, 1)$ ), and the proposition follows.  $\square$

*Proof of Theorem 2.6: Upper Bound.* Define  $n = n(r)$  to be  $\lfloor r^{-6} \rfloor$ , and define  $I(j; n)$  to be the interval  $[j/n, (j+1)/n)$  ( $j = 0, \dots, n$ ). Then, by stationarity and Proposition 2.9,

$$(2.25) \quad \mathbb{P} \left\{ \inf_{s \in G} U_s^* \leq r \right\} \leq \sum_{\substack{0 \leq j \leq n: \\ I(j; n) \cap G \neq \emptyset}} \mathbb{P} \left\{ \inf_{s \in I(j; n)} U_s^* \leq r \right\} \leq a\mu \{f^* \leq r\} M_n(G),$$

where  $M_n(G) = \#\{0 \leq j \leq n : I(j; n) \cap G \neq \emptyset\}$  defines the *Minkowski content* of  $G$ . In the companion to this paper (2004, Proposition 2.7) we proved that  $M_n(G) \leq 3K_G(1/n)$ . By monotonicity, the latter is at most  $3K_G(r^6)$ , whence the theorem.  $\square$

### 3. PROOF OF THEOREM 1.1 AND COROLLARIES 1.2 AND 1.4

We begin with some preliminary discussions. Define

$$(3.1) \quad \psi_H(G) := \int_1^\infty \frac{K_G(H^6(s))}{sH^2(s)} \exp \left( -\frac{\pi^2}{8H^2(s)} \right) ds, \quad \sigma(r) := \mu \{f^* \leq r\}.$$

Following Erdős (1942), define

$$(3.2) \quad \mathbf{e}_n = e^{n/\ln_+ n}, \quad H_n = H(\mathbf{e}_n) \quad \forall n \geq 1.$$

The “critical” function in (1.11) is  $H^2(t) = \pi^2/(8 \ln_+ \ln_+ t)$ . This, the fact that  $\pi/\sqrt{8} \in (1, 2)$ , and a familiar argument (Erdős, 1942, equations 1.2 and 3.4), together allow us to assume without loss of generality that

$$(3.3) \quad \frac{1}{\sqrt{\ln_+ n}} \leq H_n \leq \frac{2}{\sqrt{\ln_+ n}} \quad \forall n \geq 1.$$

From this we can conclude the existence of a constant  $a > 1$  such that

$$(3.4) \quad \frac{1}{a} H_n^2 \mathbf{e}_{n+1} \leq \mathbf{e}_{n+1} - \mathbf{e}_n \leq a H_{n+1}^2 \mathbf{e}_n \quad \forall n \geq 1.$$

According to our companion work (2004, eq. 2.8), for all  $r > 0$  sufficiently small,

$$(3.5) \quad K_G(\varepsilon) \leq 6K_G(2\varepsilon).$$

Because  $\mathbf{e}_{n+1} \sim \mathbf{e}_n$  as  $n \rightarrow \infty$ , (2.9), (3.4), and (3.5) together imply that

$$(3.6) \quad \sum_{n=1}^{\infty} K_G(H_n^6) \sigma(H_n) < \infty \quad \text{iff} \quad \psi_H(G) < \infty.$$

The following is the key step toward proving Theorem 1.1.

**Proposition 3.1.** *Let  $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  be decreasing and measurable. Then for all non-random Borel sets  $G \subseteq [0, 1]$ ,*

$$(3.7) \quad \liminf_{t \rightarrow \infty} \left( \inf_{s \in G} \sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right) = \begin{cases} +\infty, & \text{if } \psi_H(G) < \infty, \\ -\infty, & \text{if } \psi_H(G) = \infty. \end{cases}$$

First we assume this proposition and derive Theorem 1.1. Then, we will tidy things up by proving the technical Proposition 3.1.

Let us recall (3.1).

**Definition 3.2.** We say that  $\Psi_H(G) < \infty$  if we can decompose  $G$  as  $G = \cup_{n=1}^{\infty} G_n$ —where  $G_1, G_2, \dots$  are closed—such that for all  $n \geq 1$ ,  $\psi_H(G_n) < \infty$ . Else, we say that  $\Psi_H(G) = \infty$ .

Let us first rephrase Theorem 1.1 in the following convenient, and equivalent, form.

**Proposition 3.3.** *Let  $H : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  be decreasing and measurable and  $G \subseteq [0, 1]$  be non-random and Borel. If  $\Psi_H(G) < \infty$ , then*

$$(3.8) \quad \inf_{s \in G} \liminf_{t \rightarrow \infty} \left( \sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right) = \infty \quad \text{P-a.s.}$$

Else, the left-hand side is P-a.s. equal to  $-\infty$ .

*Proof of Theorem 1.1 in the form of Proposition 3.3.* First suppose  $\Psi_H(G)$  is finite. We can write  $G = \cup_{n=1}^{\infty} G_n$ , where the  $G_n$ 's are closed and  $\psi_H(G_n) < \infty$  for all  $n \geq 1$ . Then, according to Proposition 3.1,

$$(3.9) \quad \begin{aligned} & \inf_{s \in G_n} \liminf_{t \rightarrow \infty} \left[ \sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right] \\ & \geq \liminf_{t \rightarrow \infty} \inf_{s \in G_n} \left[ \sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right] = \infty. \end{aligned}$$

This proves that  $\inf_{s \in G} \liminf_{t \rightarrow \infty} (\sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t}) = \infty$  a.s. [P].



For the converse portion suppose  $\Psi_H(G) = \infty$ , and choose arbitrary non-random closed sets  $\{G_n\}_{n=1}^\infty$  such that  $\cup_{n=1}^\infty G_n = G$ . By definition,  $\psi_H(G_n) = \infty$  for some  $n \geq 1$ . Define for all  $T \geq 1$ ,

$$(3.10) \quad \mathcal{S}_T := \left\{ s \in [0, 1] : \inf_{t \geq T} \frac{\sup_{u \in [0, t]} |U_s(u)|}{H(t)\sqrt{t}} \leq 1 \right\}.$$

Evidently,  $\mathcal{S}_T$  is a random set for each  $T \geq 0$ . Moreover, the continuity of the Brownian sheet implies that with probability one,  $\mathcal{S}_T$  is closed for all  $T$ ; hence, so is  $\mathcal{S}_T \cap G_n$ . Because  $\psi_H(G_n) = \infty$ , Proposition 3.1 implies that almost surely,  $\mathcal{S}_T \cap G_n \neq \emptyset$ . Since  $\{\mathcal{S}_T \cap G_n\}_{T=1}^\infty$  is a decreasing sequence of non-void compact sets, they have non-void intersection. That is,  $(\cap_{T=1}^\infty \mathcal{S}_T) \cap G_n \neq \emptyset$  a.s. [P]. Replace  $H$  by  $H - H^3$  to complete the proof of Proposition 3.3.  $\square$

Now we derive Proposition 3.1. This completes our proof of Theorem 1.1. Our proof is divided naturally into two halves.

*Proof of Proposition 3.1: First Half.* Throughout this portion of the proof, we assume that  $\psi_H(G) < \infty$ .

Because  $\mathbf{e}_{n+1} \sim \mathbf{e}_n$  as  $n \rightarrow \infty$ , Theorem 2.6 and Brownian scaling together imply that

$$(3.11) \quad \begin{aligned} \mathbb{P} \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\} &= \mathbb{P} \left\{ \inf_{s \in G} U_s^* \leq H_n \sqrt{\mathbf{e}_n / \mathbf{e}_{n-1}} \right\} \\ &\leq aK_G \left( H_n^6 \left[ \frac{\mathbf{e}_n}{\mathbf{e}_{n-1}} \right]^3 \right) \sigma \left( H_n \sqrt{\frac{\mathbf{e}_n}{\mathbf{e}_{n-1}}} \right). \end{aligned}$$

According to (3.5),  $K_G(\dots) \leq 6K_G(H_n^6)$  for all  $n$  large. This and (3.4) together imply that for all  $n$  large,

$$(3.12) \quad \begin{aligned} \mathbb{P} \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\} \\ \leq aK_G(H_n^6) \sigma \left( H_n \sqrt{1 + AH_n^2} \right) \\ \leq aK_G(H_n^6) \sigma \left( H_n [1 + AH_n^2] \right). \end{aligned}$$

In accord with (2.9), for any fixed  $c \in \mathbf{R}$ ,

$$(3.13) \quad \sigma(r + cr^3) = O(\sigma(r)) \quad (r \rightarrow 0).$$

Thus, for all  $n \geq 1$ ,

$$(3.14) \quad \mathbb{P} \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\} \leq aK_G(H_n^6) \sigma(H_n).$$

Because we are assuming that  $\psi_H(G)$  is finite, (3.6) and the Borel–Cantelli lemma together imply that almost surely,  $\inf_{s \in G} \sup_{u \in [0, \mathbf{e}_{n-1}]} |U_s(u)| > H_n \sqrt{\mathbf{e}_n}$  for all but a finite number of  $n$ 's. It follows from this and a standard monotonicity argument that

$$(3.15) \quad \psi_H(G) < \infty \implies \liminf_{t \rightarrow \infty} \left[ \inf_{s \in G} \sup_{u \in [0, t]} |U_s(u)| - H(t)\sqrt{t} \right] > 0 \text{ a.s. [P].}$$

But if  $\psi_H(G)$  were finite then  $\psi_{H+H^3}(G)$  is also finite; compare (3.5) and (3.13). Thanks to (3.3),  $\lim_{t \rightarrow \infty} H^3(t)\sqrt{t} = \infty$ . Therefore, the lim inf of the preceding display is infinity. This concludes the first half of our proof of Proposition 3.1.  $\square$

In order to prove the second half of Proposition 3.1 we assume that  $\psi_H(G) = \infty$ , recall (3.1), and define

$$(3.16) \quad \begin{aligned} L_n &:= \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_n]} |U_s(u)| \leq H_n \sqrt{\mathbf{e}_n} \right\}, \\ f(z) &:= K_G(z^6) \sigma(z). \end{aligned}$$

**Lemma 3.4.** *Define for all  $j \geq i$ ,  $\lambda_{i,j} := \mathbf{e}_j / (\mathbf{e}_j - \mathbf{e}_i)$  and  $\delta_{i,j} := H_j \sqrt{\lambda_{i,j}} + H_i \sqrt{\lambda_{i,j} - 1}$ . Then, there exists  $a > 1$  such that for all  $j \geq i$ ,  $P(L_j | L_i) \leq a K_G(\delta_{i,j}^6) \sigma(\delta_{i,j})$ .*

*Proof.* Evidently,  $P(L_j | L_i)$  is at most

$$(3.17) \quad \begin{aligned} &P \left\{ \inf_{s \in G} \sup_{u \in [\mathbf{e}_i, \mathbf{e}_j]} |U_s(u)| \leq H_j \sqrt{\mathbf{e}_j} \mid L_i \right\} \\ &= P \left\{ \inf_{s \in G} \sup_{u \in [\mathbf{e}_i, \mathbf{e}_j]} |U_s(u) - U_s(\mathbf{e}_i) + U_s(\mathbf{e}_i)| \leq H_j \sqrt{\mathbf{e}_j} \mid L_i \right\} \\ &\leq P \left\{ \inf_{s \in G} \sup_{u \in [\mathbf{e}_i, \mathbf{e}_j]} |U_s(u) - U_s(\mathbf{e}_i)| \leq H_j \sqrt{\mathbf{e}_j} + H_i \sqrt{\mathbf{e}_i} \right\}. \end{aligned}$$

We have appealed to the Markov properties of the Brownian sheet in the last line. Because  $u \mapsto U_\bullet(u)$  is a  $C(\mathbf{R}_+)$ -valued Brownian motion,

$$(3.18) \quad \begin{aligned} P(L_j | L_i) &\leq P \left\{ \inf_{s \in G} \sup_{u \in [0, \mathbf{e}_j - \mathbf{e}_i]} |U_s(u)| \leq H_j \sqrt{\mathbf{e}_j} + H_i \sqrt{\mathbf{e}_i} \right\} \\ &= P \left\{ \inf_{s \in G} U_s^* \leq \delta_{i,j} \right\}. \end{aligned}$$

Theorem 2.6 completes the proof.  $\square$

Our forthcoming estimates of  $P(L_j | L_i)$  rely on the following elementary bound; see, for example, our earlier work (2003, eq. 8.30): Uniformly for all integers  $j > i$ ,

$$(3.19) \quad \mathbf{e}_j - \mathbf{e}_i \geq \mathbf{e}_i \left( \frac{j-i}{\ln i} \right) (1 + o(1)) \quad (i \rightarrow \infty).$$

**Lemma 3.5.** *There exist  $i_0 \geq 1$  and a finite  $a > 1$  such that for all  $i \geq i_0$  and  $j \geq i + \ln^{19}(j)$ ,*

$$(3.20) \quad P(L_j | L_i) \leq a P(L_j).$$

*Proof.* Thanks to (3.3) and (3.19), the following holds uniformly over all  $j > i + \ln^{19}(j)$ :  $(\mathbf{e}_j / \mathbf{e}_i) \geq (1 + o(1)) H_j^{-36}$  ( $i \rightarrow \infty$ ). Thus, uniformly over all  $j > i + \ln^{19}(j)$ ,

$$(3.21) \quad \begin{aligned} \sqrt{\lambda_{i,j}} &= \frac{1}{\sqrt{1 - (\mathbf{e}_i / \mathbf{e}_j)}} \leq \frac{1}{\sqrt{1 - (1 + o(1)) H_j^{36}}} = 1 + O(H_j^3), \\ H_i \sqrt{\lambda_{i,j} - 1} &= O(H_j^3) \quad (i \rightarrow \infty). \end{aligned}$$

Lemma 3.4 guarantees then that uniformly over all  $j > i + \ln^{19}(j)$ ,  $\delta_{i,j} \leq H_j + O(H_j^3)$ , and the big- $O$  and little- $o$  terms do not depend on the  $j$ 's in question. The lemma follows from this, equations (3.5) and (3.13), and Theorem 2.6.  $\square$

**Lemma 3.6.** *There exist  $i_1 \geq 1$  and  $a \in (0, 1)$  such that for all  $i \geq i_1$  and  $j \in [i + \ln(i), i + \ln^{19}(j)]$ ,  $P(L_j | L_i) \leq (aj^a)^{-1}$ .*

*Proof.* Equations (3.19) and (3.3) together imply that uniformly for all  $j \geq i + \ln(i)$ ,  $(\mathbf{e}_i/\mathbf{e}_j) \leq \frac{1}{2} + o(1)$  ( $i \rightarrow \infty$ ). This is equivalent to the existence of a constant  $A_{3.22}$  such that for all  $(i, j)$  in the range of the lemma,

$$(3.22) \quad \sqrt{\lambda_{i,j}} \vee \sqrt{\lambda_{i,j} - 1} \leq a.$$

Thanks to (3.3), we can enlarge the last constant  $a$ , if necessary, to ensure that for all  $(i, j)$  in the range of this lemma,  $H_i \leq aH_j$ . Therefore, Lemma 3.4 then implies that  $\delta_{i,j} = O(H_j)$ , and the big- $O$  term does not depend on the range of  $j$ 's in question. Because  $G \subseteq [0, 1]$ ,

$$(3.23) \quad K_G(\varepsilon) \leq K_{[0,1]}(\varepsilon) \sim 1/\varepsilon \quad (\varepsilon \rightarrow 0).$$

Thus, Lemma 3.4 ensures that  $P(L_j | L_i) \leq a\delta_{i,j}^{-6}\sigma(\delta_{i,j})$ . Near the origin, the function  $\delta \mapsto \delta^{-6}\sigma(\delta)$  is increasing. Because we have proved that over the range of  $(i, j)$  of this lemma  $\delta_{i,j} = O(H_j)$ , equation (2.9) asserts the existence of a universal  $\alpha > 1$  such that  $P(L_j | L_i)$  is at most  $\alpha H_j^{-6} \exp(-\alpha^{-1}H_j^{-2})$ . Equation (3.3) then completes our proof.  $\square$

**Lemma 3.7.** *There exist  $i_2 \geq 1$  and  $a > 1$  such that for all  $i \geq i_2$  and  $j \in (i, i + \ln i)$ ,  $P(L_j | L_i) \leq ae^{-(j-i)/a}$ .*

*Proof.* By (3.19),  $(\mathbf{e}_i/\mathbf{e}_j) \leq 1 - (1 + o(1))(j - i) \ln^{-1}(i)$  ( $i \rightarrow \infty$ ), where the little- $o$  term does not depend on  $j \in (i, i + \ln i)$ . Similarly,  $(\mathbf{e}_j/\mathbf{e}_i) \geq 1 + (1 + o(1))(j - i) \ln^{-1}(i)$ . Thus, as  $i \rightarrow \infty$ ,

$$(3.24) \quad \begin{aligned} \sqrt{\lambda_{i,j}} &= \frac{1}{\sqrt{1 - (\mathbf{e}_i/\mathbf{e}_j)}} \leq (1 + o(1)) \sqrt{\frac{\ln i}{j - i}} \leq \frac{2 + o(1)}{H_j \sqrt{j - i}}, \\ \sqrt{\lambda_{i,j} - 1} &= \frac{1}{\sqrt{(\mathbf{e}_j/\mathbf{e}_i) - 1}} \leq (1 + o(1)) \sqrt{\frac{\ln i}{j - i}} \leq \frac{2 + o(1)}{H_j \sqrt{j - i}}, \end{aligned}$$

by (3.3). Once again, the little- $o$  terms are all independent of  $j \in (i, i + \ln i)$ . Because  $H_i = O(H_j)$  uniformly for all  $(i, j)$  in the range considered here, Lemma 3.4 implies that uniformly for all  $j \in (i, i + \ln i)$ ,  $\delta_{i,j} = O(1/\sqrt{j - i})$ . Equation (3.23) bounds the first term on the right-hand side; (2.9) bounds the second. This and (3.3) together prove the existence of a constant  $\alpha > 1$  such that for all  $i \geq i_2$  and all  $j \in (i, i + \ln i)$ ,  $P(L_j | L_i) \leq \alpha(j - i)^3 \exp\{-(j - i)/\alpha\}$ . The lemma follows.  $\square$

*Proof of Proposition 3.1: Second Half.* According to Theorem 2.6, for all  $n$  large enough,  $P(L_n) \geq af(H_n)$ . Because  $\psi_H(G) = \infty$ , the latter estimate and (3.6) together imply that

$$(3.25) \quad \sum_{i=1}^{\infty} P(L_i) = \infty.$$

Thus, our derivation is complete once we demonstrate the following:

$$(3.26) \quad \liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^{n-1} \sum_{j=i}^n \mathbb{P}(L_i \cap L_j)}{(\sum_{i=1}^n \mathbb{P}(L_i))^2} < \infty.$$

See Chung and Erdős (1952). In fact, the preceding display holds with a lim sup in place of the lim inf. This fact follows from combining, using standard arguments, Lemmas 3.5 through 3.7.

Indeed, let  $I := \max(3, i_1, i_2, i_3)$  and  $s_n := \sum_{i=1}^n \mathbb{P}(L_i)$ . Lemma 3.5 ensures that

$$(3.27) \quad \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j > i + \ln^{19}(j)}}^n \mathbb{P}(L_j \cap L_i) = O(s_n^2).$$

By Lemma 3.6,

$$(3.28) \quad \begin{aligned} \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j \in (i + \ln(i), i + \ln^{19}(j)]}}^n \mathbb{P}(L_j \cap L_i) &\leq \frac{1}{a} \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j \in (i + \ln(i), i + \ln^{19}(j)]}}^n j^{-a} \mathbb{P}(L_i) \\ &= \sum_{i=I}^n O\left(\frac{\ln^{19}(i)}{i^a}\right) \mathbb{P}(L_i) = O(s_n). \end{aligned}$$

The big- $O$  terms do not depend on the variables  $(j, n)$ .

Finally, Lemma 3.7 implies that

$$(3.29) \quad \sum_{i=I}^{n-1} \sum_{\substack{j=i \\ j \in (i, i + \ln i]}}^n \mathbb{P}(L_j \cap L_i) \leq a \sum_{i=I}^n \sum_{j=i}^{\infty} \mathbb{P}(L_i) e^{(j-i)/a} = O(s_n).$$

We have already seen that  $s_n \rightarrow \infty$ . Thus, (3.27)–(3.29) imply (3.26), and hence the theorem. More precisely, we have proved so far that

$$(3.30) \quad \psi_H(G) = \infty \implies \liminf_{t \rightarrow \infty} \left[ \inf_{s \in G} \sup_{u \in [0, t]} |U_s(u)| - H(t) \sqrt{t} \right] < 0 \text{ a.s. } [\mathbb{P}].$$

Replace  $H$  by  $H + H^3$  to deduce that the preceding lim inf is in fact  $-\infty$ . This completes our proof of Proposition 3.1.  $\square$

We conclude this section by proving the remaining Corollaries 1.2 and 1.4.

*Proof of Corollary 1.2.* By definition,  $\mathcal{L}(H)$  holds q.s. iff  $\text{cap}_{\mathbf{R}_+}((\mathcal{L}(H))^{\mathfrak{c}}) = 0$ . Thanks to Theorem 1.1, this condition is equivalent to the existence of a non-random “closed-denumerable” decomposition  $\mathbf{R}_+ = \cup_{n=1}^{\infty} G_n$  such that for all  $n \geq 1$ ,  $\psi_H(G_n) < \infty$ . But one of the  $G_n$ ’s must contain a closed interval that has positive length. Therefore, by the translation-invariance of  $G \mapsto K_G(r)$ , there exists  $\varepsilon \in (0, 1)$  such that  $\psi_H([0, \varepsilon]) < \infty$ .

Conversely, if  $\psi_H([0, \varepsilon])$  is finite, then we can define  $G_n$  to be  $[(n-1)\varepsilon, n\varepsilon]$  ( $n \geq 1$ ) to find that  $\psi_H(G_n) = \psi_H([0, \varepsilon]) < \infty$ . Theorem 1.1 then proves that  $\text{cap}_{\mathbf{R}_+}((\mathcal{L}(H))^{\mathfrak{c}}) = 0$  iff there exists  $\varepsilon > 0$  such that  $\psi_H([0, \varepsilon]) < \infty$ . Because  $K_{[0, \varepsilon]}(r) \sim \varepsilon/r$  ( $r \rightarrow 0$ ), the corollary follows.  $\square$

*Proof of Corollary 1.4.* We can change variables to deduce that  $\psi_{H_\nu}(G)$  is finite iff  $\int_1^\infty K_G(1/s)s^{-1-(\nu/3)} ds$  converges. This and Proposition 2.8 of our companion work (2004) together imply that

$$(3.31) \quad \inf\{\nu > 0 : \psi_{H_\nu}(G) < \infty\} = 2 + 3\overline{\dim}_{\mathcal{M}} G,$$

where  $\overline{\dim}_{\mathcal{M}}$  denotes the (upper) Minkowski dimension (Mattila, 1995). By regularization (Mattila, 1995, p. 81),

$$(3.32) \quad \inf\{\nu > 0 : \Psi_{H_\nu}(G) < \infty\} = 2 + 3 \dim_{\mathcal{D}} G.$$

Theorem 1.1 now implies Corollary 1.4.  $\square$

#### REFERENCES

- Chung, Kai Lai. 1948. *On the maximum partial sums of sequences of independent random variables*, Trans. Amer. Math. Soc. **64**, 205–233.
- Chung, K. L. and P. Erdős. 1952. *On the application of the Borel-Cantelli lemma*, Trans. Amer. Math. Soc. **72**, 179–186.
- Dudley, R. M. 1973. *A Course in Empirical Processes*, École d'été de probabilités Saint-Flour, XII–1982, Springer, Berlin, pp. 1–142.
- Erdős, Paul. 1942. *On the law of the iterated logarithm*, Ann. Math. **43(2)**, 419–436.
- Fukushima, Masatoshi. 1984. *Basic properties of Brownian motion and a capacity on the Wiener space*, J. Math. Soc. Japan **36(1)**, 161–176.
- Khintchine, A. Ya. 1933. *Asymptotische Gesetz der Wahrscheinlichkeitsrechnung*, Springer, Berlin.
- Khoshnevisan, Davar. 2002. *Multiparameter Processes: An Introduction to Random Fields*, Springer, New York.
- Khoshnevisan, Davar, David A. Levin, and Pedro J. Méndez–Hernández. 2003. *On dynamical Gaussian random walks*, Ann. Probab. (To appear).
- . 2004. *Exceptional times and invariance for dynamical random walks* (Preprint).
- Lifshits, M. A. and Z. Shi. 2003. *Lower functions of an empirical process and of a Brownian sheet*, Teor. Veroyatnost. i Primenen. **48(2)**, 321–339. (Russian, with Russian summary)
- Malliavin, Paul. 1979. *Régularité de lois conditionnelles et calcul des variations stochastiques*, C.R. Acad. Sci. Paris, Sér. A-B **289**.
- Mattila, Pertti. 1995. *Geometry of Sets and Measures in Euclidean Spaces: Fractals and Rectifiability*, Cambridge University Press, Cambridge.
- Meyer, P.-A. 1982. *Note sur les processus d'Ornstein–Uhlenbeck (Appendice: Un resultat de D. Williams)*, Sémin. de Probab. XVI, Lec. Notes in Math., vol. 920, Springer, pp. 95–133.
- Mountford, T. S. 1992. *Quasi-everywhere upper functions*, Sémin. de Probab. XXVI, Lect. Notes in Math., vol. 1526, Springer, pp. 95–106.
- Tihomirov, V. M. 1963. *The works of A. N. Kolmogorov on  $\varepsilon$ -entropy of function classes and superpositions of functions*, Uspehi Mat. Nauk **18(5 (113))**, 55–92.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF UTAH, 155 S. 1400 E., SALT LAKE CITY, UT 84112–0090

*E-mail address:* [davar@math.utah.edu](mailto:davar@math.utah.edu)

*URL:* <http://www.math.utah.edu/~davar>

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF UTAH, 155 S. 1400 E., SALT LAKE CITY, UT 84112–0090

*E-mail address:* [levin@math.utah.edu](mailto:levin@math.utah.edu)

*URL:* <http://www.math.utah.edu/~levin>

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF UTAH, 155 S. 1400 E., SALT LAKE CITY, UT 84112–0090

*Current address:* Escuela de Matematica, Universidad de Costa Rica, San Pedro de Montes de Oca, Costa Rica

*E-mail address:* [mendez@math.utah.edu](mailto:mendez@math.utah.edu), [pmendez@emate.ucr.ac.cr](mailto:pmendez@emate.ucr.ac.cr)

*URL:* <http://www.math.utah.edu/~mendez>