

Probability is nothing but common sense reduced to calculation.

Probability is nothing but common sense reduced to calculation.

Pierre Simon Laplace

Probability is nothing but common sense reduced to calculation.

Pierre Simon Laplace



Last time we started to talk about probability as ratio of areas.

Last time we started to talk about probability as ratio of areas. For example, the probability that you would hit a section of a dartboard at random is the ratio of the area of that section to to the total area of the dartboard.

Last time we started to talk about probability as ratio of areas. For example, the probability that you would hit a section of a dartboard at random is the ratio of the area of that section to the total area of the dartboard. But areas can “abstractly” represent probabilities as well. We start with a fun example before moving on to a more standard ones.

Example 1. *You and your housemates have a two-bathroom house, but only one hot shower can be taken at a time. You have one housemate whose first class is at the same time as yours, and unfortunately you both like to take ten-minutes showers and start your day in the same half-hour window.*

Example 1. *You and your housemates have a two-bathroom house, but only one hot shower can be taken at a time. You have one housemate whose first class is at the same time as yours, and unfortunately you both like to take ten-minutes showers and start your day in the same half-hour window. What are the chances that you get to have an uninterrupted hot shower?*

Continuous random variables

Continuous random variables

Let's first step back and review and expand our language as developed for probability within the context of there being a finite number of outcomes; we will see that this language applies when the outcomes can vary continuously.

Continuous random variables

Let's first step back and review and expand our language as developed for probability within the context of there being a finite number of outcomes; we will see that this language applies when the outcomes can vary continuously.

Definition 2. • *A random phenomenon is one which is unpredictable but follows a regular distribution after large numbers of repetitions.*

Continuous random variables

Let's first step back and review and expand our language as developed for probability within the context of there being a finite number of outcomes; we will see that this language applies when the outcomes can vary continuously.

Definition 2. • *A random phenomenon is one which is unpredictable but follows a regular distribution after large numbers of repetitions.*

• *A random variable X represents the outcome of a*

random phenomenon.

random phenomenon.

- *The sample space of a random variable is the set of possible values for the random variable.*

random phenomenon.

- *The sample space of a random variable is the set of possible values for the random variable.*
- *The probability distribution of a random variable is the assignment of probabilities to the values in the sample space.*

random phenomenon.

- *The sample space of a random variable is the set of possible values for the random variable.*
- *The probability distribution of a random variable is the assignment of probabilities to the values in the sample space.*

Example 3. • *Our random phenomenon is throwing two dice.*

Example 3. • *Our random phenomenon is throwing two dice.*

- *Our random variable X will be the total shown on the two dice.*

- Example 3.**
- *Our random phenomenon is throwing two dice.*
 - *Our random variable X will be the total shown on the two dice.*
 - *Our sample space is $\{2, 3, 4, 5, \dots, 12\}$.*

Example 3. • *Our random phenomenon is throwing two dice.*

- *Our random variable X will be the total shown on the two dice.*
- *Our sample space is $\{2, 3, 4, 5, \dots, 12\}$.*
- *Our probability distribution is*

| X | 2 | 3 | 4 | 5 | 6 | 7 | 8 | |
|--------------------|------|------|------|------|------|------|------|----|
| <i>Probability</i> | .028 | .056 | .083 | .111 | .139 | .167 | .139 | .1 |

Example 3. • *Our random phenomenon is throwing two dice.*

- *Our random variable X will be the total shown on the two dice.*
- *Our sample space is $\{2, 3, 4, 5, \dots, 12\}$.*
- *Our probability distribution is*

| X | 2 | 3 | 4 | 5 | 6 | 7 | 8 | |
|--------------------|------|------|------|------|------|------|------|----|
| <i>Probability</i> | .028 | .056 | .083 | .111 | .139 | .167 | .139 | .1 |
| | | | | | | | | |

So, for example, a “7” is six times as likely as a “2” or a “12”.

So, for example, a “7” is six times as likely as a “2” or a “12”.

Definition 4. *A random variable is discrete if the sample space is a finite number of values. It is continuous if the sample space is a range of values.*

So, for example, a “7” is six times as likely as a “2” or a “12”.

Definition 4. *A random variable is discrete if the sample space is a finite number of values. It is continuous if the sample space is a range of values.*

The example above is a discrete random variable – why?

Example 5. *Let X be a randomly chosen number between 0 and 3.*

Example 5. *Let X be a randomly chosen number between 0 and 3.*

- *What is our random event?*

Example 5. *Let X be a randomly chosen number between 0 and 3.*

- *What is our random event? our random variable?*

Example 5. *Let X be a randomly chosen number between 0 and 3.*

- *What is our random event? our random variable?*
- *What is $P(0 \leq X \leq 3)$?*

Example 5. *Let X be a randomly chosen number between 0 and 3.*

- *What is our random event? our random variable?*
- *What is $P(0 \leq X \leq 3)$? $P(1.5 \leq X \leq 2)$?*

Example 5. *Let X be a randomly chosen number between 0 and 3.*

- *What is our random event? our random variable?*
- *What is $P(0 \leq X \leq 3)$? $P(1.5 \leq X \leq 2)$?*
- *What is $P(1 \leq X \leq 1.4 \text{ or } 2 \leq X \leq 2.5)$?*

Example 5. *Let X be a randomly chosen number between 0 and 3.*

- *What is our random event? our random variable?*
- *What is $P(0 \leq X \leq 3)$? $P(1.5 \leq X \leq 2)$?*
- *What is $P(1 \leq X \leq 1.4 \text{ or } 2 \leq X \leq 2.5)$?*

In this example X is called uniformly distributed between 0 and 3.

We can redo this example using a uniform probability density function.

We can redo this example using a uniform probability density function.

Definition 6. *1. A distribution for a continuous random variable can be any non-negative function D where the total area under the function is 1.*

We can redo this example using a uniform probability density function.

- Definition 6.** 1. *A distribution for a continuous random variable can be any non-negative function D where the total area under the function is 1.*
2. *If X is distributed according to a distribution D then $P(a \leq X \leq b)$ is the area between the line $x = a$, the line $x = b$, the graph of D , and the x -axis.*

We can redo this example using a uniform probability density function.

- Definition 6.** *1. A distribution for a continuous random variable can be any non-negative function D where the total area under the function is 1.*
- 2. If X is distributed according to a distribution D then $P(a \leq X \leq b)$ is the area between the line $x = a$, the line $x = b$, the graph of D , and the x -axis.*
- 3. The sample space for such a variable is all x where D is non-zero.*

We can redo this example using a uniform probability density function.

- Definition 6.** *1. A distribution for a continuous random variable can be any non-negative function D where the total area under the function is 1.*
- 2. If X is distributed according to a distribution D then $P(a \leq X \leq b)$ is the area between the line $x = a$, the line $x = b$, the graph of D , and the x -axis.*
- 3. The sample space for such a variable is all x where D is non-zero.*

4. *If D is equal to either 0 or one other fixed number, we say D is a uniform distribution.*

4. If D is equal to either 0 or one other fixed number, we say D is a uniform distribution.

Example 7. *Revisit the previous example.*

- *Our sample space is the closed interval $[0, 3]$.*

4. If D is equal to either 0 or one other fixed number, we say D is a uniform distribution.

Example 7. *Revisit the previous example.*

- *Our sample space is the closed interval $[0, 3]$.*
- *Our probability distribution is the function*

$$p(t) = \begin{cases} 1/3 & 0 \leq t \leq 3 \\ 0 & \text{else} \end{cases}$$

4. If D is equal to either 0 or one other fixed number, we say D is a uniform distribution.

Example 7. *Revisit the previous example.*

- *Our sample space is the closed interval $[0, 3]$.*
- *Our probability distribution is the function*

$$p(t) = \begin{cases} 1/3 & 0 \leq t \leq 3 \\ 0 & \text{else} \end{cases}$$

Example 8. *Suppose that X is distributed according to*

$$D = \begin{cases} \frac{1}{12} & 1 \leq x \leq 3 \\ \frac{1}{2} & 3 \leq x \leq 4 \\ \frac{1}{9} & 4 \leq x \leq 7 \\ 0 & \text{otherwise} \end{cases}$$

- *What is the state space?*

Example 8. Suppose that X is distributed according to

$$D = \begin{cases} \frac{1}{12} & 1 \leq x \leq 3 \\ \frac{1}{2} & 3 \leq x \leq 4 \\ \frac{1}{9} & 4 \leq x \leq 7 \\ 0 & \text{otherwise} \end{cases}$$

- What is the state space?
- What is $P(2 \leq X \leq 3)$?

Example 8. Suppose that X is distributed according to

$$D = \begin{cases} \frac{1}{12} & 1 \leq x \leq 3 \\ \frac{1}{2} & 3 \leq x \leq 4 \\ \frac{1}{9} & 4 \leq x \leq 7 \\ 0 & \text{otherwise} \end{cases}$$

- What is the state space?
- What is $P(2 \leq X \leq 3)$?
- Which is more likely, that X is between 3 and 4 or that

X is between 5 and 7?

X is between 5 and 7?

Compare the answer to the last question to the answer you would get if X were uniformly distributed.

X is between 5 and 7?

Compare the answer to the last question to the answer you would get if X were uniformly distributed.

Normal distributions

X is between 5 and 7?

Compare the answer to the last question to the answer you would get if X were uniformly distributed.

Normal distributions

Example 9. *[Main Example] Z is a number chosen with normal distribution $N(\mu, \sigma)$.*

X is between 5 and 7?

Compare the answer to the last question to the answer you would get if X were uniformly distributed.

Normal distributions

Example 9. *[Main Example] Z is a number chosen with normal distribution $N(\mu, \sigma)$. We'll use $N(0, 1)$ as our first example.*

X is between 5 and 7?

Compare the answer to the last question to the answer you would get if X were uniformly distributed.

Normal distributions

Example 9. *[Main Example] Z is a number chosen with normal distribution $N(\mu, \sigma)$. We'll use $N(0, 1)$ as our first example.*

- *Our random event is the choice of a number.*

X is between 5 and 7?

Compare the answer to the last question to the answer you would get if X were uniformly distributed.

Normal distributions

Example 9. *[Main Example] Z is a number chosen with normal distribution $N(\mu, \sigma)$. We'll use $N(0, 1)$ as our first example.*

- *Our random event is the choice of a number.*
- *Our random variable Z is the value of the number.*

X is between 5 and 7?

Compare the answer to the last question to the answer you would get if X were uniformly distributed.

Normal distributions

Example 9. *[Main Example] Z is a number chosen with normal distribution $N(\mu, \sigma)$. We'll use $N(0, 1)$ as our first example.*

- *Our random event is the choice of a number.*
- *Our random variable Z is the value of the number.*

- *Our sample space is all possible real numbers.*

- *Our sample space is all possible real numbers.*
- *Our probability distribution is the function*

$$e^{-t^2/2} / \sqrt{2\pi}.$$

- *Our sample space is all possible real numbers.*
- *Our probability distribution is the function*

$$e^{-t^2/2} / \sqrt{2\pi}.$$

- *What is $P(\mathbf{Z} > 3)$? (Hint: use 68-95-99.7 rule.)*

- *Our sample space is all possible real numbers.*
- *Our probability distribution is the function*

$$e^{-t^2/2} / \sqrt{2\pi}.$$

- *What is $P(\mathbf{Z} > 3)$? (Hint: use 68-95-99.7 rule.)*
- *What is $P(0 \leq Z \leq 2)$?*

- *Our sample space is all possible real numbers.*
- *Our probability distribution is the function*

$$e^{-t^2/2} / \sqrt{2\pi}.$$

- *What is $P(\mathbf{Z} > 3)$? (Hint: use 68-95-99.7 rule.)*
- *What is $P(0 \leq Z \leq 2)$?*
- *$P(\mathbf{Z} > 100)$?*

- *Our sample space is all possible real numbers.*
- *Our probability distribution is the function*

$$e^{-t^2/2} / \sqrt{2\pi}.$$

- *What is $P(\mathbf{Z} > 3)$? (Hint: use 68-95-99.7 rule.)*
- *What is $P(0 \leq Z \leq 2)$?*
- *$P(\mathbf{Z} > 100)$?*

Doing calculations with normal distributions works much as before.

Doing calculations with normal distributions works much as before. Instead of calculating percentiles, we are calculating probabilities.

Doing calculations with normal distributions works much as before. Instead of calculating percentiles, we are calculating probabilities.