


**MATH 634 HOMEWORK 8**  
**DUE NOVEMBER 14, 2018.**

INSTRUCTOR: ROBERT LIPSHITZ

Required problems, to turn in:


- (1) Give abstract simplicial complexes whose geometric realizations are the following simplicial complexes. Compute the simplicial homology and cohomology groups of these complexes. (Note: the shaded triangles are 2-simplices.)



- (2) Recall that  $\delta = \delta_n: C^n(X) \rightarrow C^{n+1}(X)$  is the coboundary map on the simplicial cochain complex. Prove that  $\delta^2 = 0$  or, more precisely,  $\delta_{n+1} \circ \delta_n = 0$  for all  $n \geq 0$ .
- (3) Find a simplicial complex homeomorphic to  $\mathbb{R}P^2$ .
- (4) Find a  $\Delta$ -complex homeomorphic to the Klein bottle and use it to compute the simplicial homology and cohomology of the Klein bottle.
- (5) Let  $X_\bullet$  be an abstract simplicial complex and  $v_0, \dots, v_n \in X_0$  be vertices of  $X_\bullet$  and suppose that there is a  $0 \leq k \leq n$  so that  $\{v_0, \dots, \widehat{v}_k, \dots, v_n\} \notin X_{n-1}$  but  $\{v_0, \dots, \widehat{v}_i, \dots, v_n\} \in X_{n-1}$  for all  $i \neq k$ . Then we can form a new simplicial complex  $Y_\bullet$  with
- (a)  $Y_m = X_m$  for  $m \notin \{n-1, n\}$ ,
  - (b)  $Y_{n-1} = X_{n-1} \cup \{\{v_0, \dots, \widehat{v}_k, \dots, v_n\}\}$ , and
  - (c)  $Y_n = X_n \cup \{\{v_0, \dots, v_k, \dots, v_n\}\}$ .

(In the special case  $n = 1$ ,  $Y_\bullet$  is obtained by adding a single new vertex  $w$  to  $X_0$  and an edge from some vertex  $v_0$  to  $w$ .) We say that  $Y_\bullet$  is obtained from  $X_\bullet$  by an *n-dimensional elementary expansion* and  $X_\bullet$  is obtained from  $Y_\bullet$  by an *n-dimensional elementary collapse*. More generally, given abstract simplicial complexes  $X_\bullet$  and  $Y_\bullet$ , we say that  $X_\bullet$  is *simple homotopy equivalent* to  $Y_\bullet$  if you can get from  $X_\bullet$  to  $Y_\bullet$  by a sequence of elementary expansions and elementary collapses of any dimension (and re-ordering vertices).

For example, here is a simple homotopy equivalence between two complexes:



- (a) Sketch a proof that if  $X_\bullet$  is simple homotopy equivalent to  $Y_\bullet$  then their geometric realizations are homotopy equivalent. (The converse is false.)
- (b) If  $Y_\bullet$  is obtained from  $X_\bullet$  by an elementary expansion, there is an inclusion map  $i_m: C_m(X_\bullet) \hookrightarrow C_m(Y_\bullet)$ . Construct a quotient map  $q_m: C_m(Y_\bullet) \rightarrow C_m(X_\bullet)$  so that
- the  $q_m$  form a chain map,
  - $q_m \circ i_m = \mathbb{I}_{C_m(X_\bullet)}$  and
  - there are maps  $h_m: C_m(Y_\bullet) \rightarrow C_{m+1}(Y_\bullet)$  with

$$i_m \circ q_m - \mathbb{I}_{C_m(Y_\bullet)} = \partial \circ h_m + h_{m-1} \circ \partial.$$

(Hint: for most simplices  $\sigma$ , you will probably define  $h_m(\sigma) = 0$ .)

To save time, it's okay if you only consider the case of an  $n$ -dimensional elementary expansion for  $n > 1$ : the 1-dimensional case is similar, but maybe the notation is a bit different.

- (c) Conclude that  $H_m(Y_\bullet) \cong H_m(X_\bullet)$  for each  $m$  and, consequently, that simple homotopy equivalent simplicial complexes have isomorphic homology groups.

To think about but not turn in:

- Given a simplicial map  $f: X \rightarrow Y$ , define an induced map  $f^*: H^n(Y) \rightarrow H^n(X)$  and prove your map is well-defined. Prove also that  $\mathbb{I}^* = \mathbb{I}$  and if  $g: Y \rightarrow Z$  then  $(g \circ f)^* = f^* \circ g^*$ .
- Give a simplicial complex homeomorphic to  $\mathbb{R}P^3$  and compute its simplicial homology and cohomology.
- Give a simplicial complex homeomorphic to the  $n$ -sphere and compute its simplicial homology and cohomology groups.

*E-mail address:* lipshitz@uoregon.edu