

MATH 636 HOMEWORK 3
DUE APRIL 16, 2021.

INSTRUCTOR: ROBERT LIPSHITZ

- (1) Hatcher 4.1.4 (p. 358).
- (2) Hatcher 4.1.11 (pp. 358–359).
- (3) (a) Suppose that M is a closed, connected m -manifold. Suppose further that M is *triangulated*, i.e., is an m -dimensional simplicial complex. Suppose $p \in M$ is on a codimension-1 face (or facet) $\partial_i\sigma$ of some simplex σ . Show that p is on a codimension-1 face of exactly two simplices σ, σ' .
- (b) With notation as in the previous part, let $\alpha = \sum \sigma_i \in C_m(M; \mathbb{F}_2)$ be the sum of the m -simplices in M , viewed, via their characteristic maps, as maps $\sigma_i: \Delta^m \rightarrow M$. Show that α is a cycle.
- (c) Show that for any point p in the interior of some m -simplex σ_i , the image of α in $H_m(M, M \setminus \{p\}; \mathbb{F}_2)$ is a generator. Deduce that α is the generator of $H_m(M; \mathbb{F}_2) \cong \mathbb{F}_2$ so α is a (in fact, the) mod-2 fundamental class for M .
- (d) Now, suppose that $M^m \subset N^n$ is a closed submanifold of a manifold N , and that N is triangulated in such a way that $M \subset N$ is a subcomplex. Show that $i_*[M] \in H_m(N; \mathbb{F}_2)$, the homology class represented by M , is the sum of the m -simplices in N which are contained in M . (Hint: this is easy.)

Suggested review / qualifying exam practice (not to turn in):

- (1) Hatcher 4.1.3, 4.1.5, 4.1.8, 4.1.12, 4.1.13, 4.1.14.
- (2) In class, we showed that if $f: Y \rightarrow Z$ is a weak homotopy equivalence then for any CW complex X , $f_*: [X, Y] \rightarrow [X, Z]$ is injective. We also sketched a proof that f_* is surjective. Fill in the details of that proof.
- (3) With notation as in the previous problem, fill in the proof that the map $f_*: [(X, x_0), (Y, y_0)] \rightarrow [(X, x_0), (Z, z_0)]$ of based homotopy classes of maps is bijective.

More problems to think about but not turn in:

- (1) Hatcher 4.1.10, 4.1.18.
- (2) Extend Problem 3 to the case that instead of the manifold M being triangulated, M is an n -dimensional CW complex and $\alpha \in C_n^{\text{cell}}(M; \mathbb{F}_2)$ is the sum of the n -cells in M . (You'll have to follow α through the isomorphism between cellular and singular homology.)
- (3) Extend Problem 3 to \mathbb{Z} -coefficients.

Email address: lipshitz@uoregon.edu