

**MATH 635 HOMEWORK 2**  
**DUE JANUARY 23, 2019.**

INSTRUCTOR: ROBERT LIPSHITZ

Reminder: homework is due at the *beginning* of class, handed to me or in my mailbox on the second floor of Fenton.

- (1) Hatcher 2.2.22 (p. 157).
- (2) Hatcher 2.2.23 (p. 157). (This is a special case of the Riemann-Hurwitz Formula (over  $\mathbb{C}$ ), which includes branched covers. The proof of the Riemann-Hurwitz formula is not any harder.)
- (3) Hatcher 3.1.11 (p. 205).
- (4) Give an example of a space  $X$  so that:
  - (a) For any field  $k$ ,  $\dim(H^*(X; k)) \leq 5$ .
  - (b)  $H_*(X; \mathbb{Z})$  is not finitely generated.
- (5) Hatcher 3.A.1 (p. 267).

Review / qualifying exam practice (not to turn in):

- (1) Hatcher 2.2.20, 2.2.21, 2.2.25, 3.1.1.

More problems to think about but not turn in:

- (1) Read the rest of the problems in Sections 2.2, 3.1, and 3.A and solve any that seem hard.
- (2) If you're new to Ext and Tor, solve Hatcher 3.1.1, 3.1.2, 3.1.3, 3.A.2,
- (3) Check that every time I have asked you to compute both homology and cohomology, the answers you got are consistent with the universal coefficient theorem.
- (4) Here is a more abstract proof of the universal coefficient theorem; fill in the details.
  - (a) Let  $C_*$  be a chain complex of free abelian groups, and  $H_*$  the homology of  $C_*$ , which we view as a chain complex with trivial homology. Assume that  $C_i = 0$  for all sufficiently small  $i$  (i.e.,  $C_*$  is *bounded below*). Show that there is a chain map  $C_* \rightarrow H_*$  inducing an isomorphism on homology (i.e., a *quasi-isomorphism* from  $C_*$  to  $H_*$ ).
  - (b) Given a bounded-below chain complex  $C_*$  over a ring  $R$ , choose a chain complex  $P_*$  of free  $R$ -modules which is quasi-isomorphic to  $C_*$ . For  $M$  an  $R$ -module, define  $\text{Tor}^i(C_*, M)$  to be the  $i^{\text{th}}$  homology group of  $P_* \otimes M$ . Prove that:
    - (i)  $\text{Tor}^i(C_*, M)$  is independent of the choice of  $P_*$ , up to (canonical) isomorphism.
    - (ii) If  $C_*$  is quasi-isomorphic to  $D_*$  then  $\text{Tor}^i(C_*, M) \cong \text{Tor}^i(D_*, M)$ .
  - (c) If the differential on  $C_*$  vanishes, then

$$\text{Tor}^i(C_*, M) \cong (C_i \otimes M) \oplus \text{Tor}^1(C_{i-1}, M) \oplus \text{Tor}^2(C_{i-2}, M) \oplus \cdots$$

- (d) By definition, if  $C_*$  is a chain complex of free  $R$ -modules then  $\text{Tor}^i(C_*, M) \cong H_i(C_* \otimes M) =: H_i(C_*; M)$ . So, deduce the universal coefficient theorem for homology.
- (5) Prove the universal coefficient theorem with  $\mathbb{Z}$  replaced by any PID.
- (6) Look up and prove the Riemann-Hurwitz formula.

*E-mail address:* lipshitz@uoregon.edu