



1. (1 point.) The Squeeze Theorem was invented for the purpose of torturing which of:

1. Functions.
2. Calculus students.
3. Professors who have to teach it in calculus classes.

2. (6 points) State carefully the definition of the derivative of a function.

3. (a) (9 points) If  $f(x) = x^2 + 5$ , compute the derivative  $f'(3)$  *directly from the definition of the derivative* (which you are supposed to have given above). (No credit will be given for just using the differentiation rules, but see Part (b).)

(b) (1 point) Use the differentiation rules we have learned to check your answer to part (a).

4. (10 points) Differentiate the function  $y(x) = (4x^3 + 8x) \sin(x) + \frac{1}{5}$ . (You need not do this directly from the definition.)

5. (10 points) Differentiate the function  $h(x) = \cos(3x^2 - 16x)$ . (You need not do this directly from the definition.)

6. (10 points) Find the exact value of the limit  $\lim_{x \rightarrow -4} \frac{x^2 - 16}{x^2 + x - 12}$ , or explain why this limit does not exist.

7. (20 points.) An eccentric gardener wants to build a rectangular walled enclosure. The north and south walls will cost 2 pesos per foot, the west wall will cost 5 pesos per foot, and the east wall will cost 3 pesos per foot. The gardener has 800 pesos available to build the enclosure. Use calculus to find the lengths of the south and west walls of the largest such enclosure that can be built. Include units. Verify that the solution corresponds to the largest possible area, using methods we have seen so far in the course (not the second derivative test).

8. (13 points) Use the methods of calculus to find the exact values of  $x$  at which the function  $f(x) = x^3 - 6x^2 + 16$  takes its absolute minimum and maximum on the interval  $[-1, 2]$ .

(No credit will be given for correct guesses without supporting work that is valid for general functions of the sort considered in this course.)

9. (10 points.) This problem is about using correct notation. Accordingly, almost all the credit is for correctness of notation.


Consider the problem of finding the exact value of  $\lim_{x \rightarrow -5} \frac{x^3 + 5x^2 + 4x + 20}{x + 5}$ . The method is to factor the numerator and cancel one of the factors. The factors of the numerator are  $x + 5$  and  $x^2 + 4$ .

Write out the calculation in full, in correct notation which exhibits correctly the steps of the calculation. In particular, put “=” and “lim” everywhere they belong, and nowhere else. Start by writing  $\lim_{x \rightarrow -5} \frac{x^3 + 5x^2 + 4x + 20}{x + 5}$ . Show at least the following steps:

- After factoring but before cancellation.
- After cancellation but before substituting  $x = -5$ .
- After substituting  $x = -5$  but before possible simplification.
- The simplified final result, if the result in the previous step can be simplified.

(Continued on back or on next page.)

10. (5 points/part) For the function  $y = g(x)$  graphed below, answer the following questions:



(a) Which of the following best describes  $g'(4)$ ? Why?

- (1)  $g'(4)$  does not exist.
- (2)  $g'(4)$  is close to 0.
- (3)  $g'(4)$  is positive and not close to 0.
- (4)  $g'(4)$  is negative and not close to 0.
- (5) None of the above.

(b) Does  $\lim_{x \rightarrow -5} g(x)$  exist? If so, what is it? If not, why not?

Extra credit. (15 extra credit points; grading will be harsher than on related problems on the main exam. Do not attempt this problem until you have done and checked your answer to all the ordinary problems on this exam. It will only be counted if you get a grade of B or better on the main part of this exam.)

Let  $f$  be a function such that  $f'(t) = \sqrt[3]{7 + \arctan(t)}$ . Let

$$g(x) = \sin \left( \left[ ([f(x) + 7]^{16} + 11)^{1/12} + e^{2x} \right]^{99} \right).$$

Find  $g'(x)$ .