

Kazhdan-Lusztig polynomials of matroids

Nicholas Proudfoot

University of Oregon

AMS Special Session on Arrangements of Hypersurfaces

Arrangements and Flats

Let V be a finite dimensional vector space

Arrangements and Flats

Let V be a finite dimensional vector space

\mathcal{A} a finite set of hyperplanes in V with $\bigcap_{H \in \mathcal{A}} H = \{0\}$

Arrangements and Flats

Let V be a finite dimensional vector space

\mathcal{A} a finite set of hyperplanes in V with $\bigcap_{H \in \mathcal{A}} H = \{0\}$


$F \subset V$ a flat (intersection of some hyperplanes)

Arrangements and Flats

Let V be a finite dimensional vector space

\mathcal{A} a finite set of hyperplanes in V with $\bigcap_{H \in \mathcal{A}} H = \{0\}$

$F \subset V$ a flat (intersection of some hyperplanes)



Arrangements and Flats

Definition

The **contraction of \mathcal{A} at F** is the arrangement

$$\mathcal{A}^F := \{H \cap F \mid F \not\subset H \in \mathcal{A}\}$$

in the vector space F .


Arrangements and Flats

Definition


The **contraction of \mathcal{A} at F** is the arrangement

$$\mathcal{A}^F := \{H \cap F \mid F \not\subset H \in \mathcal{A}\}$$

in the vector space F .



\mathcal{A}



\mathcal{A}^F

Definition

The **localization of \mathcal{A} at F** is the arrangement

$$\mathcal{A}_F := \{H/F \mid F \subset H \in \mathcal{A}\}$$

in the vector space V/F .



Arrangements and Flats

Definition

The **localization of \mathcal{A} at F** is the arrangement

$$\mathcal{A}_F := \{H/F \mid F \subset H \in \mathcal{A}\}$$

in the vector space V/F .



Characteristic Polynomial

Let $\chi_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ be the **characteristic polynomial** of \mathcal{A} .

Characteristic Polynomial

Let $\chi_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ be the **characteristic polynomial** of \mathcal{A} .


If V is a vector space over \mathbb{F}_q , $\chi_{\mathcal{A}}(q) = |V \setminus \bigcup_{H \in \mathcal{A}} H|$.

Characteristic Polynomial

Let $\chi_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ be the **characteristic polynomial** of \mathcal{A} .

If V is a vector space over \mathbb{F}_q , $\chi_{\mathcal{A}}(q) = |V \setminus \bigcup_{H \in \mathcal{A}} H|$.

Example



$$\chi_{\mathcal{A}}(t) = t^2 - 3t + 2$$

Kazhdan-Lusztig Polynomial

Theorem

There exists a unique way to assign to each arrangement \mathcal{A} a polynomial $P_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ subject to the following conditions:

Theorem

There exists a unique way to assign to each arrangement \mathcal{A} a polynomial $P_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ subject to the following conditions:

- *If $\dim V = 0$, $P_{\mathcal{A}}(t) = 1$*

Theorem

There exists a unique way to assign to each arrangement \mathcal{A} a polynomial $P_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ subject to the following conditions:

- *If $\dim V = 0$, $P_{\mathcal{A}}(t) = 1$*
- *If $\dim V > 0$, $\deg P_{\mathcal{A}}(t) < \frac{1}{2} \dim V$*

Kazhdan-Lusztig Polynomial

Theorem

There exists a unique way to assign to each arrangement \mathcal{A} a polynomial $P_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ subject to the following conditions:

- *If $\dim V = 0$, $P_{\mathcal{A}}(t) = 1$*
- *If $\dim V > 0$, $\deg P_{\mathcal{A}}(t) < \frac{1}{2} \dim V$*
- *$t^{\dim V} P_{\mathcal{A}}(t^{-1}) = \sum_F \chi_{\mathcal{A}_F}(t) P_{\mathcal{A}_F}(t)$.*

Kazhdan-Lusztig Polynomial

Theorem

There exists a unique way to assign to each arrangement \mathcal{A} a polynomial $P_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ subject to the following conditions:

- *If $\dim V = 0$, $P_{\mathcal{A}}(t) = 1$*
- *If $\dim V > 0$, $\deg P_{\mathcal{A}}(t) < \frac{1}{2} \dim V$*
- *$t^{\dim V} P_{\mathcal{A}}(t^{-1}) = \sum_F \chi_{\mathcal{A}_F}(t) P_{\mathcal{A}_F}(t)$.*

$P_{\mathcal{A}}(t)$ is called the **Kazhdan-Lusztig polynomial** of \mathcal{A} .

Kazhdan-Lusztig Polynomial

Theorem

There exists a unique way to assign to each arrangement \mathcal{A} a polynomial $P_{\mathcal{A}}(t) \in \mathbb{Z}[t]$ subject to the following conditions:

- *If $\dim V = 0$, $P_{\mathcal{A}}(t) = 1$*
- *If $\dim V > 0$, $\deg P_{\mathcal{A}}(t) < \frac{1}{2} \dim V$*
- $t^{\dim V} P_{\mathcal{A}}(t^{-1}) = \sum_F \chi_{\mathcal{A}_F}(t) P_{\mathcal{A}^F}(t).$

$P_{\mathcal{A}}(t)$ is called the **Kazhdan-Lusztig polynomial** of \mathcal{A} .

Remark

The theory of Kazhdan-Lusztig-Stanley polynomials provides a common generalization of these polynomials and classical Kazhdan-Lusztig polynomials.

Geometric Interpretation

$$V \hookrightarrow \bigoplus_{H \in \mathcal{A}} V/H \cong \bigoplus_{H \in \mathcal{A}} \mathbb{A}^1 \subset \prod_{H \in \mathcal{A}} \mathbb{P}^1$$

Geometric Interpretation

$$V \hookrightarrow \bigoplus_{H \in \mathcal{A}} V/H \cong \bigoplus_{H \in \mathcal{A}} \mathbb{A}^1 \subset \prod_{H \in \mathcal{A}} \mathbb{P}^1$$

Definition

We define the **Schubert variety of \mathcal{A}**

$$Y_{\mathcal{A}} := \bar{V} \subset \prod_{H \in \mathcal{A}} \mathbb{P}^1.$$

Geometric Interpretation

$$V \hookrightarrow \bigoplus_{H \in \mathcal{A}} V/H \cong \bigoplus_{H \in \mathcal{A}} \mathbb{A}^1 \subset \prod_{H \in \mathcal{A}} \mathbb{P}^1$$

Definition

We define the **Schubert variety of \mathcal{A}**

$$Y_{\mathcal{A}} := \bar{V} \subset \prod_{H \in \mathcal{A}} \mathbb{P}^1.$$

Have $H^*(Y_{\mathcal{A}}) \cong IH^*(Y_{\mathcal{A}})$, both concentrated in even degree.

Theorem (Huh-Wang, P-Xu-Young, Elias-P-Wakefield)

- $$\sum t^i \dim H^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F}$$

Geometric Interpretation

Theorem (Huh-Wang, P-Xu-Young, Elias-P-Wakefield)

- $\sum t^i \dim H^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F}$
- $\sum t^i \dim IH^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F} P_{\mathcal{A}^F}(t) =: Z_{\mathcal{A}}(t)$

Geometric Interpretation

Theorem (Huh-Wang, P-Xu-Young, Elias-P-Wakefield)

- $\sum t^i \dim H^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F}$
- $\sum t^i \dim IH^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F} P_{\mathcal{A}^F}(t) =: Z_{\mathcal{A}}(t)$
- $\sum t^i \dim \left(IH^{2i}(Y_{\mathcal{A}}) / H^2(Y_{\mathcal{A}}) \cdot IH^{2i-2}(Y_{\mathcal{A}}) \right) = P_{\mathcal{A}}(t)$

Geometric Interpretation

Theorem (Huh-Wang, P-Xu-Young, Elias-P-Wakefield)

- $\sum t^i \dim H^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F}$
- $\sum t^i \dim IH^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F} P_{\mathcal{A}^F}(t) =: Z_{\mathcal{A}}(t)$
- $\sum t^i \dim \left(IH^{2i}(Y_{\mathcal{A}}) / H^2(Y_{\mathcal{A}}) \cdot IH^{2i-2}(Y_{\mathcal{A}}) \right) = P_{\mathcal{A}}(t)$

Corollary

The polynomial $P_{\mathcal{A}}(t)$ has non-negative coefficients.

Geometric Interpretation

Theorem (Huh-Wang, P-Xu-Young, Elias-P-Wakefield)

- $\sum t^i \dim H^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F}$
- $\sum t^i \dim IH^{2i}(Y_{\mathcal{A}}) = \sum_F t^{\text{codim } F} P_{\mathcal{A}^F}(t) =: Z_{\mathcal{A}}(t)$
- $\sum t^i \dim \left(IH^{2i}(Y_{\mathcal{A}}) / H^2(Y_{\mathcal{A}}) \cdot IH^{2i-2}(Y_{\mathcal{A}}) \right) = P_{\mathcal{A}}(t)$

Corollary

The polynomial $P_{\mathcal{A}}(t)$ has non-negative coefficients.

Remark

The definition of $P_{\mathcal{A}}(t)$ makes sense for matroids, but when the matroid is not realizable, non-negativity is still a conjecture.

Work in progress by Braden-Huh-Matherne-P-Wang.

Example

Let \mathcal{A}_n be an arrangement of n generic hyperplanes in \mathbb{C}^{n-1} .

Example

Let \mathcal{A}_n be an arrangement of n generic hyperplanes in \mathbb{C}^{n-1} .

What are the flats?

Example

Let \mathcal{A}_n be an arrangement of n generic hyperplanes in \mathbb{C}^{n-1} .

What are the flats?

- For each $k < n - 1$, there are $\binom{n}{k}$ flats of codimension k .

Example

Let \mathcal{A}_n be an arrangement of n generic hyperplanes in \mathbb{C}^{n-1} .

What are the flats?

- For each $k < n - 1$, there are $\binom{n}{k}$ flats of codimension k .
For such a flat F ,

$$\mathcal{A}_n^F \cong \mathcal{A}_{n-k}$$

Example

Let \mathcal{A}_n be an arrangement of n generic hyperplanes in \mathbb{C}^{n-1} .

What are the flats?

- For each $k < n - 1$, there are $\binom{n}{k}$ flats of codimension k .
For such a flat F ,

$$\mathcal{A}_n^F \cong \mathcal{A}_{n-k}$$

and $(\mathcal{A}_n)_F$ is Boolean of rank k .

Example

Let \mathcal{A}_n be an arrangement of n generic hyperplanes in \mathbb{C}^{n-1} .

What are the flats?

- For each $k < n - 1$, there are $\binom{n}{k}$ flats of codimension k .
For such a flat F ,

$$\mathcal{A}_n^F \cong \mathcal{A}_{n-k}$$

and $(\mathcal{A}_n)_F$ is Boolean of rank k .

- There is a unique flat of codimension $n - 1$.

Example

Let \mathcal{A}_n be an arrangement of n generic hyperplanes in \mathbb{C}^{n-1} .

What are the flats?

- For each $k < n - 1$, there are $\binom{n}{k}$ flats of codimension k .
For such a flat F ,

$$\mathcal{A}_n^F \cong \mathcal{A}_{n-k}$$

and $(\mathcal{A}_n)_F$ is Boolean of rank k .

- There is a unique flat of codimension $n - 1$.

$$\begin{aligned} t^{n-1} P_{\mathcal{A}_n}(t^{-1}) &= \sum_{k=0}^{n-2} \binom{n}{k} (t-1)^k P_{\mathcal{A}_{n-k}}(t) + \chi_{\mathcal{A}_n}(t) \\ &= \sum_{k=0}^{n-2} \binom{n}{k} (t-1)^k P_{\mathcal{A}_{n-k}}(t) + \frac{(t-1)^n + (-1)^n (t-1)}{t} \end{aligned}$$

Example

Put it in a generating function:

$$\Phi(t, u) := \sum_{n=2}^{\infty} P_{\mathcal{A}_n}(t) u^{n-1}$$

Example

Put it in a generating function:

$$\Phi(t, u) := \sum_{n=2}^{\infty} P_{\mathcal{A}_n}(t) u^{n-1}$$

Then our recursion becomes

$$\begin{aligned} \Phi(t^{-1}, tu) &= \sum_{n=2}^{\infty} t^{n-1} P_{\mathcal{A}_n}(t^{-1}) u^{n-1} \\ &= \sum_{n=2}^{\infty} \left(\sum_{k=0}^{n-2} \binom{n}{k} (t-1)^k P_{\mathcal{A}_{n-k}}(t) + \chi_{\mathcal{A}_n}(t) \right) u^{n-1} \\ &= \dots \\ &= \frac{(1+u) \Phi\left(t, \frac{u}{1-u(t-1)}\right) + u(t-1)(1-u(t-1))}{(1+u)(1-u(t-1))^2} \end{aligned}$$

Example

Theorem (P-Wakefield-Young)

- $$\Phi(t, u) = \frac{2}{u} \cdot \frac{(2t+1)u - 1 + \sqrt{1 - 2(2tu+1)u + u^2}}{1 - (2tu+1)^2}$$

Example

Theorem (P-Wakefield-Young)

- $\Phi(t, u) = \frac{2}{u} \cdot \frac{(2t+1)u - 1 + \sqrt{1 - 2(2tu+1)u + u^2}}{1 - (2tu+1)^2}$
- $P_{\mathcal{A}_n}(t) = \sum_{i \geq 0} \frac{1}{i+1} \binom{n}{i} \binom{n-i-2}{i} t^i$

Example

Theorem (P-Wakefield-Young)

- $\Phi(t, u) = \frac{2}{u} \cdot \frac{(2t+1)u - 1 + \sqrt{1 - 2(2tu+1)u + u^2}}{1 - (2tu+1)^2}$
- $P_{\mathcal{A}_n}(t) = \sum_{i \geq 0} \frac{1}{i+1} \binom{n}{i} \binom{n-i-2}{i} t^i$

Corollary (Gedeon-P-Young)

The polynomial $P_{\mathcal{A}_n}(t)$ is real-rooted for all n .

Example

Theorem (P-Wakefield-Young)

- $\Phi(t, u) = \frac{2}{u} \cdot \frac{(2t+1)u - 1 + \sqrt{1 - 2(2tu+1)u + u^2}}{1 - (2tu+1)^2}$
- $P_{\mathcal{A}_n}(t) = \sum_{i \geq 0} \frac{1}{i+1} \binom{n}{i} \binom{n-i-2}{i} t^i$

Corollary (Gedeon-P-Young)

The polynomial $P_{\mathcal{A}_n}(t)$ is real-rooted for all n .

Remark

The polynomial $P_{\mathcal{A}}(t)$ is conjecturally real-rooted for any \mathcal{A} , but this is the only non-trivial family of examples for which we can prove it!

Thanks!