

The geometry of zonotopal algebras II: Orlik–Terao algebras and Schubert varieties

Colin Crowley¹

Department of Mathematics, University of Oregon, Eugene, OR 97403
and School of Mathematics, Institute for Advanced Study, Princeton, NJ 08540

Nicholas Proudfoot²

Department of Mathematics, University of Oregon, Eugene, OR 97403

Abstract. Zonotopal algebras, introduced by Postnikov–Shapiro–Shapiro, Ardila–Postnikov, and Holtz–Ron, show up in many different contexts, including approximation theory, representation theory, Donaldson–Thomas theory, and hypertoric geometry. In the first half of this paper, we construct a perfect pairing between the internal zonotopal algebra of a linear space and the reduced Orlik–Terao algebra of the Gale dual linear space. As an application, we prove a conjecture of Moseley–Proudfoot–Young that relates the reduced Orlik–Terao algebra of a graph to the cohomology of a certain configuration space. In the second half of the paper, we interpret the Macaulay inverse system of a zonotopal algebra as the space of sections of a sheaf on the Schubert variety of a linear space. As an application of this, we use an equivariant resolution of the structure sheaf of the Schubert variety inside of a product of projective lines to produce an exact sequence relating internal and external zonotopal algebras.

1 Introduction

Zonotopal algebras come in three flavors: central, internal, and external. Central zonotopal algebras and their duals play an important role in approximation theory, dating back to the work of de Boor and Höllig [dBH83]; they have also arisen in the contexts of hypertoric geometry [HS02] and Donaldson–Thomas theory [RRT23]. Internal zonotopal algebras also have applications to approximation theory [Len14], and to graphical configuration spaces [CDBHP]. External zonotopal algebras were introduced by Postnikov, Shapiro, and Shapiro in the context of differential forms on flag manifolds [PSS99]. These three types of algebras were combined and regarded as parts of a single theory by Ardila and Postnikov [AP10], and independently by Holtz and Ron [HR11], who gave them their names.

Another important algebra associated with the same data is the Orlik–Terao algebra [OT94]. This algebra has connections to linear programming [SSV13], linear codes [Toh], Kazhdan–Lusztig theory of matroids [EPW16], and the theory of resonance varieties [Sch11]; it also plays a role in the categorification of the Tutte polynomial in [Ber10]. Our first main result is that the internal zonotopal algebra of a linear space is canonically isomorphic to the dual of the reduced Orlik–Terao algebra of the Gale dual linear space (Theorem 1.8). We use this to prove a conjecture of Moseley,

¹Supported by NSF grant DMS-2039316.

²Supported by NSF grants DMS-1954050, DMS-2053243, and DMS-2344861.

Proudfoot, and Young [MPY17] relating the Orlik–Terao algebra of a graphical arrangement with the cohomology ring of a certain configuration space (Corollary 1.11), which has previously been proved only for the complete graph [Pag23].

The Schubert variety of a linear space was introduced by Ardila and Boocher [AB16], and was used to prove non-negativity of the Kazhdan–Lusztig polynomial and Z -polynomial of a realizable matroid [EPW16, PXY18] and the Dowling–Wilson conjecture for realizable matroids [HW17]. When the linear space arises from the Coxeter arrangement of a Lie algebra as in Remark 1.1, the fundamental group of the real locus of the Schubert variety coincides with the pure virtual Weyl group of the Lie algebra [IKL⁺, Jia24]. We interpret the Macaulay inverse systems of all of the zotopal algebras as groups of sections of certain sheaves on the Schubert variety (Theorem 1.14), which naturally appear in the minimal free resolution of the structure sheaf of the Schubert variety inside of a product of projective lines (Theorem 1.15). This allows us to construct an exact sequence relating internal and external zotopal algebras (Corollary 1.18), equivariant with respect to any automorphisms of the arrangement, categorifying a numerical formula for the number of spanning sets of a matroid (Remark 1.19).

1.1 Background

The basic data structure with which we will work is that of a triple $\mathcal{A} = (E, V, L)$, where E is a finite set, $V = \bigoplus_{e \in E} V_e$ is a vector space over a field \mathbb{F} of characteristic zero equipped with a decomposition into lines indexed by E , and $L \subset V$ is a linear subspace. We will refer to such a triple as a **linear space**. We define the **Gale dual** $\mathcal{A}^! := (E, V^*, L^\perp)$. The linear space \mathcal{A} has an associated matroid $M_{\mathcal{A}}$ of rank equal to the dimension of L , and $M_{\mathcal{A}^!}$ is the dual of $M_{\mathcal{A}}$. An automorphism of \mathcal{A} is by definition a linear automorphism of V that preserves the decomposition into lines (possibly permuting them) and takes L to itself. Note that the automorphism group $\text{Aut}(\mathcal{A})$ contains a distinguished copy of the multiplicative group \mathbb{G}_m , acting by scalar multiplication. There is a canonical isomorphism $\text{Aut}(\mathcal{A}) \cong \text{Aut}(\mathcal{A}^!)$ that restricts to inversion on \mathbb{G}_m .

Remark 1.1. Suppose that we are given a vector space L and a multiset $\{H_e \mid e \in E\}$ of hyperplanes indexed by E that intersect only in the origin. Then we may define a linear space \mathcal{A} by putting $V_e := L/H_e$. An automorphism of \mathcal{A} consists of a pair (σ, τ) , where σ is a linear automorphism of L and τ is a permutation of E such that $\sigma(H_e) = H_{\tau(e)}$ for all $e \in E$. An arbitrary linear space \mathcal{A} arises via this construction if and only if the projection $L \rightarrow V_e$ is surjective for all e , or equivalently if the matroid $M_{\mathcal{A}}$ is loopless. The main reason that we choose to work with linear spaces rather than hyperplane arrangements is that looplessness is not preserved by duality.

For any element $\alpha \in L$, let $m(\alpha)$ be the number of coordinates $e \in E$ such that the projection $\alpha_e \in V_e$ is nonzero, and let ρ be the minimum value of $m(\alpha)$ for $0 \neq \alpha \in L$. For any integer $k \geq -(\rho + 1)$, Ardila and Postnikov [AP10, AP15] define the ring

$$R_{\mathcal{A},k} := \text{Sym } L / \left\langle \alpha^{m(\alpha)+k+1} \mid 0 \neq \alpha \in L \right\rangle.$$

Examples of these rings include the three **zonotopal algebras** introduced and studied by Holtz and Ron [HR11]:

$$\begin{aligned} & \text{the internal zonotopal algebra } \mathcal{R}_-(\mathcal{A}) = R_{\mathcal{A},-2}, \\ & \text{the central zonotopal algebra } \mathcal{R}(\mathcal{A}) = R_{\mathcal{A},-1}, \\ & \text{and the external zonotopal algebra } \mathcal{R}_+(\mathcal{A}) = R_{\mathcal{A},0}. \end{aligned}$$

Remark 1.2. The external zonotopal algebra $\mathcal{R}_+(\mathcal{A})$ was originally defined by Postnikov, Shapiro, and Shapiro [PSS99]. Their primary motivation came from the case where \mathcal{A} arises from the Coxeter arrangement of a Lie algebra via Remark 1.1, in which case they showed that the external zonotopal algebra is isomorphic to the subalgebra of differential forms on the flag manifold generated by curvature forms of Hermitian line bundles.

Remark 1.3. The word “zonotopal” derives from the special case where \mathcal{A} is **unimodular**: this means that L^* is equipped with a lattice Λ , the projection $L \rightarrow V_e \cong \mathbb{F}$ is given by a primitive element $\chi_e \in \Lambda$, and any subset of the vectors $\{\chi_e \mid e \in E\}$ spans a saturated sublattice of Λ . Consider the **zonotope**

$$Z(\mathcal{A}) := \left\{ \sum c_H \chi_H \mid 0 \leq c_H \leq 1 \text{ for all } H \in \mathcal{A} \right\} \subset \Lambda \otimes \mathbb{R}.$$

We denote its interior by $\mathring{Z}(\mathcal{A})$, and we also consider a translation $Z(\mathcal{A}, \lambda) := Z(\mathcal{A}) + \lambda$ by some generic³ element $\lambda \in \Lambda \otimes \mathbb{R}$. Intersecting with Λ , we obtain finite subsets

$$\begin{aligned} \mathcal{Z}_+(\mathcal{A}) &:= Z(\mathcal{A}) \cap \Lambda, \\ \mathcal{Z}(\mathcal{A}, \lambda) &:= Z(\mathcal{A}, \lambda) \cap \Lambda, \\ \mathcal{Z}_-(\mathcal{A}) &:= \mathring{Z}(\mathcal{A}) \cap \Lambda. \end{aligned}$$

Given a finite subset $S \subset L^*$, its **orbit harmonics ring** $\mathcal{R}(S)$ is defined to be the coordinate ring of the scheme-theoretic limit

$$\lim_{\epsilon \rightarrow 0} (\epsilon \cdot S);$$

this may also be characterized as the associated graded ring of the filtration on $\mathbb{F}[S]$ whose i^{th} piece is the image of $\text{Sym}^{\leq i} L$ under the restriction of functions from L^* to S . Then we have canonical isomorphisms [HR11, Theorems 5.10, 3.9, 4.11]⁴

$$\mathcal{R}_-(\mathcal{A}) \cong \mathcal{R}(\mathcal{Z}_-(\mathcal{A})), \quad \mathcal{R}(\mathcal{A}) \cong \mathcal{R}(\mathcal{Z}(\mathcal{A}, \lambda)), \quad \text{and} \quad \mathcal{R}_+(\mathcal{A}) \cong \mathcal{R}(\mathcal{Z}_+(\mathcal{A})).$$

Let $C_{\mathcal{A},k} \subset \text{Sym } L^*$ be the linear dual of $R_{\mathcal{A},k}$, also known as the **Macaulay inverse system** of the defining ideal of $R_{\mathcal{A},k}$. Concretely, we may think of elements of $\text{Sym } L^*$ as polynomials and elements of $\text{Sym } L$ as differential operators, in which the natural pairing between $\text{Sym } L^*$ and $\text{Sym } L$

³Here “generic” means that $Z(\mathcal{A}, \lambda)$ contains no lattice points on its boundary.

⁴See [CDBHP, Proposition 1.1] for an alternate proof in the internal case, which extends to the other two cases.

is given by applying the differential operator to the polynomial and then evaluating the resulting polynomial at $0 \in L$. With this interpretation, we may write

$$C_{\mathcal{A},k} = \left\{ f \in \text{Sym } L^* \mid \alpha^{m(\alpha)+k+1} \cdot f = 0 \text{ for all } 0 \neq \alpha \in L \right\}.$$

Holtz and Ron use the notation

$$\mathcal{P}_-(\mathcal{A}) := C_{\mathcal{A},-2}, \quad \mathcal{P}(\mathcal{A}) := C_{\mathcal{A},-1} \quad \text{and} \quad \mathcal{P}_+(\mathcal{A}) := C_{\mathcal{A},0}.$$

Remark 1.4. We will use the convention that the gradings on all of our vector spaces are induced by the inverse action of $\mathbb{G}_m \subset \text{Aut}(\mathcal{A})$. Thus $\text{Sym } L^*$ is generated in degree 1, $\text{Sym } L$ is generated in degree -1 , and all nontrivial pairings occur between classes in opposite degrees. This is consistent with thinking of elements of $\text{Sym } L^*$ as polynomials and elements of $\text{Sym } L$ as differential operators.

The in the central ($k = -1$) case, we have some beautiful additional structure, namely a canonical ring structure on $\mathcal{P}(\mathcal{A})$, which we now describe. For each $e \in E$, choose an isomorphism $V_e \cong \mathbb{F}$, and let $\chi_e \in L^*$ denote the projection from L to V_e . For any subset $S \subset E$, let

$$\chi_S := \prod_{e \in S} \chi_e \in \text{Sym } L^*.$$

Consider the ideal⁵ $\mathcal{J}(\mathcal{A}^1) \subset \text{Sym } L^*$ generated by polynomials χ_S for all subsets $B \subset S$ that are dependent for the matroid $M_{\mathcal{A}^1}$. The quotient ring $\text{Sym } L^*/\mathcal{J}(\mathcal{A}^1)$ is the reduced Stanley–Reisner algebra $\overline{\text{SR}}(\mathcal{A}^1)$ of the independence complex of $M_{\mathcal{A}^1}$.⁶ (See Remark 1.7 for an interpretation of this ring in the context of hypertoric geometry.)

Let $\mathcal{D}(\mathcal{A}^1) \subset \text{Sym } L$ denote the Macaulay inverse system of $\mathcal{J}(\mathcal{A}^1)$, or equivalently the linear dual of $\overline{\text{SR}}(\mathcal{A}^1)$. When $\mathbb{F} = \mathbb{R}$, $\mathcal{D}(\mathcal{A}^1)$ is equal to the **Dahmen–Micchelli space**, which is the $\text{Sym } L^*$ -submodule of $\text{Sym } L$ spanned by polynomials that coincide on some open subset of L^* with the multivariate spline associated with \mathcal{A} [DCP11, Theorem 11.37]. The Dahmen–Micchelli space is of fundamental importance in approximation theory; it was first studied by de Boor and Höllig [dBH83], and it is one of the main characters in the book [DCP11]. See [HR11, Section 1.2] for a more detailed survey of the literature on the space $\mathcal{D}(\mathcal{A}^1)$.

The following theorem was proved by several authors [AS88, DR90, Jia93], and appears implicitly in [BV99] and the references therein. See [DCP11, Corollary 11.23] or [HR11, Theorem 3.8] for a nice exposition.

Theorem 1.5. *We have $\text{Sym } L^* = \mathcal{P}(\mathcal{A}) \oplus \mathcal{J}(\mathcal{A}^1)$. Equivalently, the compositions*

$$\mathcal{P}(\mathcal{A}) \hookrightarrow \text{Sym } L^* \twoheadrightarrow \overline{\text{SR}}(\mathcal{A}^1) \quad \text{and} \quad \mathcal{D}(\mathcal{A}^1) \hookrightarrow \text{Sym } L \twoheadrightarrow \mathcal{R}(\mathcal{A})$$

⁵Holtz and Ron call this ideal $\mathcal{J}(\mathcal{A})$, but this notation would lead to difficulty for us in Sections 2 and 3. Similarly, the vector space $\mathcal{D}(\mathcal{A}^1)$ in the next paragraph is called $\mathcal{D}(\mathcal{A})$ by Holtz and Ron.

⁶The Stanley–Reisner algebra is a quotient of $\text{Sym } V^*$, and the reduced Stanley–Reisner algebra is the quotient of the Stanley–Reisner algebra by the linear system of parameters $L^\perp \subset V^*$, which is a quotient of $\text{Sym } L^*$.

are isomorphisms, and the graded vector spaces $\mathcal{P}(\mathcal{A}) \cong \overline{\text{SR}}(\mathcal{A}^\dagger)$ are dual to $\mathcal{D}(\mathcal{A}^\dagger) \cong \mathcal{R}(\mathcal{A})$.

It is natural to ask whether there is an analogue of Theorem 1.5 in the internal or external settings. Indeed, Holtz and Ron define homogeneous ideals

$$\mathcal{J}_-(\mathcal{A}^\dagger), \mathcal{J}_+(\mathcal{A}^\dagger) \subset \text{Sym } L^*$$

and prove that we have [HR11, Theorems 4.10 and 5.9]

$$\mathcal{P}_-(\mathcal{A}) \oplus \mathcal{J}_-(\mathcal{A}^\dagger) = \text{Sym } L^* = \mathcal{P}_+(\mathcal{A}) \oplus \mathcal{J}_+(\mathcal{A}^\dagger).$$

The drawback to these results is that the ideals $\mathcal{J}_\pm(\mathcal{A}^\dagger)$ depend on choices. For example, $\mathcal{J}_-(\mathcal{A}^\dagger)$ depends on a choice of ordering of E , and the quotient $\text{Sym } L^*/\mathcal{J}_-(\mathcal{A}^\dagger)$ is the reduced Stanley–Reisner algebra of the broken circuit complex of $M_{\mathcal{A}^\dagger}$. (The broken circuit complex, unlike the independence complex, depends on an ordering.) The ideal $\mathcal{J}_+(\mathcal{A}^\dagger)$ depends on both an ordering and a choice of basis for L^* that is suitably generic with respect to \mathcal{A}^\dagger . In particular, the group $\text{Aut}(\mathcal{A}) = \text{Aut}(\mathcal{A}^\dagger)$ acts on the rings $\mathcal{R}_\pm(\mathcal{A})$ and on their linear duals $\mathcal{P}_\pm(\mathcal{A})$, but not on $\mathcal{J}_\pm(\mathcal{A}^\dagger)$.

Remark 1.6. Like the central zonotopal algebra, the internal zonotopal algebra is also interesting from the perspective of approximation theory. When \mathcal{A} is unimodular (Remark 1.3), Holtz and Ron used the box spline associated with \mathcal{A} to define a canonical map from $\mathcal{P}_-(\mathcal{A})$ to the space of functions on $\mathcal{Z}_-(\mathcal{A})$, and Lenz proved that this map is an isomorphism [Len14].⁷

1.2 The Orlik–Terao algebra

The first half of this paper is devoted to providing a canonical counterpart to Theorem 1.5 in the internal setting. As in the previous section, choose isomorphisms $V_e \cong \mathbb{F}$ and let $\chi_e \in L^*$ denote the projection from L to V_e . The **Orlik–Terao algebra** of \mathcal{A} , defined in [OT94] and first systematically studied in [Ter02], is the subalgebra of rational functions on L generated by $\{\chi_e^{-1} \mid e \in E\}$.⁸ Consider the map from L to $\text{OT}(\mathcal{A})$ taking $\alpha \in L$ to $\sum \chi_e(\alpha)\chi_e^{-1}$. Note that this map does not change if we scale one of the linear functionals χ_e , and is therefore completely canonical. Equivalently, we may regard $\text{OT}(\mathcal{A})$ as a quotient of $\text{Sym } V$, and the map from L is given by the inclusion of L into $V_{\mathcal{A}}$. The image of this map is a linear system of parameters [PS06, Proposition 7], meaning that $\text{OT}(\mathcal{A})$ is a free module over $\text{Sym } L$ of finite rank.⁹ The quotient of $\text{OT}(\mathcal{A})$ by the ideal generated by the image of L is called the **reduced Orlik–Terao algebra**, and denoted $\overline{\text{OT}}(\mathcal{A})$.

Remark 1.7. If $\mathbb{F} = \mathbb{Q}$ and $M_{\mathcal{A}}$ is loopless, the Orlik–Terao algebra $\text{OT}(\mathcal{A})$ is canonically isomorphic to the equivariant intersection cohomology of the hypertoric variety associated with \mathcal{A}

⁷This is very different from the isomorphism obtained from [HR11, Result 2.3 and Theorem 5.10], which depends on a choice of positive definite inner product on Λ . In particular, that isomorphism takes $1 \in \mathcal{P}_-(\mathcal{A})$ to the constant function, whereas the more interesting and canonical isomorphism in Lenz’s theorem takes $1 \in \mathcal{P}_-(\mathcal{A})$ to the restriction of the box spline to $\mathcal{Z}_-(\mathcal{A})$.

⁸If $\chi_e = 0$ for some e , then the Orlik–Terao algebra is by definition identically zero.

⁹This statement can fail in positive characteristic, which is why it is crucial that we assume that $\text{char } \mathbb{F} = 0$.

[BP09], and the Stanley–Reisner algebra of the independence complex is canonically isomorphic to the cohomology ring of an orbifold symplectic resolution of that hypertoric variety [HS02]. Passing to the reduced Orlik–Terao algebra or the reduced Stanley–Reisner algebra corresponds to passing from equivariant to ordinary cohomology.

We will apply this construction to the Gale dual linear space \mathcal{A}^\dagger . The algebra $\text{OT}(\mathcal{A}^\dagger)$ is a quotient of $\text{Sym } V^*$, and $\overline{\text{OT}}(\mathcal{A}^\dagger)$ is a quotient of $\text{Sym}(V^*/L^\perp) \cong \text{Sym } L^*$. Let $\mathcal{K}(\mathcal{A}^\dagger) \subset \text{Sym } L^*$ denote the kernel of the homomorphism to $\overline{\text{OT}}(\mathcal{A}^\dagger)$. Let $\mathcal{E}(\mathcal{A}^\dagger) \subset \text{Sym } L$ denote the Macaulay inverse system of $\mathcal{K}(\mathcal{A}^\dagger)$, consisting of differential operators in $\text{Sym } L$ that pair trivially with the polynomials in $\mathcal{K}(\mathcal{A}^\dagger)$. The following analogue of Theorem 1.5 is our first main result.

Theorem 1.8. *We have $\text{Sym } L^* = \mathcal{P}_-(\mathcal{A}) \oplus \mathcal{K}(\mathcal{A}^\dagger)$. Equivalently, the compositions*

$$\mathcal{P}_-(\mathcal{A}) \hookrightarrow \text{Sym } L^* \twoheadrightarrow \overline{\text{OT}}(\mathcal{A}^\dagger) \quad \text{and} \quad \mathcal{E}(\mathcal{A}^\dagger) \hookrightarrow \text{Sym } L \twoheadrightarrow \mathcal{R}_-(\mathcal{A})$$

are isomorphisms, and the graded vector spaces $\mathcal{P}_-(\mathcal{A}) \cong \overline{\text{OT}}(\mathcal{A}^\dagger)$ are dual to $\mathcal{E}(\mathcal{A}^\dagger) \cong \mathcal{R}_-(\mathcal{A})$.

Remark 1.9. While the statement of Theorem 1.8 is elementary, our proof requires the theory of minimal extension sheaves on the poset of flats of the arrangement \mathcal{A}^\dagger , developed in [BP09].

1.3 An application to graphs

Theorem 1.8 has a concrete application outside of the world of zonotopal algebras, which we now describe. Let Γ be a connected graph with vertex set $[n]$. Let

$$\text{Conf}(SU(2), \Gamma) := \{\kappa : [n] \rightarrow SU(2) \mid \kappa(i) \neq \kappa(j) \text{ if } i \text{ is adjacent to } j \text{ in } \Gamma\},$$

and let $X(SU(2), \Gamma)$ be the quotient of $\text{Conf}(SU(2), \Gamma)$ by the action of $SU(2)$ by simultaneous left translation.

Remark 1.10. This space can be identified with something more familiar in the case of the complete graph K_n . Since $SU(2) \setminus \{\text{id}\}$ is homeomorphic to \mathbb{R}^3 , we may use the translation action of $SU(2)$ to move $\kappa(n)$ to the identity and thus identify $X(SU(2), K_n)$ with the configuration space of $n-1$ distinct labeled points in \mathbb{R}^3 . The advantage of writing it as $X(SU(2), K_n)$ is that it reveals the fact that we have an action of \mathfrak{S}_n and not just \mathfrak{S}_{n-1} . The \mathfrak{S}_{n-1} action on the cohomology groups of $X(SU(2), K_n)$ gives (up to a sign) the Eulerian representations \mathfrak{S}_{n-1} , and the \mathfrak{S}_n action gives (up to a sign) Whitehouse’s lifts of the Eulerian representations to \mathfrak{S}_n [ER19, Proposition 2]. For arbitrary Γ , the space $X(SU(2), \Gamma)$ is more mysterious; it may be regarded as a finite approximation to a product of g copies of the $SU(2)$ affine Grassmanian, where g is the first Betti number of Γ [CDBHP, Remark 1.12].

Let $\mathcal{A}_\Gamma^\dagger := (E, C_1(\Gamma; \mathbb{Q}), H_1(\Gamma; \mathbb{Q}))$, where E is the set of edges and $C_1(\Gamma; \mathbb{Q})$ is the space of cellular 1-chains. Let \mathcal{A}_Γ be the Gale dual of $\mathcal{A}_\Gamma^\dagger$. If Γ is simple, then \mathcal{A}_Γ is the linear space associated with the corresponding graphical hyperplane arrangement, that is, the subarrangement

of the Coxeter arrangement for \mathfrak{sl}_n indexed by edges of Γ . The following statement was conjectured in [MPY17, Conjecture 2.16].

Corollary 1.11. *There exists a degree-halving isomorphism of graded $\text{Aut}(\Gamma)$ -representations*

$$H^*(X(SU(2), \Gamma); \mathbb{Q}) \cong \overline{\text{OT}}(\mathcal{A}_\Gamma).$$

Proof. By [CDBHP, Theorem 1.9], there exists an $\text{Aut}(\mathcal{A}_\Gamma)$ -equivariant vector space isomorphism $H^*(X(SU(2), \Gamma); \mathbb{Q}) \cong \mathcal{R}_-(\mathcal{A}_\Gamma^!)$ taking classes of degree $2i$ to classes of degree $-i$. Combining this with Theorem 1.8, we see that $H^{2i}(X(SU(2), \Gamma); \mathbb{Q})$ is dual to $\overline{\text{OT}}^i(\mathcal{A}_\Gamma)$ as a representation of $\text{Aut}(\Gamma) \subset \text{Aut}(\mathcal{A}_\Gamma)$. The corollary then follows from the fact that finite dimensional representations of finite groups over \mathbb{Q} are (noncanonically) self-dual. \square

Remark 1.12. The case of the complete graph is already very interesting, and was the primary motivation for the conjecture in [MPY17]. For this special case, Corollary 1.11 was proved by Pagaria [Pag23] by studying the generating functions for the symmetric functions obtained by applying the Frobenius characteristic map to the graded pieces of the representations. This is a beautiful proof, but it does not provide any canonical isomorphism or duality, nor does it extend to other graphs.

Remark 1.13. There are many examples of linear spaces over \mathbb{C} for which $\mathcal{R}_-(\mathcal{A})$ and $\overline{\text{OT}}(\mathcal{A}^!)$ are dual but not isomorphic as graded representations of a finite group of symmetries. For example, see [Ber18] for an equivariant study of $\mathcal{R}_-(\mathcal{A})$ when $\mathcal{A}^!$ is the linear space associated with the reflection hyperplanes for the complex reflection group $G(m, 1, n)$. Theorem 1.8 allows us to translate this into a study of the reduced Orlik–Terao algebra of $\mathcal{A}^!$.

1.4 The Schubert variety

Assume now that the ground field \mathbb{F} is algebraically closed. For all $e \in E$, let

$$\overline{V}_e := V_e \sqcup \{\infty\} = \mathbb{P}(V_e \oplus \mathbb{F}) \cong \mathbb{P}^1.$$

The **Schubert variety** $Y_{\mathcal{A}}$ is defined as the closure of L inside of $\mathbb{P}_{\mathcal{A}} := \prod \overline{V}_e$. The name Schubert variety indicates an analogy with Schubert varieties in Lie theory; the most important shared feature is that both varieties admit affine pavings. The translation action of L on itself extends to an action on $Y_{\mathcal{A}}$, which has finitely many orbits indexed by flats of $M_{\mathcal{A}}$, each of which is isomorphic to an affine space (Lemma 4.2).

Let F be a corank 1 flat. The corresponding orbit in $Y_{\mathcal{A}}$ has codimension 1, and we denote its closure by D_F . Let $\alpha_F \in L$ span the line in L corresponding to F , which is the stabilizer of the orbit, and let $m(F) := m(\alpha_F)$. For any integer k , consider the Weyl divisor

$$D_k := \sum_F (m(F) + k) D_F$$

along with the associated sheaf $\mathcal{O}_{Y_{\mathcal{A}}}(D_k)$.¹⁰ Concretely, a section of $\mathcal{O}_{Y_{\mathcal{A}}}(D_k)$ is a rational function on $Y_{\mathcal{A}}$ that is regular on L and has a pole of order at most $m(F) + k$ on D_F . In particular, we have a canonical inclusion $H^0(\mathcal{O}_{Y_{\mathcal{A}}}(D_k)) \subset H^0(\mathcal{O}_L) \cong \text{Sym } L^*$. Our second main result identifies this space of sections with the Macaulay inverse system $C_{\mathcal{A},k}$. Recall that $R_{\mathcal{A},k}$ and $C_{\mathcal{A},k}$ are only defined when $k \geq -(\rho + 1)$, where ρ is the smallest possible value of $m(\alpha)$.

Theorem 1.14. *For all $-(\rho+1) \leq k \leq 0$, we have $H^0(\mathcal{O}_{Y_{\mathcal{A}}}(D_k)) = C_{\mathcal{A},k}$ as subspaces of $\text{Sym } L^*$.*¹¹

1.5 Syzygies

Our primary motivation for Theorem 1.14 is that these cohomology groups appear naturally in the minimal resolution of the structure sheaf of $Y_{\mathcal{A}}$ on the ambient variety $\mathbb{P}_{\mathcal{A}}$. For any $S \subset E$, let $V_S := \bigoplus_{e \in S} V_e$, let $\pi_S : V \rightarrow V_S$ denote the projection, and let $L_S := \pi_S(L) \subset V_S$. We define the **localization** $\mathcal{A}_S := (S, V_S, L_S)$ and the **contraction** $\mathcal{A}^S := (E \setminus S, V_{E \setminus S}, L \cap V_{E \setminus S})$, so that $(\mathcal{A}^!)_S = (\mathcal{A}^{E \setminus S})^!$. At the level of matroids, these operations correspond to restriction to S and contraction of S , respectively. Let $\pi_e : \mathbb{P}_{\mathcal{A}} \rightarrow \overline{V}_e$ denote the coordinate projection, and let

$$\mathcal{O}_{\mathbb{P}_{\mathcal{A}}}(-1_S) := \bigotimes_{e \in S} \pi_e^* \mathcal{O}_{\overline{V}_e}(-1).$$

Let $M = M_{\mathcal{A}^!}$ be the matroid associated with the Gale dual of \mathcal{A} . For any vector space V , let $\det(V) := \wedge^{\dim V} V$.

Theorem 1.15. *There exists an $\text{Aut}(\mathcal{A})$ -equivariant minimal free resolution*

$$\cdots \rightarrow \mathcal{Q}_2 \rightarrow \mathcal{Q}_1 \rightarrow \mathcal{Q}_0 \rightarrow \mathcal{O}_{Y_{\mathcal{A}}} \rightarrow 0$$

of the structure sheaf $\mathcal{O}_{Y_{\mathcal{A}}}$ on $\mathbb{P}_{\mathcal{A}}$, and we have

$$\mathcal{Q}_i \cong \bigoplus_{\text{crk}_M(E \setminus S) = i} \det(L_S^\perp) \otimes H^0(\mathcal{O}_{Y_{\mathcal{A}_S}}(D_{-2}))^* \otimes \mathcal{O}_{\mathbb{P}_{\mathcal{A}}}(-1_S).^{12}$$

Remark 1.16. By Theorem 1.14, the factor $H^0(\mathcal{O}_{Y_{\mathcal{A}_S}}(D_{-2}))^*$ in Theorem 1.15 may be replaced by the internal zonotopal algebra $\mathcal{R}_-(\mathcal{A}_S)$. By Theorem 1.8, it may be replaced by $\overline{\text{OT}}((\mathcal{A}_S)^!)$.

Remark 1.17. Theorem 1.15 is a categorification of [AB16, Theorem 1.5], in which the multigraded Betti numbers of $\mathcal{O}_{Y_{\mathcal{A}}}$ are computed. The novelty of Theorem 1.15 is that it encodes the action of the group $\text{Aut}(\mathcal{A})$ on the multiplicity spaces, including the multiplicative group $\mathbb{G}_m \subset \text{Aut}(\mathcal{A})$, which is responsible for the gradings.

Though we have written the resolution in Theorem 1.15 as a direct sum over all subsets $S \subset E$, many of the terms vanish. Indeed, $\mathcal{R}_-(\mathcal{A}_S)$ is nonzero if and only if $M_{\mathcal{A}_S}$ has no coloops, which is the case if and only if $E \setminus S$ is a flat of M .

¹⁰We are using the fact that $Y_{\mathcal{A}}$ is normal (Theorem 4.1) to make sense of $\mathcal{O}_{Y_{\mathcal{A}}}(D_k)$.

¹¹When k is positive, $H^0(\mathcal{O}_{Y_{\mathcal{A}}}(D_k))$ is equal to the slightly larger space $C'_{\mathcal{A},k}$ appearing in [AP15].

¹²The action of $\text{Aut}(\mathcal{A})$ on the right-hand side of this expression will be made precise in Remark 4.7.

We conclude by considering what happens when we tensor the exact sequence in Theorem 1.15 with $\mathcal{O}(1_E) := \mathcal{O}(-1_E)^*$ and take cohomology. All of the sheaves appearing in this new exact sequence have vanishing higher cohomology and $\mathcal{O}_{Y_{\mathcal{A}}} \otimes \mathcal{O}(1_E) \cong \mathcal{O}_{Y_{\mathcal{A}}}(D_0)$, which leads to the following result relating internal and external zonotopal algebras.

Corollary 1.18. *There exists an $\text{Aut}(\mathcal{A})$ -equivariant exact sequence of vector spaces*

$$\cdots \rightarrow H^0(\mathcal{Q}_2(1_E)) \rightarrow H^0(\mathcal{Q}_1(1_E)) \rightarrow H^0(\mathcal{Q}_0(1_E)) \rightarrow \mathcal{R}_+(\mathcal{A})^* \rightarrow 0,$$

where

$$H^0(\mathcal{Q}_i(1_E)) \cong \bigoplus_{\text{crk}_M(E \setminus S)=i} \det(L_S^\perp) \otimes \mathcal{R}_-(\mathcal{A}_S) \otimes \bigotimes_{e \in E \setminus S} (V_e^* \oplus \mathbb{F}).$$

Remark 1.19. We have noted in Remark 1.17 that $\mathcal{R}_-(\mathcal{A}_S)$ vanishes unless $E \setminus S$ is a flat of M . If it is a flat, then the total dimension of $\mathcal{R}_-(\mathcal{A}_S)$ is equal to $(-1)^{\text{crk}(E \setminus S)} \mu(E \setminus S, E)$, where μ is the Möbius function on the lattice of flats of M [HR11, Theorem 5.9(1)]. The total dimension of $\mathcal{R}_+(\mathcal{A})$ is equal to the number of spanning sets of M [HR11, Theorem 4.10(1)]. Thus the exact sequence in Corollary 1.18 categorifies the equation

$$\text{number of spanning sets of } M = \sum_{\text{flats } F} \mu(F, E) 2^{|F|}.$$

This identity is obtained via Möbius inversion from the more familiar identity

$$2^{|E|} = \sum_{\text{flats } F} |\{T \subset E \mid \bar{T} = F\}|.$$

Acknowledgments: We are grateful to Andy Berget and Matt Larson for valuable conversations. We would also like to thank the creators of Macaulay2, as well as Federico Galleto for writing the BettiCharacters package, which was helpful in proving Theorem 1.15. The first author thanks the Simons Foundation for support. The second author thanks All Souls College for its hospitality during the preparation of this manuscript. Both authors thank the National Science Foundation.

2 Modules over $\text{Sym } L^\perp$

The algebra $\overline{\text{OT}}(\mathcal{A}^\dagger)$ comes with a flat deformation $\text{OT}(\mathcal{A}^\dagger)$ over the base $(L^\perp)^* \cong V_{\mathcal{A}}/L$. Our strategy for proving Theorem 1.8 will be to define a corresponding flat deformation $\mathcal{Q}_-(\mathcal{A})$ of $\mathcal{P}_-(\mathcal{A})$ over the same base, and to prove that $\mathcal{Q}_-(\mathcal{A})$ is isomorphic to $\text{OT}(\mathcal{A}^\dagger)$.

2.1 OT and $\text{SR}_<$

Fix for each $e \in E$ a nonzero vector $v_e \in V_e$, and let $w_e \in V_e^*$ be the unique vector that pairs to 1 with v_e . Let $\chi_e \in L^*$ be the linear functional obtained by projecting onto V_e and pairing with w_e ,

and let $\psi_e \in (L^\perp)^*$ be the linear functional obtained by projecting onto V_e^* and pairing with v_e . We have surjections

$$\mathrm{Sym} V \rightarrow \mathrm{OT}(\mathcal{A}) \quad \text{and} \quad \mathrm{Sym} V^* \rightarrow \mathrm{OT}(\mathcal{A}^\dagger)$$

taking v_e to χ_e^{-1} and w_e to ψ_e^{-1} . We will only be interested in $\mathrm{OT}(\mathcal{A}^\dagger)$, but we provide the two constructions in parallel to make them more transparent.

Remark 2.1. Note that scaling v_e has the effect of scaling ψ_e and inverse scaling w_e and χ_e , so these surjections are in fact independent of any choices. This observation makes it clear why $\mathrm{OT}(\mathcal{A})$ is canonically a quotient of $\mathrm{Sym} V$ rather than $\mathrm{Sym} V^*$.

Let $\mathcal{G}(\mathcal{A}^\dagger) \subset \mathrm{Sym} V^*$ be the kernel of the homomorphism to $\mathrm{OT}(\mathcal{A}^\dagger)$. For any element $\alpha \in L$, let $\mathrm{Supp}(\alpha) := \{e \mid \chi_e(\alpha) \neq 0\}$, so that $m(\alpha) = |\mathrm{Supp}(\alpha)|$. Since α vanishes on L^\perp , so does the rational function

$$\alpha \prod_{e \in \mathrm{Supp}(\alpha)} \psi_e^{-1} = \sum_{e \in \mathrm{Supp}(\alpha)} \chi_e(\alpha) \prod_{f \in \mathrm{Supp}(\alpha) \setminus e} \psi_f^{-1}.$$

Thus we have a corresponding element

$$h_\alpha := \sum_{e \in \mathrm{Supp}(\alpha)} \chi_e(\alpha) \prod_{f \in \mathrm{Supp}(\alpha) \setminus e} w_f \in \mathcal{G}(\mathcal{A}^\dagger).$$

If there exists some $\alpha \in L$ with $\mathrm{Supp}(\alpha) = \{e\}$, then h_α is a nonzero constant, and therefore $\mathcal{G}(\mathcal{A}^\dagger) = \mathrm{Sym} V^*$. This happens if and only if $\psi_e = 0$, and therefore agrees with the convention in Footnote 8 that $\mathrm{OT}(\mathcal{A}^\dagger) = 0$ in such cases.

Fix a total ordering $<$ of E , along with a monomial order on $\mathrm{Sym} V^*$ with the property that $w_e < w_f$ if and only if $e < f$. For any $0 \neq g \in \mathrm{Sym} V^*$, we define the initial term $\mathrm{in}_<(g) \in \mathrm{Sym} V^*$ to be the smallest monomial appearing in g with respect to our monomial order. Let

$$\mathcal{G}_<(\mathcal{A}^\dagger) := \left\langle \mathrm{in}_<(h_\alpha) \mid \alpha \in L \right\rangle \subset \mathrm{Sym} V^*.$$

This ideal is called the Stanley–Reisner ideal of the broken circuit complex of $M_{\mathcal{A}^\dagger}$, and the quotient

$$\mathrm{SR}_<(\mathcal{A}^\dagger) := \mathrm{Sym} V^* / \mathcal{G}_<(\mathcal{A}^\dagger)$$

is called the Stanley–Reisner algebra of the broken circuit complex.

We call $0 \neq \alpha \in L$ **minimal** if its support does not strictly contain the support of another nonzero element of L , or equivalently if the support of α is a circuit of $M_{\mathcal{A}^\dagger}$. It is clear that $\mathcal{G}_<(\mathcal{A}^\dagger)$ is generated by $\mathrm{in}_<(h_\alpha)$ for minimal α . Since $h_\alpha \in \mathcal{G}(\mathcal{A}^\dagger)$, we *a priori* have an inclusion $\mathcal{G}_<(\mathcal{A}^\dagger) \subset \mathrm{in}(\mathcal{G}(\mathcal{A}^\dagger))$. In fact, this inclusion turns out to be an isomorphism; the following result is proved using the theory of geometric vertex decompositions [KMY09].

Theorem 2.2. [PS06, Theorem 4] *The set $\{h_\alpha \mid \alpha \text{ minimal}\}$ is a universal Gröbner basis for $\mathcal{G}(\mathcal{A}^\dagger)$. That is, for any choice of ordering $<$ and any compatible choice of monomial order, we have $\mathrm{in}(\mathcal{G}(\mathcal{A}^\dagger)) = \mathcal{G}_<(\mathcal{A}^\dagger)$.*

The inclusion of L^\perp into V^* makes both $\text{OT}(\mathcal{A}^\dagger)$ and $\text{SR}_{<}(\mathcal{A}^\dagger)$ graded modules over $\text{Sym } L^\perp$, and they are both free modules by [PS06, Propositions 1 and 7]. We define the **reduced Orlik–Terao algebra** and **reduced Stanley–Reisner algebra**

$$\overline{\text{OT}}(\mathcal{A}^\dagger) := \text{OT}(\mathcal{A}^\dagger) \otimes_{\text{Sym } L^\perp} \mathbb{F} \quad \text{and} \quad \overline{\text{SR}}_{<}(\mathcal{A}^\dagger) := \text{SR}_{<}(\mathcal{A}^\dagger) \otimes_{\text{Sym } \mathcal{A}^\dagger} \mathbb{F}.$$

We may equivalently regard $\overline{\text{OT}}(\mathcal{A}^\dagger)$ and $\overline{\text{SR}}_{<}(\mathcal{A}^\dagger)$ as quotients of $\text{Sym } L^* \cong \text{Sym}(V_{\mathcal{A}^\dagger}^*/L^\perp)$ by the images of $\mathcal{G}(\mathcal{A}^\dagger)$ and $\mathcal{G}_{<}(\mathcal{A}^\dagger)$ in $\text{Sym } L^*$. These images are precisely the ideals $\mathcal{K}(\mathcal{A}^\dagger)$ and $\mathcal{J}_{<}(\mathcal{A}^\dagger)$ from Sections 1.2 and 1.1.

2.2 \mathcal{Q}_-

Recall that we have defined

$$\mathcal{P}_-(\mathcal{A}) := \left\{ f \in \text{Sym } L^* \mid \alpha^{m(\alpha)-1} \cdot f = 0 \text{ for all } 0 \neq \alpha \in L \right\}.$$

Ardila and Postnikov prove that it is enough to consider minimal α [AP15, Lemma 1], which may be regarded as an analogue of the statement that $\mathcal{G}(\mathcal{A}^\dagger)$ is generated by the classes h_α for minimal α . Our deformation $\mathcal{Q}_-(\mathcal{A})$ of $\mathcal{P}_-(\mathcal{A})$ is defined in the most naive possible way:

$$\mathcal{Q}_-(\mathcal{A}) := \left\{ f \in \text{Sym } V^* \mid \alpha^{m(\alpha)-1} \cdot f = 0 \text{ for all } 0 \neq \alpha \in L \right\} \subset \text{Sym } V^*.$$

The inclusion of L^\perp into V^* makes $\mathcal{Q}_-(\mathcal{A})$ a graded module over $\text{Sym } L^\perp$.

Lemma 2.3. *The space $\mathcal{Q}_-(\mathcal{A})$ is a free module over $\text{Sym } L^\perp$, and we have a canonical isomorphism of graded vector spaces*

$$\mathcal{P}_-(\mathcal{A}) \cong \mathcal{Q}_-(\mathcal{A}) \otimes_{\text{Sym } L^\perp} \mathbb{F}.$$

Proof. Any splitting of the inclusion of L^\perp into V^* induces a graded $\text{Sym } L^\perp$ -module isomorphism $\mathcal{Q}_-(\mathcal{A}) \cong \mathcal{P}_-(\mathcal{A}) \otimes_{\text{Sym } L^\perp} \mathbb{F}$. While this isomorphism depends on the choice of splitting, the induced isomorphism $\mathcal{P}_-(\mathcal{A}) \cong \mathcal{Q}_-(\mathcal{A}) \otimes_{\text{Sym } L^\perp} \mathbb{F}$ does not. \square

3 Sheaves on the poset of flats

Let \mathcal{L} be the poset of flats of the matroid $M_{\mathcal{A}^\dagger}$. We will assume throughout this section that $M_{\mathcal{A}^\dagger}$ is loopless, so the minimal element of \mathcal{L} is \emptyset and the maximal element is E . Consider the topology on the set \mathcal{L} in which $U \subset \mathcal{L}$ is open if and only if it is downward-closed: that is, $F \in U$ whenever $F \leq G \in U$. The minimal open neighborhood of G is the set $U_G := \{F \mid F \leq G\}$. We can also think of \mathcal{L} as the set of objects of a thin category in which $|\text{Hom}(G, F)| = 1$ if $F \leq G$ and 0 otherwise. Given a sheaf \mathcal{F} on the topological space \mathcal{L} , we can produce a functor on the category \mathcal{L} by sending a flat G to the stalk $\mathcal{F}_G = \mathcal{F}(U_G)$. This process is reversible, so sheaves are in canonical bijection with functors [BP09, Proposition 1.1].

For any flat F , recall that we have defined the localization $(\mathcal{A}^!)_F = (F, V_F^*, (L^\perp)_F)$ in Section 1.5. For ease of notation, we will write $\mathcal{A}_F^!$ for $(\mathcal{A}^!)_F$ and L_F^\perp for $(L^\perp)_F = (L \cap V_F)^\perp$, but we stress that the Gale duality comes first and the localization second. We define a sheaf \mathcal{O} on by putting

$$\mathcal{O}_F := \text{Sym } L_F^\perp$$

and taking $\mathcal{O}_G \rightarrow \mathcal{O}_F$ to be the canonical projection for any $F \leq G$. In particular, we have $\mathcal{O}_E = \text{Sym } L^\perp$ and $\mathcal{O}_\emptyset = \mathbb{F}$. We call \mathcal{O} the **structure sheaf** of \mathcal{L} , and we will be interested in sheaves of graded \mathcal{O} -modules.

3.1 Minimal extension sheaves

Our main technical tool in this paper will be the theory of minimal extension sheaves on \mathcal{L} , developed in [BP09].

Proposition 3.1. [BP09, Proposition 1.10 and Corollary 3.7] *There exists a sheaf \mathcal{F} of graded \mathcal{O} -modules on \mathcal{L} with the following properties:*

- For every $F \in \mathcal{L}$, \mathcal{F}_F is a free module over \mathcal{O}_F .
- The sheaf \mathcal{F} is flabby: for every open $U \subset \mathcal{L}$, the restriction $\mathcal{F}(\mathcal{L}) \rightarrow \mathcal{F}(U)$ is surjective.
- We have $\mathcal{F}_\emptyset \cong \mathcal{O}_\emptyset \cong \mathbb{F}$.

The sheaf \mathcal{F} is unique up to isomorphism. Furthermore, the only endomorphisms of \mathcal{F} are those given by scalar multiplication.

Remark 3.2. In [BP09], it is only asserted that all automorphisms of \mathcal{F} are given by scalar multiplication, rather than all endomorphisms. However, the proof of the stronger statement is identical; see [BP09, Lemma 1.16] and [BBFK02, Remark 1.18].

Any sheaf \mathcal{F} satisfying the conditions of Proposition 3.1 is called a **minimal extension sheaf**.

Corollary 3.3. *Suppose that \mathcal{F} and \mathcal{F}' are minimal extension sheaves. Any sheaf homomorphism $\varphi : \mathcal{F} \rightarrow \mathcal{F}'$ with the property that $\varphi_\emptyset : \mathcal{F}_\emptyset \rightarrow \mathcal{F}'_\emptyset$ is nonzero is necessarily an isomorphism.*

Proof. By Proposition 3.1, \mathcal{F} and \mathcal{F}' are isomorphic, and there is a unique isomorphism $\psi : \mathcal{F}' \rightarrow \mathcal{F}$ with the property that $\psi_\emptyset = \varphi_\emptyset^{-1}$. Then $\varphi \circ \psi$ is an endomorphism of \mathcal{F} and $\psi \circ \varphi$ is an endomorphism of \mathcal{F}' , both of which restrict to the identity at L^\perp . By the last sentence of Proposition 3.1, they are both equal to the identity, so $\varphi = \psi^{-1}$. \square

Remark 3.4. For a sheaf \mathcal{F} to be flabby, it is not enough for the stalk maps

$$\mathcal{F}_E = \mathcal{F}(\mathcal{L}) \rightarrow \mathcal{F}(U_F) = \mathcal{F}_F$$

to be surjective for all $F \in \mathcal{L}$. Rather, the map $\mathcal{F}(\mathcal{L}) \rightarrow \mathcal{F}(U)$ has to be surjective for *every* open set U , including those that are not of the form U_F . For example, the structure sheaf \mathcal{O} is almost never flabby [BP09, Example 1.14], and is therefore almost never a minimal extension sheaf.

3.2 OT and $\text{SR}_{<}$

For any flat F of the matroid $M_{\mathcal{A}^!}$, we will be interested in the rings $\text{OT}(\mathcal{A}_F^!)$ and $\text{SR}_{<}(\mathcal{A}_F^!)$, both of which are algebras over $\mathcal{O}_F = \text{Sym } L_F^\perp$. Recall that the localization $\mathcal{A}_F^!$ is Gale dual to the contraction $\mathcal{A}^{E \setminus F} = (F, V_F, L \cap V_F)$, therefore we have

$$\text{OT}(\mathcal{A}_F^!) \cong \text{Sym } V_F^* / \mathcal{G}(\mathcal{A}_F^!) \quad \text{and} \quad \text{SR}_{<}(\mathcal{A}_F^!) \cong \text{Sym } V_F^* / \mathcal{G}_{<}(\mathcal{A}_F^!),$$

where the generators of $\mathcal{G}(\mathcal{A}_F^!)$ are indexed by minimal elements $\alpha \in L \cap V_F$. These are the same as minimal elements $\alpha \in L$ with $\text{Supp}(\alpha) \subset F$.

Given a pair of flats $F \leq G$, we have a homomorphism

$$\rho_{FG} : \text{Sym } V_G^* \rightarrow \text{Sym } V_F^*$$

induced by the projection of V_G^* onto V_F^* . By definition, ρ_{FG} restricts to a homomorphism from $\text{Sym } L_G^\perp \subset \text{Sym } V_G^*$ to $\text{Sym } L_F^\perp \subset \text{Sym } V_F^*$.

Lemma 3.5. *We have*

$$\rho_{FG}(\mathcal{G}(\mathcal{A}_G^!)) \subset \mathcal{G}(\mathcal{A}_F^!) \quad \text{and} \quad \rho_{FG}(\mathcal{G}_{<}(\mathcal{A}_G^!)) \subset \mathcal{G}_{<}(\mathcal{A}_F^!),$$

thus ρ_{FG} descends to graded algebra homomorphisms

$$\text{OT}(\mathcal{A}_G^!) \rightarrow \text{OT}(\mathcal{A}_F^!) \quad \text{and} \quad \text{SR}_{<}(\mathcal{A}_G^!) \rightarrow \text{SR}_{<}(\mathcal{A}_F^!).$$

Proof. Suppose that $\alpha \in L$ is minimal and $\text{Supp}(\alpha) \subset G$. If $\text{Supp}(\alpha) \subset F$, then

$$\rho_{FG}(h_\alpha) = h_\alpha \quad \text{and} \quad \rho_{FG}(\text{in}_{<}(h_\alpha)) = \text{in}_{<}(h_\alpha).$$

Otherwise, the fact that $\text{Supp}(\alpha)$ is a circuit of $M_{\mathcal{A}^!}$ and F is a flat of $M_{\mathcal{A}^!}$ not containing $\text{Supp}(\alpha)$ implies that F is missing at least two elements of $\text{Supp}(\alpha)$, and therefore we can conclude that $\rho_{F,E}(h_\alpha) = 0 = \rho_{FG}(\text{in}_{<}(h_\alpha))$. \square

By Lemma 3.5, we may define sheaves of \mathcal{O} -modules OT and $\text{SR}_{<}$ on \mathcal{L} whose stalks at F are $\text{OT}(\mathcal{A}_F^!)$ and $\text{SR}_{<}(\mathcal{A}_F^!)$, with restriction maps given by the homomorphisms ρ_{FG} .

Proposition 3.6. [BP09, Propositions 3.9 and 3.11] *The sheaves OT and $\text{SR}_{<}$ are both minimal extension sheaves.*

Remark 3.7. Propositions 3.1 and 3.6 have the surprising consequence that there is a unique isomorphism of \mathcal{O} -modules $\text{OT} \cong \text{SR}_{<}$ that restricts to the identity at the flat \emptyset , and therefore a canonical isomorphism of graded $\text{Sym } L^\perp$ -modules

$$\text{OT}(\mathcal{A}^!) = \text{OT}_E \cong (\text{SR}_{<})_E = \text{SR}_{<}(\mathcal{A}^!).$$

This is not a ring isomorphism, and it is not easy to give an explicit formula for it.

3.3 \mathcal{Q}_-

In this section, we will be interested in the graded $\text{Sym } L_F^\perp$ -module

$$\mathcal{Q}_-(\mathcal{A}^{E \setminus F}) = \left\{ f \in \text{Sym } V_F^* \mid \alpha^{m(\alpha)-1} \cdot f = 0 \text{ for all } 0 \neq \alpha \in L \cap V_F \right\} \subset \text{Sym } V_F^*.$$

Lemma 3.8. *If $F \leq G$ are flats, then $\rho_{FG} : \text{Sym } V_G^* \rightarrow \text{Sym } V_F^*$ takes $\mathcal{Q}_-(\mathcal{A}^{E \setminus G})$ to $\mathcal{Q}_-(\mathcal{A}^{E \setminus F})$.*

Proof. If $f \in \mathcal{Q}_-(\mathcal{A}^{E \setminus G})$ and $0 \neq \alpha \in L \cap V_F$, then

$$\alpha^{m(\alpha)-1} \cdot \rho_{FG}(f) = \rho_{FG}(\alpha^{m(\alpha)-1} \cdot f) = \rho_{FG}(0) = 0.$$

This completes the proof. □

By Lemma 3.8, we may define a sheaf \mathcal{Q}_- on \mathcal{L} whose stalk at F is equal to $\mathcal{Q}_-(\mathcal{A}^{E \setminus F})$, with restriction maps given by the homomorphisms ρ_{FG} . We would like to show that \mathcal{Q}_- is a minimal extension sheaf. The fact that the stalks are free modules is proved in Lemma 2.3 and the fact that the stalk at the empty flat is isomorphic to \mathbb{F} is trivial. However, the flabbiness property is not obvious. While it is easy to show that the map $\rho_{FE} : \mathcal{Q}_-(\mathcal{A}) \rightarrow \mathcal{Q}_-(\mathcal{A}^{E \setminus F})$ is surjective, this is not enough (see Remark 3.4). This issue will be resolved in the next section by making use of a result from [HR11].

3.4 Isomorphisms

The inclusion of $\mathcal{Q}_-(\mathcal{A}^{E \setminus F})$ into $\text{Sym } V_F^*$, followed by the projection of $\text{Sym } V_F^*$ onto $\text{OT}(\mathcal{A}_F^!)$ or $\text{SR}_<(\mathcal{A}_F^!)$, define graded $\text{Sym } L_F^\perp$ -module homomorphisms

$$\varphi_F : \mathcal{Q}_-(\mathcal{A}^{E \setminus F}) \rightarrow \text{OT}(\mathcal{A}_F^!) \quad \text{and} \quad (\varphi_<)_F : \mathcal{Q}_-(\mathcal{A}^{E \setminus F}) \rightarrow \text{SR}_<(\mathcal{A}_F^!).$$

These morphisms are all compatible with the maps ρ_{FG} , and therefore they define sheaf homomorphisms

$$\varphi : \mathcal{Q}_- \rightarrow \text{OT} \quad \text{and} \quad \varphi_< : \mathcal{Q}_- \rightarrow \text{SR}_<.$$

Proposition 3.9. *The homomorphism $\varphi_< : \mathcal{Q}_- \rightarrow \text{SR}_<$ is an isomorphism.*

Proof. Fix a flat F , and consider the graded $\text{Sym } L_F^\perp$ -module homomorphism

$$(\varphi_<)_F : \mathcal{Q}_-(\mathcal{A}^{E \setminus F}) \rightarrow \text{SR}_<(\mathcal{A}_F^!).$$

Since the source and the target are both free modules, it is sufficient to prove that it becomes an isomorphism after tensoring over $\text{Sym } L_F^\perp$ with \mathbb{F} . This yields the map

$$\mathcal{P}_-(\mathcal{A}^{E \setminus F}) \rightarrow \overline{\text{SR}}_<(\mathcal{A}_F^!),$$

which is an isomorphism by [HR11, Theorem 5.9(6)]. \square

Corollary 3.10. *The sheaf \mathcal{Q}_- is a minimal extension sheaf, and $\varphi : \mathcal{Q}_- \rightarrow \text{OT}$ is an isomorphism.*

Proof. The first statement follows from Propositions 3.6 and 3.9, and the second follows from the first statement, Proposition 3.6, and Corollary 4.4. \square

Proof of Theorem 1.8. If $M_{\mathcal{A}^\dagger}$ has a loop e , then there exists $\alpha \in L$ with $\text{Supp}(\alpha) = \{e\}$. It follows that $\mathcal{P}_-(\mathcal{A}) = 0$ and $\mathcal{K}(\mathcal{A}^\dagger) = \text{Sym } L^*$, and therefore the theorem holds. Thus we may assume that $M_{\mathcal{A}^\dagger}$ is loopless, which allows us to apply the machinery of Section 3.

By Corollary 3.10, the map $\varphi_E : \mathcal{Q}_-(\mathcal{A}) \rightarrow \text{OT}(\mathcal{A}^\dagger)$ is an isomorphism. Tensoring over $\text{Sym } L^\perp$ with \mathbb{F} , we see the the composition $\mathcal{P}_-(\mathcal{A}) \hookrightarrow \text{Sym } L^* \rightarrow \overline{\text{OT}}(\mathcal{A}^\dagger)$ is an isomorphism. \square

3.5 An example

We conclude this section with an example of Theorem 1.8. We have intentionally chosen an example that has proved to be deceptive in the past. The space $\mathcal{P}_-(\mathcal{A})$ includes all of the monomials $\chi_S = \prod_{e \in S} \chi(e) \in \text{Sym } L^*$ for those subsets S such that every circuit of $M_{\mathcal{A}^\dagger}$ contains at least two elements of $E \setminus S$, and Holtz and Ron conjectured that $\mathcal{P}_-(\mathcal{A})$ is spanned by such monomials [HR11, Conjecture 6.1]. A proof of this conjecture was published in [AP10] and later retracted in [AP15], along with a counterexample. The example that we consider here is precisely the counterexample appearing in [AP15]. One can regard this as the first example for which the space $\mathcal{P}_-(\mathcal{A})$ is trickier than one might expect.

Let Γ be the graph obtained from the complete graph on three vertices by doubling each edge, and let

$$\mathcal{A} := \mathcal{A}_\Gamma^\dagger = (E, C_1(\Gamma; \mathbb{Q}), H_1(\Gamma; \mathbb{Q}))$$

be the corresponding cographical linear space (see Section 1.3). We will label the edges with the labels e_1, e_2, f_1, f_2, g_1 , and g_2 , so that $\{e_1, e_2\}$, $\{f_1, f_2\}$ and $\{g_1, g_2\}$ are the three parallel pairs. We will orient them so that each parallel pair is oppositely oriented and $\{e_1, f_1, g_1\}$ forms an oriented cycle. Then $L = H_1(\Gamma; \mathbb{Q})$ is spanned by the classes

$$\ell_e := v_{e_1} + v_{e_2}, \quad \ell_f := v_{f_1} + v_{f_2}, \quad \ell_g := v_{g_1} + v_{g_2}, \quad \text{and} \quad \ell := v_{e_1} + v_{f_1} + v_{g_1}.$$

The defining ideal

$$\left\langle \alpha^{m(\alpha)-1} \mid 0 \neq \alpha \in H_1(\Gamma; \mathbb{Q}) \right\rangle$$

of the internal zonotopal algebra turns out to be generated by the elements ℓ_e, ℓ_f, ℓ_g , and ℓ^2 , thus $\mathcal{R}_-(\mathcal{A}) \cong \mathbb{Q}[\ell]/\ell^2$. We therefore have $\mathcal{P}_-(\mathcal{A}) = \mathbb{Q}\{1, c\}$, where $c \in L^* = H^1(\Gamma; \mathbb{Q})$ is the class that pairs trivially with ℓ_e, ℓ_f , and ℓ_g , and pairs to 1 with ℓ . We have a surjection from the space $V^* = C^1(\Gamma; \mathbb{Q}) = \mathbb{Q}\{w_{e_1}, w_{e_2}, w_{f_1}, w_{f_2}, w_{g_1}, w_{g_2}\}$ to $H^1(\Gamma; \mathbb{Q}) = L^*$, but c is not the image of any basis element. Rather, c is the image of the element $(w_{e_1} - w_{e_2} + w_{f_1} - w_{f_2} + w_{g_1} - w_{g_2})/3$.

On the dual side, we have the Orlik–Terao algebra

$$\mathrm{OT}(\mathcal{A}^\dagger) \cong \mathbb{Q}[w_{e_1}, w_{e_2}, w_{f_1}, w_{f_2}, w_{g_1}, w_{g_2}] / \langle w_{e_1} + w_{e_2}, w_{f_1} + w_{f_2}, w_{g_1} + w_{g_2}, w_{e_1}w_{f_1} + w_{e_1}w_{g_1} + w_{f_1}w_{g_1} \rangle.$$

The linear space $L^\perp \subset V^*$ is generated by the classes $w_{e_1} - w_{e_2} - w_{f_1} + w_{f_2}$ and $w_{f_1} - w_{f_2} - w_{g_1} + w_{g_2}$, thus we have the reduced Orlik–Terao algebra

$$\overline{\mathrm{OT}}(\mathcal{A}^\dagger) \cong \mathrm{OT}(\mathcal{A}) / \langle w_{e_1} - w_{e_2} - w_{f_1} + w_{f_2}, w_{f_1} - w_{f_2} - w_{g_1} + w_{g_2} \rangle \cong \mathbb{Q}[w] / \langle w^2 \rangle,$$

where $w := w_{e_1} = w_{f_1} = w_{g_1} = -w_{e_2} = -w_{f_2} = -w_{g_2}$. The isomorphism $\mathcal{P}_-(\mathcal{A}) \cong \overline{\mathrm{OT}}(\mathcal{A}^\dagger)$ of Theorem 1.8 coincides with the map

$$\mathcal{P}_-(\mathcal{A}) = \mathbb{Q}\{1, c\} \rightarrow \mathbb{Q}[w] / \langle w^2 \rangle \cong \overline{\mathrm{OT}}(\mathcal{A}^\dagger)$$

that sends c to $2w$.

4 Schubert varieties

The goal of this section is to prove Theorems 1.14 and 1.15.

4.1 The local geometry of $Y_{\mathcal{A}}$

Recall that, by definition, $Y_{\mathcal{A}}$ is a subvariety of $\mathbb{P}_{\mathcal{A}} = \prod_{e \in E} \overline{V}_e$. Because $Y_{\mathcal{A}}$ is the closure of a linear space, the \mathbb{N}^E -graded multidegree of $Y_{\mathcal{A}}$ is multiplicity free [AB16, Li18]. The following statement is a consequence of the main theorem in [Bri03, Theorem 1] on multiplicity free subvarieties of flag varieties; see the proof of [BF22, Theorem 4.3] for a discussion of rational singularities.

Theorem 4.1. *The Schubert variety $Y_{\mathcal{A}}$ has rational singularities, and is therefore normal and Cohen–Macaulay.*

The additive group V acts on $\mathbb{P}_{\mathcal{A}}$, and the additive subgroup $L \subset V$ acts on $Y_{\mathcal{A}}$. For any subset $F \subset E$, let $p_F \in \mathbb{P}_{\mathcal{A}}$ be the point defined by putting

$$(p_F)_e := \begin{cases} 0 & \text{if } e \in F \\ \infty & \text{if } e \notin F. \end{cases}$$

Lemma 4.2. [Pro18, Proposition 4.10] *The following statements hold.*

- We have $p_F \in Y_{\mathcal{A}}$ if and only if F is a flat of $M_{\mathcal{A}}$.¹³
- The stabilizer in L of p_F is $L \cap V_{E \setminus F}$.

¹³Warning: All of the flats in Section 3 were flats of $M_{\mathcal{A}^\dagger}$, whereas now we are considering flats of $M_{\mathcal{A}}$.

- Every element of $Y_{\mathcal{A}}$ lies in the orbit of a unique point p_F . That is, we have

$$Y_{\mathcal{A}} = \bigsqcup_{F \text{ a flat}} L \cdot p_F \cong \bigsqcup_{F \text{ a flat}} L/(L \cap V_F) \cong \bigsqcup_{F \text{ a flat}} L_F.$$

- Given two flats F and G , L_F is contained in the closure of L_G if and only if $F \leq G$.

Lemma 4.2 describes a stratification of $Y_{\mathcal{A}}$. Our next goal is to describe $Y_{\mathcal{A}}$ in the neighborhood of a stratum. For any flat F , we have a closed embedding $\iota_F : Y_{\mathcal{A}^F} \rightarrow Y_{\mathcal{A}}$ given by the formula

$$\iota_F(p)_e := \begin{cases} 0 & \text{if } e \in F \\ p_e & \text{if } e \notin F. \end{cases}$$

Let

$$U_F := \bigsqcup_{F \leq G} L_G = Y_{\mathcal{A}} \setminus \bigcup_{F \not\leq G} \overline{L_G}.$$

The second description shows that U_F is an open subscheme of $Y_{\mathcal{A}}$.

Choose an arbitrary section $s : L_F \rightarrow L$ of the projection of L onto L_F . For any flat $G \geq F$, composing s with the projection of L onto L_G gives a splitting $s_G : L_F \rightarrow L_G$ of the projection of L_G onto L_F .

Lemma 4.3. *We have an isomorphism $\varphi : L_F \times Y_{\mathcal{A}^F} \rightarrow U_F$ given by the formula*

$$\varphi(\alpha, p) := s(\alpha) \cdot \iota_F(p).$$

Proof. The poset of flats of $M_{\mathcal{A}^F}$ is isomorphic to the interval $[F, E]$ in the poset of flats of $M_{\mathcal{A}}$, so both the source and target have stratifications indexed by the same posets. In the source, the stratum corresponding to a flat $G \geq F$ is isomorphic to $L_F \times L_G/L_F$, and in the target it is isomorphic to L_G . The map φ is compatible with these stratifications, and the restriction of φ to the stratum $L_F \times L_G/L_F$ coincides with the isomorphism induced by the splitting s_G . Since φ restricts to a stratum-by-stratum isomorphism, it is a bijection. The variety $Y_{\mathcal{A}}$ is normal by Theorem 4.1, therefore so is U_F , thus Zariski's main theorem implies that φ is an isomorphism. \square

Proof of Theorem 1.14. Consider the special case of Lemma 4.3 where F is a flat of corank 1. In this case, the subspace $L^F := L \cap V_{E \setminus F} \subset L$ is 1-dimensional, we have $Y_{\mathcal{A}^F} \cong \overline{L^F} \cong \mathbb{P}^1$, and for any $\alpha \in L_F$, $\varphi(\alpha, L^F) = s(\alpha) + L^F$ is an affine translate of L^F .

Let f be a rational function on $Y_{\mathcal{A}}$ whose restriction to L is regular. Then

$$\text{ord}_{D_F} f = \text{ord}_{L_F}(f|_{U_F}) = \text{ord}_{L_F \times \{\infty\}} \varphi^*(f|_{U_F}),$$

which is equal to the degree of f when restricted to a generic affine translate of the line $L^F \subset L$. It follows that $H^0(\mathcal{O}_{Y_{\mathcal{A}}}(D_k))$ is equal to the set of polynomials $f \in \text{Sym } L^*$ such that, for all corank

1 flats F , the restriction of f to any affine translate of L^F has degree at most $m(F) + k$. This is equal to $C_{\mathcal{A},k}$ by [AP10, Proposition 2.6] and [AP15, Lemma 1]. \square

4.2 The dualizing sheaf

Let

$$U := \bigcup_{\text{crk } F=1} U_F \subset Y_{\mathcal{A}}$$

be the union of the dense stratum and all strata of codimension 1. Let $\omega_{Y_{\mathcal{A}}}$ be the pushforward of the canonical sheaf ω_U , which is Weil divisorial and is a dualizing sheaf because $Y_{\mathcal{A}}$ is normal and Cohen–Macaulay [CLS11, Theorems 8.0.4 and 9.0.9]. Let

$$K_{\mathcal{A}} := -2 \sum_{\text{crk } F=1} D_F.$$

Proposition 4.4. *There is a canonical isomorphism $\omega_{Y_{\mathcal{A}}} \cong \det(L^*) \otimes \mathcal{O}_{Y_{\mathcal{A}}}(K_{\mathcal{A}})$.*

Proof. Let $\Omega \in \det(L^*)$ be a nonvanishing volume form on L , which is unique up to scale, and which we can regard as a rational volume form on $Y_{\mathcal{A}}$. Then

$$\omega_{Y_{\mathcal{A}}} \cong \det(L^*) \otimes \mathcal{O}_{Y_{\mathcal{A}}}(D),$$

where D is the divisor of poles of Ω . It remains only to show that $D = K_{\mathcal{A}}$.

Since D is preserved by the action of L , we have $D = \sum_F c_F D_F$ for some collection of integers $\{c_F\}$. Fix a particular corank 1 flat F . By Lemma 4.3, there exists an isomorphism

$$U_F \cong L_F \times Y_{\mathcal{A}^F} \cong L_F \times \mathbb{P}^1,$$

where the inclusion $L \subset U_F \cong L_F \times \mathbb{P}^1$ is a linear isomorphism onto $L_F \times \mathbb{A}^1$. Therefore c_F is equal to the order of the volume form dz at $\infty \in \mathbb{P}^1$, which is equal to -2 . \square

4.3 Equivariant minimal free resolution

Before proving Theorem 1.15, we explain why it is possible to choose a minimal free resolution of $\mathcal{O}_{Y_{\mathcal{A}}}$ that is equivariant with respect to the action of $\text{Aut}(\mathcal{A})$. Since $\text{Aut}(\mathcal{A})$ is an extension of a finite group by a torus, this reduces to a statement about finite group equivariant multigraded minimal free resolutions of modules over rings. Let

$$R = \bigoplus_{\mathbf{a} \in \mathbb{N}^k} R_{\mathbf{a}}$$

be a multigraded \mathbb{F} -algebra. Let Γ be a finite group acting on both \mathbb{N}^k and R , with the property that $\gamma \cdot R_{\mathbf{a}} \subset R_{\gamma(\mathbf{a})}$ for all $\gamma \in \Gamma$ and $\mathbf{a} \in \mathbb{N}^k$. Let M be a finitely generated Γ -equivariant multigraded R -module.

Lemma 4.5. *The module M admits a minimal free resolution of the form*

$$\dots \rightarrow Q_2 \otimes R \rightarrow Q_1 \otimes R \rightarrow Q_0 \otimes R \rightarrow M \rightarrow 0,$$

where each Q_i is a Γ -representation and the maps are Γ -equivariant.

Proof. Consider the surjection $M \rightarrow M \otimes_R \mathbb{F}$ of multigraded representations of Γ . By averaging, we can construct a Γ -equivariant section $s_0 : M \otimes_R \mathbb{F} \rightarrow M$. Let $Q_0 := M \otimes_R \mathbb{F}$, and consider the map

$$d_0 := s_0 \otimes \text{id} : Q_0 \otimes R \rightarrow M.$$

Let $K_0 := \ker d_0$, and let s_1 be a Γ -equivariant section of the surjection $K_0 \rightarrow K_0 \otimes_R \mathbb{F}$. Now let $Q_1 := K_0 \otimes_R \mathbb{F}$, and define $d_1 := s_1 \otimes \text{id}$. Continuing this process produces the desired resolution. \square

We will apply Lemma 4.5 to

$$R = \bigotimes_{e \in E} \text{Sym}(V_e \oplus \mathbb{F})$$

with $k = |E| + c$, where $c = \dim \text{Aut}(\mathcal{A})$ is equal to the number of connected components of the matroid $M_{\mathcal{A}}$, and Γ is equal to the group of connected components of $\text{Aut}(\mathcal{A})$. This gives us the following result.

Corollary 4.6. *Any $\text{Aut}(\mathcal{A})$ -equivariant coherent sheaf on $\mathbb{P}_{\mathcal{A}}$ admits a minimal free resolution by $\text{Aut}(\mathcal{A})$ -equivariant sheaves with $\text{Aut}(\mathcal{A})$ -equivariant differentials.*

4.4 Identifying the syzygies

For each element $e \in E$, define the divisor $D_0(e) := \sum_{e \notin F} D_F$, so that $\pi_e^* \mathcal{O}_{\overline{V}_e}(1) \cong \mathcal{O}_{Y_{\mathcal{A}}}(D_0(e))$. For each subset $S \subset E$, define $D_0(S) := \sum_{e \in S} D_0(e)$, so that

$$\mathcal{O}_{Y_{\mathcal{A}}}(1_S) := \bigotimes_{e \in S} \pi_e^* \mathcal{O}_{\overline{V}_e}(1) \cong \mathcal{O}_{Y_{\mathcal{A}}}(D_0(S)).$$

Remark 4.7. Let $\text{Aut}(\mathcal{A})_S \subset \text{Aut}(\mathcal{A})$ be the stabilizer of the subset $S \subset E$. The natural embedding of $\mathcal{O}_{Y_{\mathcal{A}}}(D_0(S))$ into the sheaf of rational functions on $Y_{\mathcal{A}}$ induces an $\text{Aut}(\mathcal{A})_S$ -linearization of $\mathcal{O}_{Y_{\mathcal{A}}}(1_S)$. In particular when $S = E$, it induces an $\text{Aut}(\mathcal{A})$ -linearization of $\mathcal{O}_{Y_{\mathcal{A}}}(1_E)$. It also induces an $\text{Aut}(\mathcal{A})$ -linearization of the right-hand side of the equation in Theorem 1.15.

Let

$$\dots \rightarrow \mathcal{Q}_2 \rightarrow \mathcal{Q}_1 \rightarrow \mathcal{Q}_0 \rightarrow \mathcal{O}_{Y_{\mathcal{A}}} \rightarrow 0$$

be an $\text{Aut}(\mathcal{A})$ -equivariant minimal free resolution of the structure sheaf $\mathcal{O}_{Y_{\mathcal{A}}}$ on $\mathbb{P}_{\mathcal{A}}$, which exists by Corollary 4.6. As in the statement of Theorem 1.15, let $M = M_{\mathcal{A}'}$. It follows from [AB16, Theorem 1.5] that

$$\mathcal{Q}_i \cong \bigoplus_{\text{crk}_M(E \setminus S) = i} \mathcal{Q}_S \otimes \mathcal{O}_{\mathbb{P}_{\mathcal{A}}}(-1_S),$$

where Q_S is a representation of $\text{Aut}(\mathcal{A})_S$.

Lemma 4.8. *For any subset $S \subset E$, there is a canonical isomorphism*

$$Q_S \otimes H^{|S|}(\mathbb{P}_{\mathcal{A}_S}, \mathcal{O}(-2_S)) \cong H^{\text{rk}_{M_{\mathcal{A}}}(S)}(Y_{\mathcal{A}}, \mathcal{O}(-1_S)).$$

Proof. We begin by noting that the Künneth theorem provides an isomorphism

$$H^{|S|}(\mathbb{P}_{\mathcal{A}_S}, \mathcal{O}(-2_S)) \cong H^{|S|}(\mathbb{P}_{\mathcal{A}}, \mathcal{O}(-2_S)).$$

Now consider the resolution

$$0 \rightarrow \mathcal{Q}_{\text{rk } M}(-1_S) \rightarrow \cdots \rightarrow \mathcal{Q}_2(-1_S) \rightarrow \mathcal{Q}_1(-1_S) \rightarrow \mathcal{Q}_0(-1_S) \rightarrow \mathcal{O}_{Y_{\mathcal{A}}}(-1_S) \rightarrow 0$$

of $\mathcal{O}_{Y_{\mathcal{A}}}(-1_S)$. Since all cohomology groups of $\mathcal{O}_{\mathbb{P}^1}(-1)$ vanish, the Künneth theorem implies that the same is true of $\mathcal{O}_{\mathbb{P}_{\mathcal{A}}}(-1_S - 1_T)$ unless $S = T$. It follows that the cohomology of $\mathcal{Q}_i(-1_S)$ vanishes unless $i = \text{crk}_M(E \setminus S)$, in which case it is isomorphic to

$$Q_S \otimes H^{|S|}(\mathbb{P}_{\mathcal{A}}, \mathcal{O}_{\mathbb{P}_{\mathcal{A}}}(-2_S)).$$

If we regard $\mathcal{Q}_{\bullet}(-1_S)$ as a complex of sheaves with $\mathcal{Q}_i(-1_S)$ sitting in degree $-i$, this complex is quasi-isomorphic to $\mathcal{O}_{Y_{\mathcal{A}}}(-1_S)$, and we therefore have

$$Q_S \otimes H^{|S|}(\mathbb{P}_{\mathcal{A}}, \mathcal{O}_{\mathbb{P}_{\mathcal{A}}}(-2_S)) \cong \mathbb{H}^{|S| - \text{crk}_M(E \setminus S)}(\mathbb{P}_{\mathbb{A}}, \mathcal{Q}_{\bullet}(-1_S)) \cong H^{|S| - \text{crk}_M(E \setminus S)}(Y_{\mathcal{A}}, \mathcal{O}(-1_S)).$$

The lemma now follows from the fact that $|S| - \text{crk}_M = \text{rk}_{M^!}(S) = \text{rk}_{M_{\mathcal{A}}}(S)$. \square

Consider the projection map $\pi_S : \mathbb{P}_{\mathcal{A}} \rightarrow \mathbb{P}_{\mathcal{A}_S}$ along with its restriction $\pi_S : Y_{\mathcal{A}} \rightarrow Y_{\mathcal{A}_S}$.

Lemma 4.9. *The pullback*

$$\pi_S^* : H^{\text{rk}_{M_{\mathcal{A}}}(S)}(Y_{\mathcal{A}_S}, \mathcal{O}(-1_S)) \rightarrow H^{\text{rk}_{M_{\mathcal{A}}}(S)}(Y_{\mathcal{A}}, \mathcal{O}(-1_S))$$

is an isomorphism.

Proof. Choose $T \subset E \setminus S$ such that $|T| = \text{crk}_{M_{\mathcal{A}}}(S)$ and $S \cup T$ is a spanning set for $M_{\mathcal{A}}$. Then $\pi_S : Y_{\mathcal{A}} \rightarrow Y_{\mathcal{A}_S}$ can be factored as a birational proper morphism

$$\pi_{S \cup T} : Y_{\mathcal{A}} \rightarrow Y_{\mathcal{A}_{S \cup T}}$$

followed by a projection

$$\rho_{S,T} : Y_{\mathcal{A}_{S \cup T}} \cong Y_{\mathcal{A}_S} \times (\mathbb{P}^1)^T \rightarrow Y_{\mathcal{A}_S}.$$

Because Schubert varieties have rational singularities (Theorem 4.1), the induced map

$$\pi_{S \cup T}^* : H^{\text{rk}_{M_{\mathcal{A}}}(S)}(Y_{\mathcal{A}_{S \cup T}}, \mathcal{O}(-1_S)) \rightarrow H^{\text{rk}_{M_{\mathcal{A}}}(S)}(Y_{\mathcal{A}_S}, \mathcal{O}(-1_S))$$

is also an isomorphism. Thus the composition $\pi_S^* = \pi_{S \cup T}^* \circ \rho_{S \cup T}^*$ is an isomorphism. \square

Proof of Theorem 1.15. By Lemmas 4.8 and 4.9, we have

$$Q_S \otimes H^{|S|}(\mathbb{P}_{\mathcal{A}_S}, \mathcal{O}(-2_S)) \cong H^{\text{rk}_{M_{\mathcal{A}}(S)}}(Y_{\mathcal{A}_S}, \mathcal{O}(-1_S)). \quad (1)$$

By Proposition 4.4, we have

$$\omega_{\mathbb{P}_{\mathcal{A}_S}} \cong \det(V_S^*) \otimes \mathcal{O}(-2_S) \quad \text{and} \quad \omega_{Y_{\mathcal{A}_S}} \cong \det(L_S^*) \otimes \mathcal{O}(K_{\mathcal{A}_S}) \cong \det(L_S^*) \otimes \mathcal{O}(-1_S) \otimes \mathcal{O}(D_{-2}),$$

thus we can rewrite Equation (1) as

$$\det(V_S) \otimes Q_S \otimes H^{|S|}(\mathbb{P}_{\mathcal{A}_S}, \omega_{\mathbb{P}_{\mathcal{A}_S}}) \cong \det(L_S) \otimes H^{\text{rk}_{M_{\mathcal{A}}(S)}}(Y_{\mathcal{A}_S}, \omega_{Y_{\mathcal{A}_S}} \otimes \mathcal{O}(-D_{-2})).$$

By Serre duality, we have

$$\begin{aligned} \det(V_S) \otimes Q_S &\cong \det(V_S) \otimes Q_S \otimes H^{|S|}(\mathbb{P}_{\mathcal{A}_S}, \omega_{\mathbb{P}_{\mathcal{A}_S}}) \\ &\cong \det(L_S) \otimes H^{\text{rk}_{M_{\mathcal{A}}(S)}}(Y_{\mathcal{A}_S}, \omega_{Y_{\mathcal{A}_S}} \otimes \mathcal{O}(-D_{-2})) \\ &\cong \det(L_S) \otimes H^0(Y_{\mathcal{A}_S}, \mathcal{O}(D_{-2}))^*. \end{aligned}$$

The theorem now follows from the isomorphism $\det(L_S) \cong \det(V_S) \otimes \det(L_S^\perp)$. \square

Proof of Corollary 1.18. Consider the resolution

$$\cdots \rightarrow \mathcal{Q}_2(1_E) \rightarrow \mathcal{Q}_1(1_E) \rightarrow \mathcal{Q}_0(1_E) \rightarrow \mathcal{O}_{Y_{\mathcal{A}}}(1_E) \rightarrow 0$$

obtained from Theorem 1.15 by tensoring with $\mathcal{O}(1_E)$. By [Bri03], all of the sheaves $\mathcal{Q}_i(1_E)$ as well as $\mathcal{O}_{Y_{\mathcal{A}}}(1_E)$ have vanishing higher cohomology, so the hypercohomology spectral sequence tells us that their sections form an exact sequence. Finally, $H^0(\mathcal{O}_{Y_{\mathcal{A}}}(1_E)) \cong \mathcal{R}_+(A)^*$ by Theorem 1.14. \square

4.5 The complete graph

Let $\mathcal{A} := \mathcal{A}_{K_n}^\dagger$ be the cographical linear space associated with the complete graph K_n . Then $M = M_{\mathcal{A}} = M_{K_n}$ is the braid matroid, whose flats of corank i are indexed by partitions of the set $[n]$ into $i + 1$ parts. Given such a partition $\pi = (\pi_0, \dots, \pi_i)$, the associated flat F_π is the union of the complete graphs on the parts of π , whereas the complement $S_\pi := E \setminus F_\pi$ is the complete multipartite graph consisting of edges that go between the blocks. Thus we have

$$\begin{aligned} H^0(\mathcal{Q}_i(1_E)) &\cong \bigoplus_{\text{crk}_M E \setminus S = i} \det(L_{S^\perp}^\perp) \otimes \mathcal{R}_-(\mathcal{A}_S) \otimes \bigotimes_{e \notin S} (V_e^* \oplus \mathbb{Q}) \\ &\cong \bigoplus_{\substack{\pi = (\pi_0, \dots, \pi_i) \\ [n] = \pi_0 \sqcup \dots \sqcup \pi_i}} \det(L_{S_\pi}^\perp) \otimes \mathcal{R}_-(\mathcal{A}_{S_\pi}) \otimes \bigotimes_{e \in F_\pi} (V_e^* \oplus \mathbb{Q}). \end{aligned}$$

Let $\mathfrak{S}_\pi \subseteq \mathfrak{S}_n$ be the subgroup of permutations that preserves the set partition π . We have a homomorphism $\rho_\pi : \mathfrak{S}_\pi \rightarrow \mathfrak{S}_{i+1}$ that records the way in which an element \mathfrak{S}_π permutes the blocks of π , with kernel equal to the Young subgroup $\mathfrak{S}_{\pi_0} \times \cdots \times \mathfrak{S}_{\pi_i}$. Each of the tensor factors above is a representation of \mathfrak{S}_π , and we will analyze each one individually.

Let Γ_π be the graph obtained from K_n by contracting the edges in F_π ; this is a loopless graph with vertex set $\{0, \dots, i\}$ and $|\pi_j| \cdot |\pi_k|$ edges between distinct vertices j and k . Then

$$\mathcal{A}_{S_\pi} = (\mathcal{A}_{K_n}^!)_{S_\pi} = (\mathcal{A}_{K_n}^{F_\pi})^! \cong \mathcal{A}_{\Gamma_\pi}^!.$$

This implies that

$$L_{S_\pi} = H_1(\Gamma_\pi; \mathbb{Q}) \subset C_1(\Gamma_\pi; \mathbb{Q}),$$

and therefore

$$L_{S_\pi}^\perp = B^1(\Gamma_\pi; \mathbb{Q}) = \ker(C^1(\Gamma_\pi; \mathbb{Q}) \rightarrow H^1(\Gamma_\pi; \mathbb{Q})).$$

The space of coboundaries of a graph is canonically isomorphic to that of its simplification, thus

$$L_{S_\pi}^\perp = B^1(\Gamma_\pi; \mathbb{Q}) \cong B^1(K_{i+1}; \mathbb{Q}).$$

This is isomorphic to the irreducible representation $V_{[i,1]}$ of \mathfrak{S}_{i+1} , hence

$$\det(L_{S_\pi}^\perp) \cong \wedge^i(V_{[i,1]}) \cong \text{sign}_{\mathfrak{S}_{i+1}},$$

which we regard as a representation of \mathfrak{S}_π via the homomorphism $\rho_\pi : \mathfrak{S}_\pi \rightarrow \mathfrak{S}_{i+1}$. Since the multiplicative group $\mathbb{G}_m \subset \text{Aut}(\mathcal{A})$ acts on $L_{S_\pi}^\perp$ with weight 1, we will write

$$\det(L_{S_\pi}^\perp) \cong \text{sign}_{\mathfrak{S}_{i+1}}[-i]$$

to indicate that it is sitting in degree i .

Simplifying a loopless graph by deleting parallel edges does not affect the internal zonotopal algebra of the cographical linear space; this can be seen in many different ways, for example from [CDBHP, Theorem 1.9]. Thus we have \mathfrak{S}_π -equivariant isomorphisms

$$\mathcal{R}_-(\mathcal{A}_{S_\pi}) \cong \mathcal{R}_-(\mathcal{A}_{\Gamma_\pi}^!) \cong \mathcal{R}_-(\mathcal{A}_{K_{i+1}}^!) \cong \overline{\text{OT}}(\mathcal{A}_{K_{i+1}})^*,$$

where the last isomorphism follows from Theorem 1.8. Note that the reduced Orlik–Terao algebra is generated in degree 1, and dualizing negates degrees.

Remark 4.10. The graded \mathfrak{S}_{i+1} -representation $\overline{\text{OT}}(\mathcal{A}_{K_{i+1}})$ is studied in [MPY17], where in particular one can find an explicit algorithm to compute it recursively. Alternatively, $\mathcal{R}_-(\mathcal{A}_{K_{i+1}}^!)$ is isomorphic in degree $-j$ to $H^{2j}(X(SU(2), K_{i+1}); \mathbb{Q})$ [Ber18, CDBHP], which is Whitehouse’s lift of an Eulerian representation of \mathfrak{S}_i , tensored with $\text{sign}_{\mathfrak{S}_{i+1}}$ [ER19, Proposition 2].

Finally, for any subgraph Γ of K_n , let

$$\Theta_\Gamma := \bigotimes_{e \in E(\Gamma)} V_e^*;$$

this is a (possibly nontrivial) 1-dimensional representation of $\text{Aut}(\Gamma)$ sitting in degree $|E(\Gamma)|$. Then

$$\bigotimes_{e \in F_\pi} (V_e^* \oplus \mathbb{Q}) \cong \bigoplus_{\Gamma \in \mathcal{F}_\pi} \Theta_\Gamma,$$

where \mathcal{F}_π is the set of all subgraphs of F_π . Putting it all together, we have

$$H^0(\mathcal{Q}_i(1_E)) \cong \bigoplus_{\substack{\pi=(\pi_0, \dots, \pi_i) \\ [n]=\pi_0 \sqcup \dots \sqcup \pi_i}} \text{sign}_{\mathfrak{S}_{i+1}}[i] \otimes \overline{\text{OT}}(\mathcal{A}_{K_{i+1}})^* \otimes \bigoplus_{\Gamma \in \mathcal{F}_\pi} \Theta_\Gamma,$$

where the first two tensor factors are regarded as representations of \mathfrak{S}_π via the homomorphism ρ_π .

Corollary 4.11. *We have an equality*

$$\begin{aligned} \mathcal{R}_+(\mathcal{A}_{K_n}^!)^* &= \sum_{i=0}^n (-1)^i \bigoplus_{\substack{\pi=(\pi_0, \dots, \pi_i) \\ [n]=\pi_0 \sqcup \dots \sqcup \pi_i}} \text{sign}_{\mathfrak{S}_{i+1}}[i] \otimes \overline{\text{OT}}(\mathcal{A}_{K_{i+1}})^* \otimes \bigoplus_{\Gamma \in \mathcal{F}_\pi} \Theta_\Gamma \\ &= \bigoplus_{\Gamma} \Theta_\Gamma \otimes \bigoplus_{\pi: \Gamma \in \mathcal{F}_\pi} (-1)^{|\pi|-1} \text{sign}_{\mathfrak{S}_{|\pi|}}[|\pi|-1] \otimes \overline{\text{OT}}(\mathcal{A}_{K_{|\pi|}})^* \end{aligned}$$

of graded virtual representations of \mathfrak{S}_n .

Remark 4.12. The dimension of the left hand side of Corollary 4.11 is the number of connected graphs on $[n]$, and the dimension of $\overline{\text{OT}}(\mathcal{A}_{K_{i+1}})$ is the Möbius function value $\mu(\pi, \hat{1})$ of the partition lattice, where $\hat{1}$ is the single block partition. In this case, Remark 1.19 says that Corollary 4.11 categorifies the formula

$$\text{number of connected graphs on } [n] = \sum_{\pi=(\pi_0, \dots, \pi_i)} \mu(\pi, \hat{1}) \cdot 2^{\binom{|\pi_0|}{2} + \dots + \binom{|\pi_i|}{2}},$$

which is obtained via Möbius inversion from the formula expressing the total number of graphs on $[n]$ as the sum over $\pi = (\pi_0, \dots, \pi_i)$ of the number of graphs with connected components π .

Example 4.13. Consider the special case where $n = 3$. We will work with the first of the two expressions for $\mathcal{R}_+(\mathcal{A}_{K_3}^!)^*$ in Corollary 4.11.

- When $i = 0$, π is the partition of $[3]$ into a single block, $\text{sign}_{\mathfrak{S}_1}[0] \otimes \overline{\text{OT}}(\mathcal{A}_{K_1})^*$ is trivial, and \mathcal{F}_π consists of all simple graphs on the set $[3]$. The empty graph contributes the trivial representation $V_{[3]}$ in degree 0. The three graphs with one edge contribute the representation $V_{[2,1]} \oplus V_{[1,1,1]}$ in degree 1; the appearance of the sign representation is due to the fact that, if e is a single edge, then the stabilizer of e acts nontrivially on the line V_e^* . The three graphs

with two edges contribute the representation $V_{[2,1]} \oplus V_{[3]}$ in degree 2; the appearance of the trivial representation is due to the fact that, if e and f is a pair of edges, then the stabilizer of the set $\{e, f\}$ acts trivially on the line $V_e^* \otimes V_f^*$. Finally, the graph with all three edges contributes $\Theta_{K_3} \cong V_{[1,1,1]}$ in degree 3.

- When $i = 1$, π ranges over the three different partitions of $[3]$ into two blocks. For each such π , the homomorphism ρ_π is trivial (the two blocks have different sizes and therefore cannot be permuted), so $\text{sign}_{\mathfrak{S}_2}[-1] \otimes \overline{\text{OT}}(\mathcal{A}_{K_2})^* \cong \mathbb{Q}[-1]$. For each π , \mathcal{F}_π consists of two graphs, one of which is empty and one of which contains a single edge. The empty graph contributes the trivial representation of $\mathfrak{S}_\pi \cong S_2$ in degree 0; tensoring with $\mathbb{Q}[-1]$ and inducing to \mathfrak{S}_3 gives $V_{[3]} \oplus V_{[2,1]}$ in degree 1. The graph with one edge contributes the sign representation of $\mathfrak{S}_\pi \cong S_2$ in degree 1; tensoring with $\mathbb{Q}[-1]$ and inducing to \mathfrak{S}_3 gives $V_{[2,1]} \oplus V[1, 1, 1]$ in degree 2.
- When $i = 2$, π is the partition of $[3]$ into three blocks, $\text{sign}_{\mathfrak{S}_2}[2]$ is the sign representation in degree 2, and $\overline{\text{OT}}(\mathcal{A}_{K_3})^*$ is trivial in degree 0 and sign in degree -1 . The set \mathcal{F}_π contains only the empty graph, and Θ_\emptyset is trivial in degree zero. Thus the total contribution when $i = 2$ is $V_{[1,1,1]} \otimes V_{[1,1,1]} \cong V_{[3]}$ in degree 1 plus $V_{[1,1,1]} \otimes V_{[3]} \cong V_{[1,1,1]}$ in degree 2.

Taking the alternating sum, we find that $\mathcal{R}_+(\mathcal{A}_{K_3}^!)^*$ is isomorphic to the trivial representation of \mathfrak{S}_3 in degrees 0 and 2 plus the sign representation of \mathfrak{S}_3 in degrees 1 and 3. This can be confirmed by direct calculation, as the external zonotopal algebra of $\mathcal{A}_{K_3}^!$ is generated by a single element α in degree -1 that is not fixed by \mathfrak{S}_3 , modulo the relation $\alpha^4 = 0$.

References

- [AB16] Federico Ardila and Adam Boocher, *The closure of a linear space in a product of lines*, J. Algebraic Combin. **43** (2016), no. 1, 199–235.
- [AP10] Federico Ardila and Alexander Postnikov, *Combinatorics and geometry of power ideals*, Trans. Amer. Math. Soc. **362** (2010), no. 8, 4357–4384.
- [AP15] ———, *Correction to “Combinatorics and geometry of power ideals”: two counterexamples for power ideals of hyperplane arrangements [mr2608410]*, Trans. Amer. Math. Soc. **367** (2015), no. 5, 3759–3762.
- [AS88] A. A. Akopyan and A. A. Saakyan, *A system of differential equations that is related to the polynomial class of translates of a box spline*, Mat. Zametki **44** (1988), no. 6, 705–724, 861. MR 983544
- [BBFK02] Gottfried Barthel, Jean-Paul Brasselet, Karl-Heinz Fieseler, and Ludger Kaup, *Combinatorial intersection cohomology for fans*, Tohoku Math. J. (2) **54** (2002), no. 1, 1–41.

- [Ber10] Andrew Berget, *Products of linear forms and Tutte polynomials*, European J. Combin. **31** (2010), no. 7, 1924–1935.
- [Ber18] ———, *Internal zonotopal algebras and the monomial reflection groups $G(m, 1, n)$* , J. Combin. Theory Ser. A **159** (2018), 1–25.
- [BF22] Andrew Berget and Alex Fink, *Equivariant K -theory classes of matrix orbit closures*, Int. Math. Res. Not. IMRN (2022), no. 18, 14105–14133.
- [BP09] Tom Braden and Nicholas Proudfoot, *The hypertoric intersection cohomology ring*, Invent. Math. **177** (2009), no. 2, 337–379.
- [Bri03] Michel Brion, *Multiplicity-free subvarieties of flag varieties*, Commutative algebra (Grenoble/Lyon, 2001), Contemp. Math., vol. 331, Amer. Math. Soc., Providence, RI, 2003, pp. 13–23.
- [BV99] Michel Brion and Michèle Vergne, *Arrangement of hyperplanes. I. Rational functions and Jeffrey-Kirwan residue*, Ann. Sci. École Norm. Sup. (4) **32** (1999), no. 5, 715–741. MR 1710758
- [CDBHP] Colin Crowley, Galen Dorpalen-Barry, André Henriques, and Nicholas Proudfoot, *The geometry of zonotopal algebras i: cohomology of graphical configuration spaces*, arXiv:2502.12768.
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [dBH83] C. de Boor and K. Höllig, *B -splines from parallelepipeds*, J. Analyse Math. **42** (1982/83), 99–115.
- [DCP11] Corrado De Concini and Claudio Procesi, *Topics in hyperplane arrangements, polytopes and box-splines*, Universitext, Springer, New York, 2011.
- [DR90] N. Dyn and A. Ron, *Local approximation by certain spaces of exponential polynomials, approximation order of exponential box splines, and related interpolation problems*, Trans. Amer. Math. Soc. **319** (1990), no. 1, 381–403. MR 956032
- [EPW16] Ben Elias, Nicholas Proudfoot, and Max Wakefield, *The Kazhdan-Lusztig polynomial of a matroid*, Adv. Math. **299** (2016), 36–70.
- [ER19] Nicholas Early and Victor Reiner, *On configuration spaces and Whitehouse’s lifts of the Eulerian representations*, J. Pure Appl. Algebra **223** (2019), no. 10, 4524–4535. MR 3958103
- [HR11] Olga Holtz and Amos Ron, *Zonotopal algebra*, Adv. Math. **227** (2011), no. 2, 847–894.

- [HS02] Tamás Hausel and Bernd Sturmfels, *Toric hyperKähler varieties*, Doc. Math. **7** (2002), 495–534 (electronic).
- [HW17] June Huh and Botong Wang, *Enumeration of points, lines, planes, etc*, Acta Math. **218** (2017), no. 2, 297–317.
- [IKL⁺] Aleksei Ilin, Joel Kamnitzer, Yu Li, Piotr Przytycki, and Leonid Rybnikov, *Equivariant incidence algebras and equivariant Kazhdan–Lusztig–Stanley theory*, arXiv:2009.06696.
- [Jia93] Rong Qing Jia, *A dual basis for the integer translates of an exponential box spline*, Rocky Mountain J. Math. **23** (1993), no. 1, 223–242. MR 1212738
- [Jia24] Leo Jiang, *Real matroid Schubert varieties, zonotopes, and virtual Weyl groups*, Sémin. Lothar. Combin. **91B** (2024), Art. 71, 9.
- [KMY09] Allen Knutson, Ezra Miller, and Alexander Yong, *Gröbner geometry of vertex decompositions and of flagged tableaux*, J. Reine Angew. Math. **630** (2009), 1–31.
- [Len14] Matthias Lenz, *Interpolation, box splines, and lattice points in zonotopes*, Int. Math. Res. Not. IMRN (2014), no. 20, 5697–5712.
- [Li18] Binglin Li, *Images of rational maps of projective spaces*, Int. Math. Res. Not. IMRN (2018), no. 13, 4190–4228. MR 3829180
- [MPY17] Daniel Moseley, Nicholas Proudfoot, and Ben Young, *The Orlik-Terao algebra and the cohomology of configuration space*, Exp. Math. **26** (2017), no. 3, 373–380.
- [OT94] Peter Orlik and Hiroaki Terao, *Commutative algebras for arrangements*, Nagoya Math. J. **134** (1994), 65–73.
- [Pag23] Roberto Pagaria, *The Frobenius characteristic of the Orlik-Terao algebra of type A*, Int. Math. Res. Not. IMRN (2023), no. 13, 11577–11591.
- [Pro18] Nicholas Proudfoot, *The algebraic geometry of Kazhdan-Lusztig-Stanley polynomials*, EMS Surv. Math. Sci. **5** (2018), no. 1, 99–127.
- [PS06] Nicholas Proudfoot and David Speyer, *A broken circuit ring*, Beiträge Algebra Geom. **47** (2006), no. 1, 161–166.
- [PSS99] Alexander Postnikov, Boris Shapiro, and Mikhail Shapiro, *Algebras of curvature forms on homogeneous manifolds*, Differential topology, infinite-dimensional Lie algebras, and applications, Amer. Math. Soc. Transl. Ser. 2, vol. 194, Amer. Math. Soc., Providence, RI, 1999, pp. 227–235.
- [PXY18] Nicholas Proudfoot, Yuan Xu, and Ben Young, *The Z-polynomial of a matroid*, Electron. J. Combin. **25** (2018), no. 1, Paper 1.26, 21.

- [RRT23] Markus Reineke, Brendon Rhoades, and Vasu Tewari, *Zonotopal algebras, orbit harmonics, and Donaldson-Thomas invariants of symmetric quivers*, Int. Math. Res. Not. IMRN (2023), no. 23, 20169–20210.
- [Sch11] Hal Schenck, *Resonance varieties via blowups of \mathbb{P}^2 and scrolls*, Int. Math. Res. Not. IMRN (2011), no. 20, 4756–4778.
- [SSV13] Raman Sanyal, Bernd Sturmfels, and Cynthia Vinzant, *The entropic discriminant*, Adv. Math. **244** (2013), 678–707.
- [Ter02] Hiroaki Terao, *Algebras generated by reciprocals of linear forms*, J. Algebra **250** (2002), no. 2, 549–558.
- [Toh] Ștefan O. Tohăneanu, *Generalized Hamming weights and minimal shifts of Orlik–Terao algebras*, arXiv:2311.03688.